

Datasheet: MCA1972F

Description:	MOUSE ANTI PIG CD18a:FITC
Specificity:	CD18a
Other names:	INTEGRIN BETA 2 CHAIN
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	PNK-I
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species

Pig

Species Cross Reactivity

Reacts with: Camel

N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

Product Form

Ig Fraction conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid

Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525

Preparation

Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant

Buffer Solution

Phosphate buffered saline

Preservative Stabilisers	0.09% sodium azide (NaN ₃) 1% bovine serum albumin
Approx. Protein Concentrations	IgG concentration 0.1 mg/ml
Immunogen	Porcine large granular lymphocytes.
External Database Links	<p>UniProt: P53714 Related reagents</p> <p>Entrez Gene: 396943 ITGB2 Related reagents</p>
Synonyms	CD18
RRID	AB_2296305
Fusion Partners	Spleen cells from immunized Balb/c mice were fused with cells of the mouse P3-X63-Ag8.653 myeloma cell line.
Specificity	<p>Mouse anti Pig CD18a, clone PNK-I recognizes porcine CD18a. PNK-I was clustered as CD18a at the Second International Workshop to Define Swine Cluster of Differentiation (CD) Antigens (Saalmuller et al. 1998). Clone PNK-I immunoprecipitates proteins of ~166 kDa, ~155 kDa and ~95 kDa under non-reducing conditions, specifically recognizing the 95 kDa protein, consistent with the integrin β2 chain (CD18). PNK-I inhibits porcine NK cell activity independent of any effect on antibody dependent cellular cytotoxicity (Dato and Kim 1990).</p> <p>CD18 is a single pass type I transmembrane protein and is expressed on all leukocytes and is involved in a variety of cell functions. CD18 acts as a receptor for several ICAM molecules effecting intercellular adhesion functions, it is also involved in the recognition of a variety of extracellular substrate molecules.</p> <p>CD18 acts as a receptor for a number of leukotoxins produced by fungi and bacteria. Clone PNK-I is able to ameliorate the effects of these leukotoxins by blocking binding of the toxins to the CD18 receptor (Chen et al. 2011).</p>
Flow Cytometry	Use 10µl of the suggested working dilution to label 10 ⁶ cells in 100µl
References	<ol style="list-style-type: none"> 1. Dato, M.E. & Kim, Y.B. (1990) Characterization and utilization of a monoclonal antibody inhibiting porcine natural killer cell activity for isolation of natural killer and killer cells. J Immunol. 144 (11): 4452-62. 2. Haverson, K. et al. (1999) T-cell populations in the pig intestinal lamina propria: memory cells with unusual phenotypic characteristics. Immunology 96: 66-73. 3. Vanden Bergh, P.G. et al. (2008) Probing of Actinobacillus pleuropneumoniae ApxIIIA toxin-dependent cytotoxicity towards mammalian peripheral blood mononucleated cells.

[BMC Res Notes 1:121.](#)

4. Ebdrup, L. *et al.* (2008) Dynamic expression of the signal regulatory protein alpha and CD18 on porcine PBMC during acute endotoxaemia. [Scand J Immunol. 68:430-7.](#)
5. Vanden Bergh, P.G. *et al.* (2009) Porcine CD18 mediates Actinobacillus pleuropneumoniae ApxIII species-specific toxicity. [Vet Res. 40:1-10.](#)
6. Chen, Z.W. *et al.* (2011) Mechanisms underlying Actinobacillus pleuropneumoniae exotoxin ApxI induced expression of IL-1 β , IL-8 and TNF- α in porcine alveolar macrophages. [Vet Res. 42:25.](#)

Further Reading	1. Piriou-Guzylack, L. (2008) Membrane markers of the immune cells in swine: an update. Vet Res. 39:54.
Storage	<p>This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.</p> <p>Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.</p>
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/MCA1972F 10041
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:FITC \(MCA928F\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M412405:221111'

Printed on 12 Sep 2023