Datasheet: MCA1973GA BATCH NUMBER 154334

Description:	MOUSE ANTI PIG CD203a
Specificity:	CD203a
Other names:	SWC9
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	PM18-7
Isotype:	lgG1
Quantity:	0.1 mg

Product Details

Applications This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u>.

		Yes	No	Not Determined	Suggested Dilution	
	Flow Cytometry				1/25 - 1/200	
	Immunohistology - Frozen					
	Immunohistology - Paraffin					
	ELISA					
	Immunoprecipitation	-				
	Western Blotting					
	Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.					
Target Species	Pig					
Product Form	Purified IgG - liquid					

Preparation	Purified IgG prepared by affinity chromatography on Protein A
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)

Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Porcine alveolar macrophages.
Fusion Partners	Spleen cells from immunized mice were fused with P3X63-Ag8-653 murine myeloma cells (Kearney <i>et al.</i> 1979).
Specificity	Mouse anti Pig CD203a, clone PM18-7 recognizes porcine CD203a, originally clustered as SWC9 at the Second International Swine CD Workshop (<u>Dominguez <i>et al.</i> 1998</u>) and later identified as the porcine homologue of human ecto-nucleotidepyrophosphatase / phosphodiesterase 1 or <u>ENPP1</u> (<u>Petersen <i>et al.</i> 2007</u>).
	Mouse anti Pig CD203a was originally reported to immunoprecipitate two bands, one of ~;205 kDa and one of ~130 kDa (<u>Dominguez <i>et al.</i> 1998</u>) under both reducing and non-reducing conditions. Subsequent studies suggest that CD203a migrates as a homodimer of ~260 kDa under non-reducing conditions and a 130 kDa monomer under reducing conditions (<u>Petersen <i>et al.</i> 2007</u>) from preparations of porcine alveolar macrophages.
	CD203a is expressed widely in macrophage populations with notably high levels on alveolar macrophages (<u>Petersen <i>et al.</i> 2007</u> , <u>Hwang <i>et al.</i> 2015</u>), it is not expressed on monocyte populations (<u>McCullough <i>et al.</i> 1997</u> , <u>Hwang <i>et al.</i> 2015</u>).
	SWC1a, expressed at very much higher levels on monocytes than mature macrophages and CD203a (SWC9), expressed exclusively on mature tissue macrophages have been used as markers of monocyte-macrophage differentiation (<u>Sanchez <i>et al.</i> 1999</u>).
Flow Cytometry	Use 10ul of the suggested working dilution to label 1×10^6 cells in 100ul
References	 McCullough, K.C. <i>et al.</i> (1997) Phenotype of porcine monocytic cells: modulation of surface molecule expression upon monocyte differentiation into macrophages. <u>Vet</u> <u>Immunol Immunopathol. 58 (3-4): 265-75.</u> McCullough, K.C. <i>et al.</i> (1999) Intermediate stages in monocyte-macrophage differentiation modulate phenotype and susceptibility to virus infection. <u>Immunology. 98</u> (2): 203-12. Boersma, W.J. <i>et al.</i> (2001) Summary of workshop findings for porcine B-cell markers. <u>Vet Immunol Immunopathol. 80 (1-2): 63-78.</u> Domínguez, J. <i>et al.</i> (1998) Porcine myelomonocytic markers: summary of the Second International Swine CD Workshop. <u>Vet Immunol Immunopathol. 60 (3-4): 329-41.</u> Dominguez, J. <i>et al.</i> (1998) Workshop studies with monoclonal antibodies identifying a novel porcine differentiation antigen, SWC9. <u>Vet Immunol Immunopathol. 60 (3-4): 343-9.</u> Petersen, C.B. <i>et al.</i> (2007) Porcine ecto-nucleotide pyrophosphatase/phosphodiesterase 1 (NPP1/CD203a): cloning, transcription, expression, mapping, and identification of an NPP1/CD203a epitope for swine workshop cluster 9
	mapping, and identification of an NPP1/CD203a epitope for swine workshop cluster 9 (SWC9) monoclonal antibodies. <u>Dev Comp Immunol. 31 (6): 618-31.</u>

	7. Basta, S. et al. (1999) Modulation of monocytic cell activity and virus susceptibility
	during differentiation into macrophages. <u>J Immunol. 162 (7): 3961-9.</u>
	8. Gimeno, M. et al. (2011) Cytokine profiles and phenotype regulation of antigen
	presenting cells by genotype-I porcine reproductive and respiratory syndrome virus
	isolates. Vet Res. 42: 9.
	9. Sánchez, C. <i>et al.</i> (1999) The porcine 2A10 antigen is homologous to human CD163
	and related to macrophage differentiation. J Immunol. 162 (9): 5230-7.
	10 Cantu E et al. (2006) Depletion of pulmonary intravascular macrophages prevents
	hyperacute pulmonary xenograft dysfunction Transplantation 81 (8): 1157-64
	11 Basta S. et al. (2001) Linopolysaccharide and phorbol 12-myristate 13-acetate both
	impair monocyte differentiation, relating cellular function to virus suscentibility
	12 Lithgow B et al. (2014) Correlation of call surface marker expression with African
	12. Litingow, P. et al. (2014) Correlation of cell surface marker expression with Amcan
	22. On drag leave. D. et al. (2012) Drag sturie characterization of the mean state
	13. Ondrackova, P. et al. (2013) Phenotypic characterisation of the monocyte
	suppopulations in healthy adult pigs and < I> Salmonelia-Infected piglets by seven-colour
	flow cytometry. <u>Res Vet Sci. 94: 240 - 5.</u>
	14. Tsai, Y.C. <i>et al.</i> (2014) Differences in the expression of innate immune response-
	modulating genes in blood monocytes between subclinically porcine circovirus type s
	(PCV2)-infected and PCV2-free pigs prior to and after lipopolysaccharide stimulation in
	<i>vitro</i> <u>Taiwan Veterinary Journal. 40 (01): 37-48.</u>
	15. Hwang, J.H.et al. (2015) Characterization of monoclonal antibodies against porcine
	pulmonary alveolar macrophages of gnotobiotic miniature swine. <u>Biochem Biophys Res</u>
	<u>Commun. 461 (2): 427-34.</u>
	16. Shao, L. et al. (2016) Tissue-specific mRNA expression profiles of porcine Toll-like
	receptors at different ages in germ-free and conventional pigs. <u>Vet Immunol</u>
	Immunopathol. 171: 7-16.
	17. Burkard, C. <i>et al.</i> (2017) Precision engineering for PRRSV resistance in pigs:
	Macrophages from genome edited pigs lacking CD163 SRCR5 domain are fully resistant
	to both PRRSV genotypes while maintaining biological function, PLoS Pathog, 13 (2):
	e1006206
	18 Fernández-Caballero T <i>et al.</i> (2018) Phenotypic and functional characterization of
	norcine hone marrow monocyte subsets. Dev Comp Immunol. 81: 95-104
	10. Souttor C.A. et al. (2018) Phonotypic and functional modulations of porcino
	19. Sautter, C.A. <i>et al.</i> (2010) Phenotypic and functional modulations of porcine
	macrophages by interferons and interfeukin-4. Dev Comp immunor. 64. 161-92.
Further Reading	1 Piriou-Guzvlack L & Salmon H (2008) Membrane markers of the immune cells in
	swine: an undate. Vet Res. 39 (6): 54
	Swille. all apartes. <u>verifies. 66 (6). 64.</u>
Storage	Store at +4°C or at -20°C if preferred.
	Storage in frost-free freezers is not recommended.
	This product should be stored undiluted. Avoid repeated freezing and thawing as this may
	denature the antibody. Should this product contain a precipitate we recommend
	microcentrifugation before use.
Guarantee	12 months from date of despatch
Health And Safety	Material Safety Datasheet documentation #10040 available at:

Information	https://www.bio-rad-antibodies.com/SDS/MCA1973GA 10040			
Regulatory	For research purposes only			

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12)	RPE		
Goat Anti Mouse IgG IgA IgM (STAR87) <u>HRP</u>			
Goat Anti Mouse IgG (STAR76)	RPE		
Goat Anti Mouse IgG (STAR70)	FITC		
Goat Anti Mouse IgG (H/L) (STAR117)	Alk. Phos., DyLight®488, DyLight®550,		
	DyLight®650, DyLight®680, DyLight®800,		
	FITC, HRP		
Goat Anti Mouse IgG (STAR77)	HRP		
Rabbit Anti Mouse IgG (STAR13)	HRP		
Rabbit Anti Mouse IgG (STAR9)	<u>FITC</u>		
Goat Anti Mouse IgG (Fc) (STAR120)	FITC, HRP		
Recommended Negative Controls			

MOUSE IgG1 NEGATIVE CONTROL (MCA928)

North & South	Tel: +1 800 265 7376 Worldwide	Tel: +44 (0)186	5 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751	Fax: +44 (0)186	65 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-rad.com	Email: antibody	_sales_uk@bio-rad.	.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M366035:200529'

Printed on 26 May 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint