

Datasheet: MCA1973GA

Description:	MOUSE ANTI PIG CD203a
Specificity:	CD203a
Other names:	SWC9
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	PM18-7
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/25 - 1/200
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting			▪	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Pig
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% sodium azide (NaN ₃)

Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Porcine alveolar macrophages.
Fusion Partners	Spleen cells from immunized mice were fused with P3X63-Ag8-653 murine myeloma cells (Kearney et al. 1979).
Specificity	<p>Mouse anti Pig CD203a, clone PM18-7 recognizes porcine CD203a, originally clustered as SWC9 at the Second International Swine CD Workshop (Dominguez et al. 1998) and later identified as the porcine homologue of human ecto-nucleotidetriphosphatase / phosphodiesterase 1 or ENPP1 (Petersen et al. 2007).</p> <p>Mouse anti Pig CD203a was originally reported to immunoprecipitate two bands, one of ~205 kDa and one of ~130 kDa (Dominguez et al. 1998) under both reducing and non-reducing conditions. CD203a migrates as a homodimer of ~260 kDa under non-reducing conditions and a 130 kDa monomer under reducing conditions (Petersen et al. 2007) from preparations of porcine alveolar macrophages.</p> <p>CD203a is expressed widely in macrophage populations with notably high levels on alveolar macrophages (Petersen et al. 2007, Hwang et al. 2015), it is not expressed on monocyte populations (McCullough et al. 1997, Hwang et al. 2015).</p> <p>SWC1a, expressed at very much higher levels on monocytes than mature macrophages and CD203a (SWC9), expressed exclusively on mature tissue macrophages have been used as markers of monocyte-macrophage differentiation (Sanchez et al. 1999).</p>
Flow Cytometry	Use 10µl of the suggested working dilution to label 1x10 ⁶ cells in 100µl
References	<ol style="list-style-type: none"> 1. McCullough, K.C. <i>et al.</i> (1997) Phenotype of porcine monocytic cells: modulation of surface molecule expression upon monocyte differentiation into macrophages. Vet Immunol Immunopathol. 58 (3-4): 265-75. 2. Domínguez, J. <i>et al.</i> (1998) Porcine myelomonocytic markers: summary of the Second International Swine CD Workshop. Vet Immunol Immunopathol. 60 (3-4): 329-41. 3. Sánchez, C. <i>et al.</i> (1999) The porcine 2A10 antigen is homologous to human CD163 and related to macrophage differentiation. J Immunol. 162 (9): 5230-7. 4. Basta, S. <i>et al.</i> (1999) Modulation of monocytic cell activity and virus susceptibility during differentiation into macrophages. J Immunol. 162 (7): 3961-9. 5. Basta, S. <i>et al.</i> (2001) Lipopolysaccharide and phorbol 12-myristate 13-acetate both impair monocyte differentiation, relating cellular function to virus susceptibility. Immunology. 103 (4): 488-97. 6. Boersma, W.J. <i>et al.</i> (2001) Summary of workshop findings for porcine B-cell markers. Vet Immunol Immunopathol. 80 (1-2): 63-78. 7. Cantu, E. <i>et al.</i> (2006) Depletion of pulmonary intravascular macrophages prevents hyperacute pulmonary xenograft dysfunction. Transplantation. 81 (8): 1157-64. 8. Petersen, C.B. <i>et al.</i> (2007) Porcine ecto-nucleotide

- pyrophosphatase/phosphodiesterase 1 (NPP1/CD203a): cloning, transcription, expression, mapping, and identification of an NPP1/CD203a epitope for swine workshop cluster 9 (SWC9) monoclonal antibodies. [Dev Comp Immunol. 31 \(6\): 618-31.](#)
9. Gimeno, M. *et al.* (2011) Cytokine profiles and phenotype regulation of antigen presenting cells by genotype-I porcine reproductive and respiratory syndrome virus isolates. [Vet Res. 42: 9.](#)
 10. Ondrackova, P. *et al.* (2013) Phenotypic characterisation of the monocyte subpopulations in healthy adult pigs and *Salmonella*-infected piglets by seven-colour flow cytometry. [Res Vet Sci. 94: 240 - 5.](#)
 11. Tsai, Y.C. *et al.* (2014) Differences in the expression of innate immune response-modulating genes in blood monocytes between subclinically porcine circovirus type s (PCV2)-infected and PCV2-free pigs prior to and after lipopolysaccharide stimulation *in vitro* [Taiwan Vet J. 40 \(01\): 37-48.](#)
 12. Hwang, J.H. *et al.* (2015) Characterization of monoclonal antibodies against porcine pulmonary alveolar macrophages of gnotobiotic miniature swine. [Biochem Biophys Res Commun. 461 \(2\): 427-34.](#)
 13. Shao, L. *et al.* (2016) Tissue-specific mRNA expression profiles of porcine Toll-like receptors at different ages in germ-free and conventional pigs. [Vet Immunol Immunopathol. 171: 7-16.](#)
 14. Burkard, C. *et al.* (2017) Precision engineering for PRRSV resistance in pigs: Macrophages from genome edited pigs lacking CD163 SRCR5 domain are fully resistant to both PRRSV genotypes while maintaining biological function. [PLoS Pathog. 13 \(2\): e1006206.](#)
 15. Fernández-Caballero, T. *et al.* (2018) Phenotypic and functional characterization of porcine bone marrow monocyte subsets. [Dev Comp Immunol. 81: 95-104.](#)
 16. Sautter, C.A. *et al.* (2018) Phenotypic and functional modulations of porcine macrophages by interferons and interleukin-4. [Dev Comp Immunol. 84: 181-92.](#)
 17. Zimmermann, C.E. *et al.* (2021) Characterization of porcine mesenchymal stromal cells and their proliferative and osteogenic potential in long-term culture. [J Stem Cells Regen Med. 17 \(2\): 49-55.](#)
 18. Jarosova, R. *et al.* (2022) Cytokine expression by CD163+ monocytes in healthy and *Actinobacillus pleuropneumoniae*-infected pigs. [Res Vet Sci. 152: 1-9.](#)
 19. Petitpas, K. *et al.* (2022) Genetic modifications designed for xenotransplantation attenuate sialoadhesin-dependent binding of human erythrocytes to porcine macrophages. [Xenotransplantation. 29 \(6\): e12780.](#)
 20. Álvarez, B. *et al.* (2023) Porcine Macrophage Markers and Populations: An Update. [Cells. 12 \(16\): 2103.](#)
 21. Nieto-Pelegri, E. *et al.* (2020) Porcine CLEC12B is expressed on alveolar macrophages and blood dendritic cells. [Dev Comp Immunol. 111: 103767.](#)
 22. Boschetto, F. *et al.* (2024) Protocol for extracting and isolating porcine bone-marrow-derived macrophages from ribs. [STAR Protoc. 5 \(2\): 103085.](#)

Further Reading

1. Piriou-Guzylack, L. & Salmon, H. (2008) Membrane markers of the immune cells in swine: an update. [Vet Res. 39 \(6\): 54.](#)

Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for

short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA1973GA 10040
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...)	RPE
Goat Anti Mouse IgG IgA IgM (STAR87...)	HRP
Goat Anti Mouse IgG (STAR76...)	RPE
Goat Anti Mouse IgG (STAR70...)	FITC
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight®488 , DyLight®550 , DyLight®650 , DyLight®680 , DyLight®800 , FITC , HRP
Goat Anti Mouse IgG (STAR77...)	HRP
Rabbit Anti Mouse IgG (STAR13...)	HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)
'M412412:221111'

Printed on 26 May 2024