

Datasheet: MCA1984

BATCH NUMBER 149446

Description:	MOUSE ANTI HUMAN CD173
Specificity:	CD173
Other names:	BLOOD GROUP H TYPE 2
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	BRIC231
Isotype:	IgG1
Quantity:	0.2 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting			▪	
Immunofluorescence	▪			
Haemagglutination	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species

Human

Species Cross Reactivity

Reacts with: Pig

N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

Product Form

Purified IgG - liquid

Buffer Solution	TRIS buffered glycine
Preservative Stabilisers	0.09% Sodium Azide
Approx. Protein Concentrations	IgG concentration 1.0mg/ml
Immunogen	Human erythroleukemic cell line (HEL) established from a 30 year old patient with relapsed erythroleukemia following treatment for Hodgkin lymphoma.
RRID	AB_323275
Specificity	Mouse anti Human CD173 antibody, clone BRIC231 recognizes human type 2 H blood group antigen, also known as CD173. Active H substances in man, are expressed by many cells and tissues and also by erythrocytes.
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
References	<ol style="list-style-type: none"> 1. Clausen, H. and Hakomori, S. (1989) ABH and related histo-blood group antigens; immunochemical differences in carrier isotypes and their distribution. Vox Sang 56(1): 1-20. 2. Siegel, G. <i>et al.</i> (2013) Phenotype, donor age and gender affect function of human bone marrow-derived mesenchymal stromal cells. BMC Med. 11: 146. 3. Hotta, H. <i>et al.</i> (2013) Lewis y antigen is expressed in oral squamous cell carcinoma cell lines and tissues, but disappears in the invasive regions leading to the enhanced malignant properties irrespective of sialyl-Lewis x. Glycoconj J. 30: 585-97. 4. Guix, S. <i>et al.</i> (2007) Norwalk virus RNA is infectious in mammalian cells. J Virol. 81: 12238-48. 5. Sharpe, C. <i>et al.</i> (2014) Mixed field reactions in ABO and Rh typing chimerism likely resulting from twin haematopoiesis. Blood Transfus. 12: 608-10. 6. Cheetham, S. <i>et al.</i> (2007) Binding patterns of human norovirus-like particles to buccal and intestinal tissues of gnotobiotic pigs in relation to A/H histo-blood group antigen expression. J Virol. 81: 3535-44. 7. Hutson, A.M. <i>et al.</i> (2003) Norwalk virus-like particle hemagglutination by binding to h histo-blood group antigens. J Virol. 77: 405-15. 8. Matsumoto, S. <i>et al.</i> (2015) A Cytotoxic Antibody Recognizing Lacto-N-fucopentaose I (LNFP I) on Human Induced Pluripotent Stem (hiPS) Cells. J Biol Chem. 290 (33): 20071-85. 9. Schäfer R, <i>et al.</i> (2020) Modulating endothelial adhesion and migration impacts stem cell therapies efficacy. EBioMedicine. 60:102987. 10. Lin, R-J. <i>et al.</i> (2019) B3GALT5 Knockout Alters Glycosphingolipid Profile and Facilitates Transition to Human Naïve Pluripotency (December 27, 2019). Stem Cell Dec 27 [Epub ahead of print - Reviewed]
Storage	Store at +4°C or at -20°C if preferred. This product should be stored undiluted.

Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee	12 months from date of despatch
------------------	---------------------------------

Health And Safety Information	Material Safety Datasheet documentation #10072 available at: https://www.bio-rad-antibodies.com/SDS/MCA1984 10072
--------------------------------------	--

Regulatory	For research purposes only
-------------------	----------------------------

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...)	RPE
Goat Anti Mouse IgG IgA IgM (STAR87...)	HRP
Goat Anti Mouse IgG (STAR76...)	RPE
Goat Anti Mouse IgG (STAR70...)	FITC
Rabbit Anti Mouse IgG (STAR13...)	HRP
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC
Goat Anti Mouse IgG (STAR77...)	HRP
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight®488 , DyLight®550 , DyLight®650 , DyLight®680 , DyLight®800 , FITC , HRP

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M366046:200529'

Printed on 19 Mar 2025