

Datasheet: MCA1999S

BATCH NUMBER 180425

Description:	MOUSE ANTI DOG CD8 ALPHA
Specificity:	CD8 ALPHA
Format:	S/N
Product Type:	Monoclonal Antibody
Clone:	CA9.JD3
Isotype:	IgG2a
Quantity:	2 ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat
Immunohistology - Frozen	▪			Neat - 1/10
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation	▪			
Functional Assays (1)			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1) Removal of sodium azide is recommended prior to use in functional assays.

Target Species	Dog
Product Form	Tissue Culture Supernatant - liquid
Preservative Stabilisers	0.1% Sodium Azide (NaN ₃)
Immunogen	Canine Thymocytes
External Database Links	UniProt: P33706 Related reagents

Entrez Gene:[403157](#) CD8A [Related reagents](#)

RRID AB_323370

Fusion Partners Spleen cells from immunised Balb/c mice were fused with cells of the P3X63-Ag.653 myeloma cell line.

Specificity **Mouse anti Dog CD8 alpha antibody, clone CA9.JD3** recognizes the canine CD8 alpha chain which is expressed by thymocytes, peripheral T cells in the blood and lymphoid organs .

Mouse anti Dog CD8 α clone CA9.JD3 has been reported to inhibit cytotoxic T lymphocyte function ([Cobbold *et al.* 1994](#)). CA9.JD3 immunoprecipitates a heterodimer of ~32 kDa and ~36 kDa (reduced) from canine Tumor cells and from thymocytes.

Flow Cytometry Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul.

Histology Positive Control Tissue Spleen, lymph node

References

1. Moore, P.F. *et al.* (1992) Monoclonal antibodies specific for canine CD4 and CD8 define functional T-lymphocyte subsets and high-density expression of CD4 by canine neutrophils. [Tissue Antigens 40\(2\): 75-85.](#)
2. Cobbold, S. & Metcalfe, S. (1994) Monoclonal antibodies that define canine homologues of human CD antigens: summary of the First International Canine Leukocyte Antigen Workshop (CLAW). [Tissue Antigens. 43 \(3\): 137-54.](#)
3. Veenhof, E.Z. *et al.* (2011) Characterisation of T cell phenotypes, cytokines and transcription factors in the skin of dogs with cutaneous adverse food reactions. [Vet J. 187 \(3\): 320-4.](#)
4. Yuasa, K. *et al.* (2007) Injection of a recombinant AAV serotype 2 into canine skeletal muscles evokes strong immune responses against transgene products. [Gene Ther. 14: 1249-60.](#)
5. Huyghe, S. *et al.* (2015) The Microscopic Structure of the Omentum in Healthy Dogs: The Mystery Unravelling. [Anat Histol Embryol. Jul 23. \[Epub ahead of print\]](#)
6. Lin Shiow-Chen *et al.* (2014) Immune Characterization of Peripheral Blood Mononuclear cells of the Dogs Restored from Inoculation of Canine Transmissible Venereal Tumor Cells. [Tai Vet J. 40 \(04\): 181-90.](#)
7. Constantinoiu, C.C. *et al.* (2015) Mucosal tolerance of the hookworm *Ancylostoma caninum* in the gut of naturally infected wild dogs. [Parasite Immunol. Jul 27 \[Epub ahead of print\].](#)

Storage Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing

as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10336 available at: https://www.bio-rad-antibodies.com/SDS/MCA1999S10336
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...)	RPE
Goat Anti Mouse IgG IgA IgM (STAR87...)	HRP
Goat Anti Mouse IgG (STAR76...)	RPE
Rabbit Anti Mouse IgG (STAR13...)	HRP
Goat Anti Mouse IgG (STAR70...)	FITC
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight®488 , DyLight®550 , DyLight®650 , DyLight®680 , DyLight®800 , FITC , HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC
Goat Anti Mouse IgG (STAR77...)	HRP
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP

Recommended Negative Controls

[MOUSE IgG2a NEGATIVE CONTROL \(MCA929\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M366060:200529'

Printed on 19 Jan 2024