

Datasheet: MCA199B BATCH NUMBER 241014

Description:	RAT ANTI MOUSE IgM HEAVY CHAIN:Biotin
Specificity:	IgM HEAVY CHAIN
Format:	Biotin
Product Type:	Monoclonal Antibody
Clone:	LO-MM-9
Isotype:	lgG2a
Quantity:	0.5 mg

Product Details

Applications	This product has been re derived from testing withi communications from the information. For general	in our labo e originato protocol r	oratories, ors. Please	peer-reviewed publica refer to references in	ations or personal ndicated for further
	rad-antibodies.com/proto				
		Yes	No	Not Determined	Suggested Dilution
	Flow Cytometry			• _	
	Immunohistology - Frozen				
	Immunohistology - Paraffin ELISA				250ng/ml
	Western Blotting	-		-	25019/111
	Where this antibody has	not boon	tootod for	uno in o porticular too	photos this doos not
	•			•	•
	necessarily exclude its us		•		•
	a guide only. It is recomn				for use in their own
	system using appropriate	e negative	positive c	controls.	
Target Species	Mouse				
Product Form	Purified IgG conjugated t	o Biotin -	liquid		
Preparation	Purified IgG prepared by culture supernatant	affinity cł	nromatogra	aphy on mouse IgM/S	epharose 4B from tissue
Buffer Solution	Phosphate buffered salin	е			
Preservative Stabilisers	0.1% Sodium Azide (NaN 50% Glycerol	N ₃)			
Approx. Protein Concentrations	IgG concentration 1 mg/r	nl			

Immunogen	Polyclonal and monoclonal mouse IgM.
External Database	UniProt:
Links	
	P01872 Related reagents
	P01873 Related reagents
	Entrez Gene:
	16019 Ighm Related reagents
	16019 Ighm Related reagents
RRID	AB_321870
Fusion Partners	Spleen cells from immunised LOU/c rats were fused with cells of the IR983F rat myeloma cell line.
Specificity	Rat anti Mouse IgM, clone LO-MM-9 is directed against the mu chain of mouse IgM.
ELISA	This product may be used in an indirect ELISA or as a secondary reagent in a sandwich ELISA together with <u>MCA1293</u> as the capture reagent and and Streptavidin:HRP (<u>STAR5B</u>) as the detection reagent.
References	 Achtman AH <i>et al.</i> (2003) <i>Plasmodium chabaudi chabaudi</i> infection in mice induces strong B cell responses and striking but temporary changes in splenic cell distribution. J Immunol. 171 (1): 317-24. Bazin, H. <i>et al.</i> (1984) Rat monoclonal antibodies. II. A rapid and efficient method of purification from ascitic fluid or serum. J Immunol Methods. 71: 9-16. Bazin, H. <i>et al.</i> (1974) Three classes and four (sub)classes of rat immunoglobulins: IgM, IgA, IgE and IgG1, IgG2a, IgG2b, IgG2c. Eur J Immunol. 4 (1): 44-8. Bermejo, D.A. <i>et al.</i> (2011) <i>Trypanosoma cruzi</i> infection induces a massive extrafollicular and follicular splenic B-cell response which is a high source of non-parasite-specific antibodies. Immunology. 132 (1): 123-33. Busund, L.T. <i>et al.</i> (2003) Spontaneously formed tumorigenic hybrids of Meth A sarcoma cells and macrophages <i>in vivo</i>. Int J Cancer. 106 (2): 153-9. Huang, C.H. <i>et al.</i> (2011) Airway inflammation and IgE production induced by dust mite allergen-specific memory/effector Th2 cell line can be effectively attenuated by IL-35. J Immunol. 187: 462-71. Sreekanthreddy, P. <i>et al.</i> (2015) A three-dimensional model of the human blood-brain barrier to analyse the transport of nanoparticles and astrocyte/endothelial interactions. F1000Res. 4: 1279.
Storage	Store at +4°C or at -20°C if preferred. This product should be stored undiluted.
	Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

I				heet documentation #10 htibodies.com/SDS/MCA	.t:	
Regulatory For research purpose		purpose	s only			
North & South America	Tel: +1 800 265 73 Fax: +1 919 878 3		Vorldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50
Email: antibody_sales_us@bio-rad.com		om	Email: antibody_sales_uk@bio	Email: antibody_sales_de@bio-rad.con		
To find a b						p-rad-antibodies.com/datas

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint