

## Datasheet: MCA199F

**BATCH NUMBER 151254**

<b>Description:</b>	RAT ANTI MOUSE IgM HEAVY CHAIN:FITC
<b>Specificity:</b>	IgM HEAVY CHAIN
<b>Format:</b>	FITC
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	LO-MM-9
<b>Isotype:</b>	IgG2a
<b>Quantity:</b>	0.5 mg

### Product Details

#### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			5ug/ml

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Mouse						
<b>Product Form</b>	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid						
<b>Max Ex/Em</b>	<table border="1"> <thead> <tr> <th>Fluorophore</th> <th>Excitation Max (nm)</th> <th>Emission Max (nm)</th> </tr> </thead> <tbody> <tr> <td>FITC</td> <td>490</td> <td>525</td> </tr> </tbody> </table>	Fluorophore	Excitation Max (nm)	Emission Max (nm)	FITC	490	525
Fluorophore	Excitation Max (nm)	Emission Max (nm)					
FITC	490	525					
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on mouse IgM/Sepharose 4B from tissue culture supernatant						
<b>Buffer Solution</b>	Phosphate buffered saline						
<b>Preservative Stabilisers</b>	0.1% Sodium Azide 50% Glycerol						
<b>Approx. Protein Concentrations</b>	IgG concentration 1 mg/ml						

**Immunogen** Polyclonal and monoclonal mouse IgM.

---

**External Database**

**Links**

**UniProt:**

[P01872](#) [Related reagents](#)

[P01873](#) [Related reagents](#)

**Entrez Gene:**

[16019](#) Ighm [Related reagents](#)

[16019](#) Ighm [Related reagents](#)

---

**RRID** AB\_321871

---

**Fusion Partners** Spleen cells from immunised LOU/c rats were fused with cells of the IR983F rat myeloma cell line.

---

**Specificity** **Rat anti Mouse IgM, clone LO-MM-9** is directed against the mu chain of mouse IgM.

---

**References**

1. Achtman AH *et al.* (2003) *Plasmodium chabaudi chabaudi* infection in mice induces strong B cell responses and striking but temporary changes in splenic cell distribution. [J Immunol. 171 \(1\): 317-24.](#)
  2. Bazin, H. *et al.* (1984) Rat monoclonal antibodies. II. A rapid and efficient method of purification from ascitic fluid or serum. [J Immunol Methods. 71: 9-16.](#)
  3. Bazin, H. *et al.* (1974) Three classes and four (sub)classes of rat immunoglobulins: IgM, IgA, IgE and IgG1, IgG2a, IgG2b, IgG2c. [Eur J Immunol. 4 \(1\): 44-8.](#)
  4. Bermejo, D.A. *et al.* (2011) *Trypanosoma cruzi* infection induces a massive extrafollicular and follicular splenic B-cell response which is a high source of non-parasite-specific antibodies. [Immunology. 132 \(1\): 123-33.](#)
  5. Busund, L.T. *et al.* (2003) Spontaneously formed tumorigenic hybrids of Meth A sarcoma cells and macrophages *in vivo*. [Int J Cancer. 106 \(2\): 153-9.](#)
  6. Huang, C.H. *et al.* (2011) Airway inflammation and IgE production induced by dust mite allergen-specific memory/effector Th2 cell line can be effectively attenuated by IL-35. [J Immunol. 187: 462-71.](#)
  7. Sreekanthreddy, P. *et al.* (2015) A three-dimensional model of the human blood-brain barrier to analyse the transport of nanoparticles and astrocyte/endothelial interactions. [F1000Res. 4: 1279.](#)
- 

**Storage** Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

---

**Guarantee** 12 months from date of despatch

---

**Health And Safety Information**      Material Safety Datasheet documentation #10328 available at:  
<https://www.bio-rad-antibodies.com/SDS/MCA199F10328>

---

**Regulatory**                      For research purposes only

---

## Related Products

### Recommended Negative Controls

[RAT IgG2a NEGATIVE CONTROL:FITC \(MCA1212F\)](#)

### Recommended Useful Reagents

[MOUSE SEROBLOCK FcR \(BUF041A\)](#)

[MOUSE SEROBLOCK FcR \(BUF041B\)](#)

**North & South**      Tel: +1 800 265 7376

**America**              Fax: +1 919 878 3751

                            Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

                            Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

                            Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)  
'M366062:200529'

**Printed on 18 Jan 2024**

---

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)