

## Datasheet: MCA199F

**BATCH NUMBER 162086**

<b>Description:</b>	RAT ANTI MOUSE IgM HEAVY CHAIN:FITC
<b>Specificity:</b>	IgM HEAVY CHAIN
<b>Format:</b>	FITC
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	LO-MM-9
<b>Isotype:</b>	IgG2a
<b>Quantity:</b>	0.5 mg

### Product Details

#### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			5ug/ml

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Mouse						
<b>Product Form</b>	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid						
<b>Max Ex/Em</b>	<table border="1"> <thead> <tr> <th>Fluorophore</th> <th>Excitation Max (nm)</th> <th>Emission Max (nm)</th> </tr> </thead> <tbody> <tr> <td>FITC</td> <td>490</td> <td>525</td> </tr> </tbody> </table>	Fluorophore	Excitation Max (nm)	Emission Max (nm)	FITC	490	525
Fluorophore	Excitation Max (nm)	Emission Max (nm)					
FITC	490	525					
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on mouse IgM/Sepharose 4B from tissue culture supernatant						
<b>Buffer Solution</b>	Phosphate buffered saline						
<b>Preservative Stabilisers</b>	0.1% Sodium Azide 50% Glycerol						
<b>Approx. Protein Concentrations</b>	IgG concentration 1 mg/ml						

**Immunogen** Polyclonal and monoclonal mouse IgM.

---

**External Database**

**Links**

**UniProt:**

[P01872](#) [Related reagents](#)

[P01873](#) [Related reagents](#)

**Entrez Gene:**

[16019](#) Ighm [Related reagents](#)

[16019](#) Ighm [Related reagents](#)

---

**RRID** AB\_321871

---

**Fusion Partners** Spleen cells from immunized LOU/c rats were fused with cells of the IR983F rat myeloma cell line.

---

**Specificity** **Rat anti Mouse IgM, clone LO-MM-9** is directed against the mu chain of mouse IgM.

---

**References**

1. Bazin, H. *et al.* (1974) Three classes and four (sub)classes of rat immunoglobulins: IgM, IgA, IgE and IgG1, IgG2a, IgG2b, IgG2c. [Eur J Immunol. 4 \(1\): 44-8.](#)
  2. Bazin, H. *et al.* (1984) Rat monoclonal antibodies. II. A rapid and efficient method of purification from ascitic fluid or serum. [J Immunol Methods. 71: 9-16.](#)
  3. Busund, L.T. *et al.* (2003) Spontaneously formed tumorigenic hybrids of Meth A sarcoma cells and macrophages *in vivo*. [Int J Cancer. 106 \(2\): 153-9.](#)
  4. Achtman AH *et al.* (2003) *Plasmodium chabaudi chabaudi* infection in mice induces strong B cell responses and striking but temporary changes in splenic cell distribution. [J Immunol. 171 \(1\): 317-24.](#)
  5. Huang, C.H. *et al.* (2011) Airway inflammation and IgE production induced by dust mite allergen-specific memory/effector Th2 cell line can be effectively attenuated by IL-35. [J Immunol. 187: 462-71.](#)
  6. Bermejo, D.A. *et al.* (2011) *Trypanosoma cruzi* infection induces a massive extrafollicular and follicular splenic B-cell response which is a high source of non-parasite-specific antibodies. [Immunology. 132 \(1\): 123-33.](#)
  7. Sreekanthreddy, P. *et al.* (2015) A three-dimensional model of the human blood-brain barrier to analyse the transport of nanoparticles and astrocyte/endothelial interactions. [F1000Res. 4: 1279.](#)
- 

**Storage**

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

---

**Guarantee**

12 months from date of despatch

---

**Health And Safety**

Material Safety Datasheet documentation #10328 available at:

**Information** <https://www.bio-rad-antibodies.com/SDS/MCA199F10328>

---

**Regulatory** For research purposes only

---

## Related Products

### Recommended Negative Controls

[RAT IgG2a NEGATIVE CONTROL:FITC \(MCA1212F\)](#)

### Recommended Useful Reagents

[MOUSE SEROBLOCK FcR \(BUF041A\)](#)

[MOUSE SEROBLOCK FcR \(BUF041B\)](#)

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M385021:210513'

**Printed on 18 Jan 2024**

---

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)