

## Datasheet: MCA199F

Description:	RAT ANTI MOUSE IgM HEAVY CHAIN:FITC
Specificity:	IgM HEAVY CHAIN
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	LO-MM-9
Isotype:	lgG2a
Quantity:	0.5 mg

## **Product Details**

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .							
		Yes	No	Not Determined	Suggested Dilution			
	Flow Cytometry	-			5ug/ml			
	Where this antibody has not been tested for use in a particular technique this does not							
	necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.							
Target Species	Mouse							
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid							
Max Ex/Em	Fluorophore	Excitation Ma	ax (nm)	Emission Max (nm)				
	FITC	490		525				
Preparation	Purified IgG prepared by affinity chromatography on mouse IgM/Sepharose 4B from tissue culture supernatant							
Buffer Solution	Phosphate buffered saline							
Preservative	0.1% Sodium Azide							
Stabilisers	50% Glycerol							
Approx. Protein Concentrations	IgG concentration 1 m	g/ml						

Immunogen	Polyclonal and monoclonal mouse IgM.					
External Database Links	UniProt:					
	P01872 Related reagents					
	P01873 Related reagents					
	Entrez Gene:					
	16019 Ighm Related reagents					
	<u>16019</u> Ighm <u>Related reagents</u>					
RRID	AB_321871					
Fusion Partners	Spleen cells from immunized LOU/c rats were fused with cells of the IR983F rat myeloma cell line.					
Specificity	Rat anti Mouse IgM, clone LO-MM-9 is directed against the mu chain of mouse IgM.					
References	<ol> <li>Bazin, H. <i>et al.</i> (1974) Three classes and four (sub)classes of rat immunoglobulins: IgM, IgA, IgE and IgG1, IgG2a, IgG2b, IgG2c. <u>Eur J Immunol. 4 (1): 44-8.</u></li> <li>Bazin, H. <i>et al.</i> (1984) Rat monoclonal antibodies. II. A rapid and efficient method of purification from ascitic fluid or serum. <u>J Immunol Methods. 71: 9-16.</u></li> <li>Busund, L.T. <i>et al.</i> (2003) Spontaneously formed tumorigenic hybrids of Meth A sarcoma cells and macrophages <i>in vivo</i>. Int J Cancer. 106 (2): 153-9.</li> <li>Achtman AH <i>et al.</i> (2003) <i>Plasmodium chabaudi chabaudi</i> infection in mice induces strong B cell responses and striking but temporary changes in splenic cell distribution. J Immunol. 171 (1): 317-24.</li> <li>Huang, C.H. <i>et al.</i> (2011) Airway inflammation and IgE production induced by dust mite allergen-specific memory/effector Th2 cell line can be effectively attenuated by IL-35. J Immunol. 187: 462-71.</li> <li>Bermejo, D.A. <i>et al.</i> (2011) <i>Trypanosoma cruzi</i> infection induces a massive extrafollicular and follicular splenic B-cell response which is a high source of non-parasite- specific antibodies. Immunology. 132 (1): 123-33.</li> <li>Sreekanthreddy, P. <i>et al.</i> (2015) A three-dimensional model of the human blood-brain barrier to analyse the transport of nanoparticles and astrocyte/endothelial interactions. F1000Res. 4: 1279.</li> </ol>					
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in					
	frost-free freezers is not recommended. This product is photosensitive and should be protected from light.					
Guarantee	12 months from date of despatch					
Health And Safety	Material Safety Datasheet documentation #10328 available at:					

Informati	on <u>https://</u> 10328								
Regulato	<b>ry</b> For res	For research purposes only							
Relate	d Products								
Recomm	nended Negative (	Controls							
RAT IgG2	NEGATIVE CONTRO	L:FITC (MCA121	1 <u>2F)</u>						
Recomm	nended Useful Rea	agents							
MOUSE S	EROBLOCK FcR (BUF	041A)							
MOUSE S	EROBLOCK FcR (BUF	<u>041B)</u>							
North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody sales us@t	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody sales uk@bid	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com				
To find a b	0		0		io-rad-antibodies.com/datasheets				

## Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint