

Datasheet: MCA2028

Description:	MOUSE ANTI HUMAN CD29
Specificity:	CD29
Other names:	INTEGRIN BETA 1 CHAIN
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	12G10
Isotype:	IgG1
Quantity:	0.2 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/25 - 1/50
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA	▪			10ug/ml
Immunofluorescence	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species

Human

Species Cross Reactivity

Reacts with: Mink, Rabbit

Does not react with: Rat, Mouse

N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

Product Form

Purified IgG - liquid

Preparation

Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant

Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% sodium azide (NaN ₃)
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Purified human beta1 integrin preparation from HT1080 fibrosarcoma cell extract
External Database Links	<p>UniProt: P05556 Related reagents</p> <p>Entrez Gene: 3688 ITGB1 Related reagents</p>
Synonyms	FNRB, MDF2, MSK12
RRID	AB_321646
Fusion Partners	Spleen cells from an immunised BALB/c mice were fused with cells of the X63/Ag8.653 mouse myeloma cell line
Specificity	<p>Mouse anti Human CD29 monoclonal antibody, clone 12G10 recognizes human CD29 also known as beta1 integrin or VLA-4 subunit alpha. CD29 is a ~130 kDa under reducing, ~115 kDa under non-reducing conditions single pass type I transmembrane glycoprotein. CD29 acts as the common beta subunit of the heterodimeric very late antigens 1-6, complexing with CD49a-f respectively where it forms part of the receptors for laminin , collagen and fibronectin. the VLA heterodimers mediate cell-cell and cell-matrix interactions.</p> <p>Mouse anti Human CD29, clone 12G10 binding to cells adhering via VLA-4 results in actin cytoskeletal disruption and subsequent inhibition of attachment and spreading whilst 12G10 binding to cells adhering via VLA-5 results in enhancement of both these processes (Humphries et al. 2005). Clone 12G10 enhances alpha 5 beta 1-fibronectin interactions and binds to a region of CD25 containing the binding epitopes of several other anti CD29 antibody clones. However, unlike these, binding of 12G10 is enhanced in the presence of ligands such as fibronectin fragments (Mould et al. 1995). Binding of antibody clone 12G10 to the integrin β1 subunit is affected by divalent cations and the binding epitope appears to be located around residues 207-218 in the b1 subunit putative A-domain (Mould et al. 1998).</p>
References	<ol style="list-style-type: none"> 1. Mould, A.P. <i>et al.</i> (1995) Regulation of integrin alpha 5 beta 1 function by anti-integrin antibodies and divalent cations. Biochem Soc Trans. 23 (3): 395S. 2. Mould, A.P. <i>et al.</i> (1998) Regulation of integrin function: evidence that bivalent-cation-induced conformational changes lead to the unmasking of ligand-binding sites within

- integrin alpha5 beta1. [Biochem J. 331 \(Pt 3\) \(Pt 3\): 821-8.](#)
3. Sodek, K.L. *et al.* (2009) Compact spheroid formation by ovarian cancer cells is associated with contractile behavior and an invasive phenotype. [Int J Cancer. 124: 2060-70.](#)
 4. Matthews, B.D. *et al.* (2010) Ultra-rapid activation of TRPV4 ion channels by mechanical forces applied to cell surface [beta]1 integrins. [Integr Biol \(Camb\). 2: 435-42.](#)
 5. Aasted, B. *et al.* (2007) Reactivity of monoclonal antibodies to human CD antigens with cells from mink. [Vet Immunol Immunopathol. 119: 27-37.](#)
 6. Kawaguchi, N. *et al.* (2003) ADAM12 induces actin cytoskeleton and extracellular matrix reorganization during early adipocyte differentiation by regulating beta1 integrin function. [J Cell Sci. 116: 3893-904.](#)
 7. Loughran, G. *et al.* (2005) Mystique is a new insulin-like growth factor-I-regulated PDZ-LIM domain protein that promotes cell attachment and migration and suppresses Anchorage-independent growth. [Mol Biol Cell. 2005 Apr;16\(4\):1811-22.](#)
 8. Werner, J. *et al.* (2012) Expression of integrins and Toll-like receptors in cervical cancer: Effect of infectious agents. [Innate Immun. 18: 55-69.](#)
 9. Iba, K. *et al.* (2000) The cysteine-rich domain of human ADAM 12 supports cell adhesion through syndecans and triggers signaling events that lead to beta1 integrin-dependent cell spreading. [J Cell Biol. 149: 1143-56.](#)
 10. Meng, X. *et al.* (2005) Evidence for the presence of a low-mass beta1 integrin on the cell surface. [J Cell Sci. 118: 4009-16.](#)
 11. Whittard, J.D. and Akiyama, S.K. (2001) Positive regulation of cell-cell and cell-substrate adhesion by protein kinase A. [J Cell Sci. 114: 3265-72.](#)
 12. Rodriguez-Teja, M. *et al.* (2015) AGE-modified basement membrane cooperates with Endo180 to promote epithelial cell invasiveness and decrease prostate cancer survival. [J Pathol. 235 \(4\): 581-92.](#)
 13. Zhong, C. *et al.* (1998) Rho-mediated contractility exposes a cryptic site in fibronectin and induces fibronectin matrix assembly. [J Cell Biol. 141: 539-51.](#)
 14. Zhou, J. *et al.* (2008) Salvicine inactivates beta 1 integrin and inhibits adhesion of MDA-MB-435 cells to fibronectin via reactive oxygen species signaling. [Mol Cancer Res. 6: 194-204.](#)
 15. Thodeti, C.K. *et al.* (2003) ADAM12/syndecan-4 signaling promotes beta 1 integrin-dependent cell spreading through protein kinase Calpha and RhoA. [J Biol Chem. 278: 9576-84.](#)
 16. Lee, H. *et al.* (2006) A critical role for the membrane-type 1 matrix metalloproteinase in collagen phagocytosis. [Mol Biol Cell. 17: 4812-26.](#)
 17. Gravelle, S. *et al.* (2010) Up-regulation of integrin expression in lung adenocarcinoma cells caused by bacterial infection: in vitro study. [Innate Immun. 2010 Feb;16\(1\):14-26.](#)
 18. Jović, M. *et al.* (2007) EHD1 regulates beta1 integrin endosomal transport: effects on focal adhesions, cell spreading and migration. [J Cell Sci. 120: 802-14.](#)
 19. Newsome, P.N. *et al.* (2004) Serum from patients with fulminant hepatic failure causes hepatocyte detachment and apoptosis by a beta(1)-integrin pathway. [Hepatology. 40: 636-45.](#)
 20. Piccinno, M.S. *et al.* (2013) Adipose stromal/stem cells assist fat transplantation reducing necrosis and increasing graft performance. [Apoptosis. 18 \(10\): 1274-89.](#)
 21. Lee, J. *et al.* (2013) Phloridzin isolated from *Acanthopanax senticosus* promotes proliferation of alpha6 integrin (CD 49f) and beta1 integrin (CD29) enriched for a primary

keratinocyte population through the ERK-mediated mTOR pathway. [Arch Dermatol Res. 305 \(8\): 747-54.](#)

22. Xie, G. *et al.* (2021) Characterization of HIV-induced remodeling reveals differences in infection susceptibility of memory CD4⁺ T cell subsets *in vivo*. [Cell Rep. 35 \(4\): 109038.](#)

23. Mastrolia, I. *et al.* (2022) Autologous Marrow Mesenchymal Stem Cell Driving Bone Regeneration in a Rabbit Model of Femoral Head Osteonecrosis. [Pharmaceutics. 14\(10\):2127.](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10040 available at: 10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR77...) [HRP](#)
Rabbit Anti Mouse IgG (STAR12...) [RPE](#)
Goat Anti Mouse IgG (STAR70...) [FITC](#)
Goat Anti Mouse IgG IgA IgM (STAR87...) [Alk. Phos.](#), [HRP](#)
Goat Anti Mouse IgG (STAR76...) [RPE](#)
Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight®488](#), [DyLight®550](#),
[DyLight®650](#), [DyLight®680](#), [DyLight®800](#),
[FITC](#), [HRP](#)
Rabbit Anti Mouse IgG (STAR13...) [HRP](#)
Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)
Rabbit Anti Mouse IgG (STAR9...) [FITC](#)

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M412529:221112'

Printed on 17 Jan 2023
