

## Datasheet: MCA2035S BATCH NUMBER 159900

| Description:  | ption: MOUSE ANTI DOG CD45 |  |  |
|---------------|----------------------------|--|--|
| Specificity:  | CD45                       |  |  |
| Other names:  | LCA                        |  |  |
| Format:       | S/N                        |  |  |
| Product Type: | Monoclonal Antibody        |  |  |
| Clone:        | CA12.10C12                 |  |  |
| lsotype:      | lgG1                       |  |  |
| Quantity:     | 2 ml                       |  |  |
|               |                            |  |  |

## **Product Details**

| Applications                   | This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal |   |                                      |                                    |                          |  |  |
|--------------------------------|--|---|--------------------------------------|------------------------------------|--------------------------|--|--|
|                                | communications from the originators. Please refer to references indicated for further  |   |                                      |                                    |                          |  |  |
|                                | Ŭ  | information. For general protocol recommendations, please visit <u>www.bio-</u> |                                      |                                    |                          |  |  |
|                                | rad-antibodies.com/proto   |   |                                      |                                    |                          |  |  |
|                                |  | Yes   | No                                   | Not Determined                     | Suggested Dilution       |  |  |
|                                | Flow Cytometry   | •   |                                      |                                    | Neat                     |  |  |
|                                | Immunohistology - Frozen   | •   |                                      |                                    |                          |  |  |
|                                | Immunohistology - Paraffin<br>(1)  | •   |                                      |                                    |                          |  |  |
|                                | ELISA  |   |                                      | •                                  |                          |  |  |
|                                | Immunoprecipitation  |   |                                      | •                                  |                          |  |  |
|                                | necessarily exclude its u  | se in suc   | h procedu                            | res. Suggested workin              | g dilutions are given as |  |  |
|                                | a guide only. It is recommoded a guide normal system using appropriate (1) <b>This product require</b>   | nended tl<br>e negative   | nat the us<br>e/positive             | er titrates the antibody controls. | for use in their own     |  |  |
| Target Species                 | a guide only. It is recomm<br>system using appropriate   | nended tl<br>e negative   | nat the us<br>e/positive             | er titrates the antibody controls. | for use in their own     |  |  |
| Target Species<br>Product Form | a guide only. It is recomm<br>system using appropriate<br>(1) <b>This product require</b>  | nended tl<br>e negative<br>es heat n  | nat the us<br>e/positive<br>nediated | er titrates the antibody controls. | for use in their own     |  |  |
|                                | a guide only. It is recomm<br>system using appropriate<br>(1) <b>This product require</b><br>Dog   | nended tl<br>e negative<br>es heat n  | nat the us<br>e/positive<br>nediated | er titrates the antibody controls. | for use in their own     |  |  |
| Product Form Preservative      | a guide only. It is recomm<br>system using appropriate<br>(1) <b>This product require</b><br>Dog<br>Tissue Culture Supernate   | nended tl<br>e negative<br>es heat n  | nat the us<br>e/positive<br>nediated | er titrates the antibody controls. | for use in their own     |  |  |

|                                  | CD45, clustered at the firstInternational Canine Leukocyte Workshop ( <u>Cobbold <i>et al.</i></u> <u>1994</u> ). Mouse anti Dog CD45 antibody, clone CD12.10C12 recognizes a glycoprotein of approximately 200 kDa expressed by cells of haemapoietic origin.   |
|----------------------------------|--|
| Flow Cytometry                   | Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul.  |
| References                       | <ol> <li>Caniatti, M. <i>et al.</i> (1996) Canine lymphoma: immunocytochemical analysis of<br/>fine-needle aspiration biopsy. <u>Vet Pathol. 33 (2): 204-12.</u></li> <li>Cobbold, S. &amp; Metcalfe, S. (1994) Monoclonal antibodies that define canine<br/>homologues of human CD antigens: summary of the First International Canine Leukocyte<br/>Antigen Workshop (CLAW). <u>Tissue Antigens. 43 (3): 137-54.</u></li> <li>Ramos-Vara, J.A. &amp; Miller, M.A. (2007) Immunohistochemical detection of protein gene<br/>product 9.5 (PGP 9.5) in canine epitheliotropic T-cell lymphoma (<i>mycosis fungoides</i>). <u>Vet<br/>Pathol. 44 (1): 74-9.</u></li> <li>Caswell JL, <i>et al.</i> (2005) Graft-versus-host disease in severe combined<br/>immunodeficiency/beige mice administered canine leukocytes. <u>Can J Vet Res. 69(4):<br/>246-52.</u></li> <li>Lane, L.V. <i>et al.</i> (2012) Canine intravascular lymphoma with overt leukemia. <u>Vet Clin</u></li> </ol> |
|                                  | <ul> <li>Pathol. Mar;41(1):84-91.</li> <li>6. Trindade, A.B. <i>et al.</i> (2017) Mesenchymal-like stem cells in canine ovary show high differentiation potential. <u>Cell Prolif. 50(6):e12391.</u></li> <li>7. Tsumagari, S. <i>et al.</i> (2007) Characterization of CD34+ cells from canine umbilical cord blood, bone marrow leukocytes, and peripheral blood by flow cytometric analysis. <u>J Vet Med Sci. 69 (11): 1207-9.</u></li> <li>8. Lombardi, P. <i>et al.</i> (2019) Clinical and Immunological Response in Dogs Naturally Infected by <i>L. infantum</i> Treated with a Nutritional Supplement. <u>Animals (Basel). 9 (8) Jul 30 [Epub ahead of print].</u></li> </ul>  |
| Storage                          | This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.  |
|                                  | Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.  |
| Guarantee                        | 12 months from date of despatch  |
| Health And Safety<br>Information | Material Safety Datasheet documentation #10336 available at:<br>https://www.bio-rad-antibodies.com/SDS/MCA2035S<br>10336   |
| Regulatory                       | For research purposes only   |

## **Related Products**

## **Recommended Secondary Antibodies**

Rabbit Anti Mouse IgG (STAR12...)RPEGoat Anti Mouse IgG IgA IgM (STAR87...)HRP

| Goat Anti Mouse IgG (STAR76)   | RPE  |  |  |  |  |  |  |
|--|--|--|--|--|--|--|--|
| Goat Anti Mouse IgG (STAR70)   | FITC   |  |  |  |  |  |  |
| Goat Anti Mouse IgG (H/L) (STAR117)  | Alk. Phos., DyLight®488, DyLight®550,                          |  |  |  |  |  |  |
|  | DyLight®650, DyLight®680, DyLight®80                           | <u>)0</u> ,  |  |  |  |  |  |
|  | FITC, HRP  |  |  |  |  |  |  |
| Goat Anti Mouse IgG (STAR77)   | HRP  |  |  |  |  |  |  |
| Goat Anti Mouse IgG (Fc) (STAR120)   | FITC, HRP  |  |  |  |  |  |  |
| Rabbit Anti Mouse IgG (STAR13)   | HRP  |  |  |  |  |  |  |
| Rabbit Anti Mouse IgG (STAR9)  | <u>FITC</u>  |  |  |  |  |  |  |
| Recommended Negative Controls  |  |  |  |  |  |  |  |
| MOUSE IgG1 NEGATIVE CONTROL (MCA928)   |  |  |  |  |  |  |  |
| North & South         Tel: +1 800 265 7376         Worldwid           America         Fax: +1 919 878 3751 | de Tel: +44 (0)1865 852 700 Europe<br>Fax: +44 (0)1865 852 739 | Tel: +49 (0) 89 8090 95 21<br>Fax: +49 (0) 89 8090 95 50 |  |  |  |  |  |

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M383318:210513'

Email: antibody\_sales\_uk@bio-rad.com

Email: antibody\_sales\_de@bio-rad.com

Printed on 15 Apr 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint

Email: antibody\_sales\_us@bio-rad.com