

Datasheet: MCA2037S

BATCH NUMBER 168404

Description:	MOUSE ANTI DOG MHC CLASS II MONOMORPHIC
Specificity:	MHC CLASS II MONOMORPHIC
Format:	S/N
Product Type:	Monoclonal Antibody
Clone:	CA2.1C12
Isotype:	IgG1
Quantity:	2 ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation			▪	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Dog
Product Form	Tissue culture supernatant - liquid
Preservative Stabilisers	0.09% sodium azide (NaN ₃)
RRID	AB_323367

Specificity

Mouse anti Dog MHC Class II Monomorphic antibody, clone CA2.1C12 recognizes a monomorphic epitope on canine MHC Class II which was classified at the First Canine Leucocyte Antigen Workshop (CLAW) [Cobbold *et al.* 1992]. In dogs, MHC Class II is expressed by all peripheral blood mononuclear cells. The major histocompatibility complex (MHC) is a cluster of genes that are important in the immune response to infections. In

dogs, this is referred to as the dog leukocyte antigen (DLA) region.

Flow Cytometry Use 10µl of the suggested working dilution to label 10⁶ cells in 100µl

- References**
1. Caniatti, M. *et al.* (1996) Canine lymphoma: immunocytochemical analysis of fine-needle aspiration biopsy. [Vet Pathol. 33: 204-12.](#)
 2. McDonough, S.P. and Moore, P.F. (2000) Clinical, hematologic, and immunophenotypic characterization of canine large granular lymphocytosis. [Vet Pathol. 37: 637-46.](#)
 3. Pumarola, M. *et al.* (2004) Canine inflammatory myopathy: analysis of cellular infiltrates. [Muscle Nerve. 29: 782-9.](#)
 4. Isotani, M. *et al.* (2006) Efficient generation of canine bone marrow-derived dendritic cells. [J Vet Med Sci. 68: 809-14.](#)
 5. Yuasa, K. *et al.* (2007) Injection of a recombinant AAV serotype 2 into canine skeletal muscles evokes strong immune responses against transgene products. [Gene Ther. 14: 1249-60.](#)
 6. Wang, Y.S. *et al.* (2007) Characterization of canine monocyte-derived dendritic cells with phenotypic and functional differentiation. [Can J Vet Res. 71: 165-74.](#)
 7. Liu, C.C. *et al.* (2008) Transient downregulation of monocyte-derived dendritic-cell differentiation, function, and survival during tumoral progression and regression in an in vivo canine model of transmissible venereal tumor. [Cancer Immunol Immunother. 57: 479-91.](#)
 8. Huang, Y.C. *et al.* (2008) CD5-low expression lymphocytes in canine peripheral blood show characteristics of natural killer cells. [J Leukoc Biol. 84: 1501-10.](#)
 9. Ricklin, Gutzwiller. M.E. *et al.* (2010) Comparative analysis of canine monocyte- and bone-marrow-derived dendritic cells. [Vet Res. 41: 40.](#)
 10. Veenhof, E.Z. *et al.* (2011) Characterisation of T cell phenotypes, cytokines and transcription factors in the skin of dogs with cutaneous adverse food reactions. [Vet J. 187 \(3\): 320-4.](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10336 available at: <https://www.bio-rad-antibodies.com/SDS/MCA2037S>
10336

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR77...) [HRP](#)

Rabbit Anti Mouse IgG (STAR12...) [RPE](#)
Goat Anti Mouse IgG IgA IgM (STAR87...) [Alk. Phos.](#), [HRP](#)
Goat Anti Mouse IgG (STAR76...) [RPE](#)
Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)
Rabbit Anti Mouse IgG (STAR13...) [HRP](#)
Rabbit Anti Mouse IgG (STAR9...) [FITC](#)
Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight®488](#), [DyLight®550](#),
[DyLight®650](#), [DyLight®680](#), [DyLight®800](#),
[FITC](#), [HRP](#)
Goat Anti Mouse IgG (STAR70...) [FITC](#)

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	-----------------------------------------------------------------------------------------------------------------------------------------	------------------	-------------------------------------------------------------------------------------------------------------------------------------------------	---------------	-----------------------------------------------------------------------------------------------------------------------------------------------------

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M418614:230427'

Printed on 26 Jun 2024

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)