Datasheet: MCA2041GA BATCH NUMBER 1709

Description:	MOUSE ANTI BOVINE CD172a
Specificity:	CD172a
Other names:	SIRP ALPHA
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	CC149
Isotype:	lgG2b
Quantity:	0.1 mg

Product Details

Applications This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u>.

		Yes	No	Not Determined	Suggested Dilution
	Flow Cytometry				1/100 - 1/200
	Immunohistology - Frozen			•	
	Immunohistology - Paraffin				
	ELISA				
	Immunoprecipitation				
	Western Blotting			•	
	Where this antibody has	not been	tested for	r use in a particular teo	chnique this does not
	necessarily exclude its us				•
	•		•		•
	a guide only. It is recomm	nended that the user titrates the antibo			bay for use in their own
	system using appropriate	negative	e/positive	controls.	
Target Species	Bovine				
Product Form	Purified IgG - liquid				
Preparation	Purified IgG prepared by supernatant	affinity cl	hromatog	raphy on Protein G fro	m tissue culture
Buffer Solution	Phosphate buffered salin	е			

Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
External Database Links	UniProt: <u>O46631</u> Related reagents
	Entrez Gene: <u>327666</u> SIRPA <u>Related reagents</u>
Synonyms	MYD1, PTPNS1, SHPS1, SIRP
Specificity	Mouse anti Bovine CD172a antibody, clone CC149 recognizes bovine CD172a, also known as MyD-1 antigen and SIRPA. CD172a is a ~55 kDa single pass type 1 membrane protein belonging to the family of signal regulatory proteins (SIRP). CD172a has been identified as the receptor for CD47.
	Bovine CD172a is strongly expressed by splenic macrophages, monocytes and a subset of afferent lymph veiled cells (ALVC) and by dendritic cells in the skin.
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
References	 Howard, C.J. <i>et al.</i> (1999) Dendritic cells in cattle: phenotype and function. <u>Vet Immunol Immunopathol. 72 (1-2): 119-24.</u> Price, S.J. & Hope, J.C. (2009) Enhanced secretion of interferon-gamma by bovine gammadelta T cells induced by coculture with <i>Mycobacterium bovis</i>-infected dendritic cells: evidence for reciprocal activating signals. <u>Immunology. 126:201-8</u> Waters, W.R. (2009) Signal regulatory protein alpha (SIRPalpha) cells in the adaptive response to ESAT-6/CFP-10 protein of tuberculous mycobacteria. <u>PLoS One. 4: e6414.</u> Brackenbury, L.S. <i>et al.</i> (2005) Identification of a cell population that produces alpha/beta interferon <i>in vitro</i> and <i>in vivo</i> in response to noncytopathic bovine viral diarrhea virus. <u>J Virol. 79: 7738-44.</u> Smith, R. <i>et al.</i> (2003) A novel MyD-1 (SIRP-1alpha) signaling pathway that inhibits LPS-induced TNFalpha production by monocytes. <u>Blood.102: 2532-40.</u> Jensen, K. <i>et al.</i> (2014) Comparison of small interfering RNA (siRNA) delivery into bovine monocyte-derived macrophages by transfection and electroporation. <u>Vet Immunol Immunopathol. 158 : 224-32.</u> Tahoun, A. <i>et al.</i> (2015) Functional analysis of bovine TLR5 and association with IgA responses of cattle following systemic immunisation with H7 flagella. <u>Vet Res. 46: 9.</u> Hussen J <i>et al.</i> (2014) The chemokine CCL5 induces selective migration of bovine classical monocytes and drives their differentiation into LPS-hyporesponsive macrophages <i>in vitro</i>. <u>Dev Comp Immunol. 47 (2): 169-77.</u> Eger, M. <i>et al.</i> (2015) An <i>in vitro</i> model to assess the immunosuppressive effect of tick saliva on the mobilization of inflammatory monocyte-derived cells. <u>Vet Res. 46 (1)</u>:

	 <u>117.</u> 11. Pridans, C. <i>et al.</i> (2016) A Csf1r-EGFP Transgene Provides Monocyte Subsets in Sheep. <u>J Immunol. 197 (6): 2297-305.</u> 12. Herry, V. <i>et al.</i> (2017) Local immunization impacts the response <i>Escherichia coli</i> mastitis. <u>Sci Rep. 7 (1): 3441.</u> 	
Storage	Store at +4°C or at -20°C if preferred.	
	This product should be stored undiluted.	
	Storage in frost free freezers is not recommended. Avoid repea as this may denature the antibody. Should this product contain recommend microcentrifugation before use.	• •
Guarantee	12 months from date of despatch	
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA2041GA 10040	
Regulatory	For research purposes only	

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12)	RPE	
Goat Anti Mouse IgG IgA IgM (STAR87) <u>HRP</u>		
Goat Anti Mouse IgG (STAR76)	RPE	
Goat Anti Mouse IgG (STAR70)	<u>FITC</u>	
Goat Anti Mouse IgG (H/L) (STAR117)	Alk. Phos., DyLight®488, DyLight®550,	
	DyLight®650, DyLight®680, DyLight®800,	
	FITC, HRP	
Rabbit Anti Mouse IgG (STAR9)	<u>FITC</u>	
Goat Anti Mouse IgG (STAR77)	HRP	
Goat Anti Mouse IgG (Fc) (STAR120)	FITC, HRP	
Rabbit Anti Mouse IgG (STAR13)	HRP	
Recommended Negative Controls		

MOUSE IgG2b NEGATIVE CONTROL (MCA691)

Recommended Useful Reagents

BOVINE DENDRITIC CELL GROWTH KIT (PBP014KZZ) BOVINE DENDRITIC CELL GROWTH KIT (PBP015KZZ) MOUSE ANTI BOVINE CD205:FITC (MCA1651F) MOUSE ANTI BOVINE CD205 (MCA1651GA)

North & South	Tel: +1 800 265 7376
America	Fax: +1 919 878 3751

Worldwide

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M366086:200529'

Printed on 21 Mar 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint