

## Datasheet: MCA2042GA

<b>Description:</b>	MOUSE ANTI BOVINE CD63
<b>Specificity:</b>	CD63
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	CC25
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	0.1 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/10 - 1/25
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting			▪	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Bovine
<b>Product Form</b>	Purified IgG - liquid
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.09% sodium azide (NaN <sub>3</sub> )
<b>Carrier Free</b>	Yes

Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Bovine PBMC
External Database Links	<p><b>UniProt:</b>  <a href="#">Q9XSK2</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">404156</a>    CD63    <a href="#">Related reagents</a></p>
Specificity	<p><b>Mouse anti Bovine CD63 antibody, clone CC25</b> recognizes the bovine homologue of human CD63, a 237 amino acid multipass transmembrane glycoprotein and member of the tetraspanin TM4SF protein family with a predicted molecular weight of ~26 kDa. CD63 is also known as lysosome associated membrane glycoprotein 3 or LAMP-3.</p> <p>CD63 along with other TM4SF members including CD9, CD61 and CD151 can form specific interactions with phosphoinositide 4-kinase, suggesting a role for CD63 in the recruitment of phosphoinositide 4-kinase to specific membrane sites (<a href="#">Yauch and Hemler 2000</a>). CD63 is expressed on the cell surface of platelets and basophils, along with activated macrophages, monocytes and granulocytes.</p> <p>Mouse anti Bovine CD63, clone CC25 acts as a specific marker for bovine lysosomes and has been used for the identification and quantitation of phagosome-lysosome fusion in models of bacterial insult (<a href="#">Souza et al. 2013</a>).</p>
Flow Cytometry	Use 10µl of the suggested working dilution to label 10 <sup>6</sup> cells in 100µl
References	<ol style="list-style-type: none"> <li>1. Brooke, G.P. <i>et al.</i> (1999) Molecular cloning of cattle CD63 and evidence for high level expression on subpopulations of dendritic cells. <a href="#">Immunogenetics. 49 (9): 812-4.</a></li> <li>2. Colino, J. and Snapper, C.M. (2006) Exosomes from bone marrow dendritic cells pulsed with diphtheria toxoid preferentially induce type 1 antigen-specific IgG responses in naive recipients in the absence of free antigen. <a href="#">J Immunol. 177: 3757-62.</a></li> <li>3. Souza, C.D. <i>et al.</i> (2007) Role of the mitogen-activated protein kinase pathway in the differential response of bovine monocytes to <i>Mycobacterium avium</i> subsp. paratuberculosis and <i>Mycobacterium avium</i> subsp. <i>avium</i>. <a href="#">Microbes Infect. 9: 1545-52.</a></li> <li>4. Weiss, D.J. <i>et al.</i> (2008) Bovine monocyte TLR2 receptors differentially regulate the intracellular fate of <i>Mycobacterium avium</i> subsp. paratuberculosis and <i>Mycobacterium avium</i> subsp. <i>avium</i>. <a href="#">J Leukoc Biol. 83: 48-55.</a></li> <li>5. Souza, C. <i>et al.</i> (2013) Mannosylated lipoarabinomannans from <i>Mycobacterium avium</i> subsp. Paratuberculosis alters the inflammatory response by bovine macrophages and suppresses killing of mycobacterium avium subsp. Avium organisms. <a href="#">PLoS One 8: e75924.</a></li> <li>6. Wolf, T. <i>et al.</i> (2015) The Intestinal Transport of Bovine Milk Exosomes Is Mediated by Endocytosis in Human Colon Carcinoma Caco-2 Cells and Rat Small Intestinal IEC-6 Cells. <a href="#">J Nutr. 145 (10): 2201-6.</a></li> <li>7. Carretta MD <i>et al.</i> (2016) Butyric acid stimulates bovine neutrophil functions and</li> </ol>

- potentiates the effect of platelet activating factor. [Vet Immunol Immunopathol. 176: 18-27.](#)
8. Kusuma, R.J. *et al.* (2016) Human vascular endothelial cells transport foreign exosomes from cow's milk by endocytosis. [Am J Physiol Cell Physiol. 310 \(10\): C800-7.](#)
9. Mobley, C.B. *et al.* (2017) Whey protein-derived exosomes increase protein synthesis and hypertrophy in C<sub>2</sub>-C<sub>12</sub> myotubes. [J Dairy Sci. 100 \(1\): 48-64.](#)
10. Gillan, V. *et al.* (2019) Characterisation of infection associated microRNA and protein cargo in extracellular vesicles of *Theileria annulata*. infected leukocytes. [Cell Microbiol. 21 \(1\): e12969.](#)
11. Fiorenza, M.F. *et al.* (2021) Neutrophils recognize and amplify IFNT signals derived from day 7 bovine embryo for stimulation of ISGs expression *in vitro*.: A possible implication for the early maternal recognition of pregnancy. [Biochem Biophys Res Commun. 553: 37-43.](#)

<b>Storage</b>	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.  Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.
<b>Guarantee</b>	12 months from date of despatch
<b>Health And Safety Information</b>	Material Safety Datasheet documentation #10040 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA2042GA">https://www.bio-rad-antibodies.com/SDS/MCA2042GA</a> 10040
<b>Regulatory</b>	For research purposes only

## Related Products

### Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR77...)	<a href="#">HRP</a>
Rabbit Anti Mouse IgG (STAR12...)	<a href="#">RPE</a>
Goat Anti Mouse IgG IgA IgM (STAR87...)	<a href="#">Alk. Phos.</a> , <a href="#">HRP</a>
Goat Anti Mouse IgG (STAR76...)	<a href="#">RPE</a>
Goat Anti Mouse IgG (Fc) (STAR120...)	<a href="#">FITC</a> , <a href="#">HRP</a>
Rabbit Anti Mouse IgG (STAR13...)	<a href="#">HRP</a>
Rabbit Anti Mouse IgG (STAR9...)	<a href="#">FITC</a>
Goat Anti Mouse IgG (H/L) (STAR117...)	<a href="#">Alk. Phos.</a> , <a href="#">DyLight®488</a> , <a href="#">DyLight®550</a> , <a href="#">DyLight®650</a> , <a href="#">DyLight®680</a> , <a href="#">DyLight®800</a> , <a href="#">FITC</a> , <a href="#">HRP</a>
Goat Anti Mouse IgG (STAR70...)	<a href="#">FITC</a>

### Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a> 'M412574:221114'	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

**Printed on 10 Jan 2025**

---

© 2025 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)