

## Datasheet: MCA2045GA

**BATCH NUMBER 159483**

<b>Description:</b>	MOUSE ANTI HUMAN CD177
<b>Specificity:</b>	CD177
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	MEM-166
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	0.1 mg

### Product Details

#### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/50 - 1/100
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting (1)	▪			Non-reducing conditions

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

**(1)MEM-166 recognizes CD177 under non-reducing conditions**

<b>Target Species</b>	Human
<b>Species Cross Reactivity</b>	<p>Reacts with: Rhesus Monkey</p> <p><b>N.B.</b> Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.</p>
<b>Product Form</b>	Purified IgG - liquid
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein A

<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.09% Sodium Azide
<b>Approx. Protein Concentrations</b>	IgG concentration 1.0 mg/ml
<b>Immunogen</b>	Human granulocytes.
<b>External Database Links</b>	<p><b>UniProt:</b>  <a href="#">Q8N6Q3</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">57126</a>    CD177    <a href="#">Related reagents</a></p>
<b>Synonyms</b>	NB1, PRV1
<b>RRID</b>	AB_324462
<b>Specificity</b>	<b>Mouse anti Human CD177 antibody, clone MEM-166</b> recognizes human CD177 (neutrophil glycoprotein NB1). The neutrophil NB1 antigen is expressed by 97% of the caucasian population. Antibodies against NB1 have been implicated in the pathology of neonatal alloimmune neutropenia ( <a href="#">Lalezari et al. 1971</a> ).
<b>Flow Cytometry</b>	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul.
<b>References</b>	<ol style="list-style-type: none"> <li>Caruccio, L. <i>et al.</i> (2003) Expression of human neutrophil antigen-2a (NB1) is increased in pregnancy. <a href="#">Transfusion. 43 (3): 357-63.</a></li> <li>Kissel, K. <i>et al.</i> (2002) Molecular basis of NB1 (HNA-2a, CD177) deficiency. <a href="#">Blood. 99 (11): 4231-3.</a></li> <li>Jerke, U. <i>et al.</i> (2011) Complement receptor Mac-1 is an adaptor for NB1 (CD177)-mediated PR3-ANCA neutrophil activation. <a href="#">J Biol Chem. 286 (9): 7070-81.</a></li> <li>Dillon, M. <i>et al.</i> (2008) Expression of the GPI-anchored receptor Prv-1 enhances thrombopoietin and IL-3-induced proliferation in hematopoietic cell lines. <a href="#">Leuk Res. 32: 811-9.</a></li> <li>Gabillet, J. <i>et al.</i> (2012) Proteinase 3, the autoantigen in granulomatosis with polyangiitis, associates with calreticulin on apoptotic neutrophils, impairs macrophage phagocytosis, and promotes inflammation. <a href="#">J Immunol. 189: 2574-83.</a></li> <li>Drewniak, A. <i>et al.</i> (2008) Granulocyte concentrates: prolonged functional capacity during storage in the presence of phenotypic changes. <a href="#">Haematologica. 93:1058-67.</a></li> <li>Drewniak, A. <i>et al.</i> (2009) Changes in gene expression of granulocytes during in vivo granulocyte colony-stimulating factor/dexamethasone mobilization for transfusion purposes. <a href="#">Blood. 113: 5979-98.</a></li> <li>Sachs, U.J. <i>et al.</i> (2007) The neutrophil-specific antigen CD177 is a counter-receptor for platelet endothelial cell adhesion molecule-1 (CD31). <a href="#">J Biol Chem. 282: 23603-12.</a></li> <li>Johansson, Å.C. <i>et al.</i> (2016) Impaired phagocytosis and reactive oxygen species production in phagocytes is associated with systemic vasculitis. <a href="#">Arthritis Res Ther. 18 (1):</a></li> </ol>

[92.](#)

10. Nishimura, M. *et al.* (2007) Detection of anti-CD32 alloantibody in donor plasma implicated in development of transfusion-related acute lung injury. [Cell Biochem Funct. 25 \(2\): 179-83.](#)

11. Pliyev, B.K. & Menshikov, M. (2012) Comparative evaluation of the role of the adhesion molecule CD177 in neutrophil interactions with platelets and endothelium. [Eur J Haematol. 89 \(3\): 236-44.](#)

12. Onodera, R. *et al.* (2017) Anti-human neutrophil antigen-1a, -1b, and -2 antibodies in neonates and children with immune neutropenias analyzed by extracted granulocyte antigen immunofluorescence assay. [Transfusion. 57 \(11\): 2586-94.](#)

13. Bayat, B. *et al.* (2021) Transfusion of Target Antigens to Pre-Immunized Recipients: A New Mechanism in Transfusion-Related Acute Lung Injury. [Blood Adv. bloodadvances.2020003843.](#)

---

**Storage** This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

---

**Guarantee** 12 months from date of despatch

---

**Health And Safety Information** Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/MCA2045GA>  
10040

---

**Regulatory** For research purposes only

---

## Related Products

### Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...) [RPE](#)  
Goat Anti Mouse IgG IgA IgM (STAR87...) [HRP](#)  
Goat Anti Mouse IgG (STAR76...) [RPE](#)  
Rabbit Anti Mouse IgG (STAR13...) [HRP](#)  
Goat Anti Mouse IgG (STAR70...) [FITC](#)  
Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight@488](#), [DyLight@550](#),  
[DyLight@650](#), [DyLight@680](#), [DyLight@800](#),  
[FITC](#), [HRP](#)  
Rabbit Anti Mouse IgG (STAR9...) [FITC](#)  
Goat Anti Mouse IgG (STAR77...) [HRP](#)  
Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)

### Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

**North & South** Tel: +1 800 265 7376

**Worldwide**

Tel: +44 (0)1865 852 700

**Europe**

Tel: +49 (0) 89 8090 95 21

To

**America** Fax: +1 919 878 3751

Fax: +44 (0)1865 852 739

Fax: +49 (0) 89 8090 95 50

find a

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://bio-rad-antibodies.com/datasheets)

'M383639:210513'

**Printed on 18 Jan 2024**

---

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)