

Datasheet: MCA2045PE

BATCH NUMBER 153454

Description:	MOUSE ANTI HUMAN CD177:RPE
Specificity:	CD177
Format:	RPE
Product Type:	Monoclonal Antibody
Clone:	MEM-166
Isotype:	lgG1
Quantity:	100 TESTS

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	•			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human			
Species Cross	Reacts with: Rhesu	s Monkey		
Reactivity	reactivity is derived	from testing within our I	ions may vary between species. Cross aboratories, peer-reviewed publications ors. Please refer to references indicated	
Product Form	Purified IgG conjug	ated to R. Phycoerythrir	(RPE) - lyophilized	
Reconstitution	Reconstitute with 1	ml distilled water.		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)	
	RPE 488nm laser	496	578	
Preparation	Purified IgG prepare	ed by affinity chromatog	raphy on Protein G.	
Buffer Solution	Phosphate buffered	l saline		

Preservative
Stabilisers

0.09% Sodium Azide

1% Bovine Serum Albumin

5% Sucrose

Immunogen

Human granulocytes.

External Database

Links

UniProt:

Q8N6Q3 Related reagents

Entrez Gene:

57126 CD177 Related reagents

Synonyms

NB1, PRV1

RRID

AB 323430

Specificity

Mouse anti Human CD177 antibody, clone MEM-166 recognizes human CD177 (neutrophil glycoprotein NB1). The neutrophil NB1 antigen is expressed by 97% of the caucasian population. Antibodies against NB1 have been implicated in the pathology of neonatal alloimmune neutropenia (<u>Lalezari et al. 1971</u>).

Flow Cytometry

Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul.

References

- 1. Caruccio, L. *et al.* (2003) Expression of human neutrophil antigen-2a (NB1) is increased in pregnancy. Transfusion. 43 (3): 357-63.
- 2. Kissel, K. *et al.* (2002) Molecular basis of NB1 (HNA-2a, CD177) deficiency. <u>Blood. 99</u> (11): 4231-3.
- 3. Jerke, U. *et al.* (2011) Complement receptor Mac-1 is an adaptor for NB1 (CD177)-mediated PR3-ANCA neutrophil activation. <u>J Biol Chem. 286 (9): 7070-81.</u>
- 4. Dillon, M. *et al.* (2008) Expression of the GPI-anchored receptor Prv-1 enhances thrombopoietin and IL-3-induced proliferation in hematopoietic cell lines. <u>Leuk Res. 32: 811-9.</u>
- 5. Gabillet, J. *et al.* (2012) Proteinase 3, the autoantigen in granulomatosis with polyangiitis, associates with calreticulin on apoptotic neutrophils, impairs macrophage phagocytosis, and promotes inflammation. J Immunol. 189: 2574-83.
- 6. Drewniak, A. *et al.* (2008) Granulocyte concentrates: prolonged functional capacity during storage in the presence of phenotypic changes. <u>Haematologica</u>. 93:1058-67.
- 7. Drewniak, A. *et al.* (2009) Changes in gene expression of granulocytes during in vivo granulocyte colony-stimulating factor/dexamethasone mobilization for transfusion purposes. Blood. 113: 5979-98.
- 8. Sachs, U.J. *et al.* (2007) The neutrophil-specific antigen CD177 is a counter-receptor for platelet endothelial cell adhesion molecule-1 (CD31). J Biol Chem. 282: 23603-12.
- 9. Johansson, Å.C. *et al.* (2016) Impaired phagocytosis and reactive oxygen species production in phagocytes is associated with systemic vasculitis. <u>Arthritis Res Ther. 18 (1):</u> 92.
- 10. Nishimura, M. *et al.* (2007) Detection of anti-CD32 alloantibody in donor plasma implicated in development of transfusion-related acute lung injury. Cell Biochem Funct. 25

(2): 179-83.

- 11. Pliyev, B.K. & Menshikov, M. (2012) Comparative evaluation of the role of the adhesion molecule CD177 in neutrophil interactions with platelets and endothelium. Eur J Haematol. 89 (3): 236-44.
- 12. Onodera, R. et al. (2017) Anti-human neutrophil antigen-1a, -1b, and -2 antibodies in neonates and children with immune neutropenias analyzed by extracted granulocyte antigen immunofluorescence assay. Transfusion. 57 (11): 2586-94.

Storage

Store at +4°C.

DO NOT FREEZE.

This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.

Information
Health And Safety

Guarantee

12 months from date of despatch

Material Safety Datasheet documentation #20487 available at:

https://www.bio-rad-antibodies.com/SDS/MCA2045PE

20487

Regulatory

For research purposes only

Related Products

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL:RPE (MCA928PE)

Email: antibody_sales_us@bio-rad.com

Recommended Useful Reagents

HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

America

North & South Tel: +1 800 265 7376 Fax: +1 919 878 3751 Worldwide

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M375408:210104'

Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint