

Datasheet: MCA2045PE

Description:	MOUSE ANTI HUMAN CD177:RPE
Specificity:	CD177
Format:	RPE
Product Type:	Monoclonal Antibody
Clone:	MEM-166
Isotype:	IgG1
Quantity:	100 TESTS

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human		
Species Cross Reactivity	Reacts with: Rhesus Monkey N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.		
Product Form	Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilized		
Reconstitution	Reconstitute with 1 ml distilled water.		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	RPE 488nm laser	496	578
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant		

Buffer Solution	Phosphate buffered saline
Preservative	0.09% sodium azide (NaN ₃)
Stabilisers	1% bovine serum albumin 5% sucrose
Immunogen	Human granulocytes.
External Database Links	UniProt: Q8N6Q3 Related reagents Entrez Gene: 57126 CD177 Related reagents
Synonyms	NB1, PRV1
RRID	AB_323430
Specificity	Mouse anti Human CD177 antibody, clone MEM-166 recognizes human CD177 (neutrophil glycoprotein NB1). The neutrophil NB1 antigen is expressed by 97% of the caucasian population. Antibodies against NB1 have been implicated in the pathology of neonatal alloimmune neutropenia (Lalezari et al. 1971).
Flow Cytometry	Use 10µl of the suggested working dilution to label 10 ⁶ cells in 100µl
References	1. Kissel, K. <i>et al.</i> (2002) Molecular basis of NB1 (HNA-2a, CD177) deficiency. Blood. 99 (11): 4231-3. 2. Caruccio, L. <i>et al.</i> (2003) Expression of human neutrophil antigen-2a (NB1) is increased in pregnancy. Transfusion. 43 (3): 357-63. 3. Nishimura, M. <i>et al.</i> (2007) Detection of anti-CD32 alloantibody in donor plasma implicated in development of transfusion-related acute lung injury. Cell Biochem Funct. 25 (2): 179-83. 4. Sachs, U.J. <i>et al.</i> (2007) The neutrophil-specific antigen CD177 is a counter-receptor for platelet endothelial cell adhesion molecule-1 (CD31). J Biol Chem. 282: 23603-12. 5. Drewniak, A. <i>et al.</i> (2008) Granulocyte concentrates: prolonged functional capacity during storage in the presence of phenotypic changes. Haematologica. 93:1058-67. 6. Dillon, M. <i>et al.</i> (2008) Expression of the GPI-anchored receptor Prv-1 enhances thrombopoietin and IL-3-induced proliferation in hematopoietic cell lines. Leuk Res. 32: 811-9. 7. Drewniak, A. <i>et al.</i> (2009) Changes in gene expression of granulocytes during in vivo granulocyte colony-stimulating factor/dexamethasone mobilization for transfusion purposes. Blood. 113: 5979-98. 8. Jerke, U. <i>et al.</i> (2011) Complement receptor Mac-1 is an adaptor for NB1 (CD177)-mediated PR3-ANCA neutrophil activation. J Biol Chem. 286 (9): 7070-81. 9. Pliyev, B.K. & Menshikov, M. (2012) Comparative evaluation of the role of the adhesion molecule CD177 in neutrophil interactions with platelets and endothelium. Eur J Haematol. 89 (3): 236-44.

10. Gabillet, J. *et al.* (2012) Proteinase 3, the autoantigen in granulomatosis with polyangiitis, associates with calreticulin on apoptotic neutrophils, impairs macrophage phagocytosis, and promotes inflammation. [J Immunol. 189: 2574-83.](#)
11. Johansson, Å.C. *et al.* (2016) Impaired phagocytosis and reactive oxygen species production in phagocytes is associated with systemic vasculitis. [Arthritis Res Ther. 18 \(1\): 92.](#)
12. Onodera, R. *et al.* (2017) Anti-human neutrophil antigen-1a, -1b, and -2 antibodies in neonates and children with immune neutropenias analyzed by extracted granulocyte antigen immunofluorescence assay. [Transfusion. 57 \(11\): 2586-94.](#)
13. Bayat, B. *et al.* (2021) Transfusion of Target Antigens to Pre-Immunized Recipients: A New Mechanism in Transfusion-Related Acute Lung Injury. [Blood Adv. bloodadvances.2020003843.](#)

Storage	This product is shipped at ambient temperature. Store at +4°C. DO NOT FREEZE. This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #20487 available at: https://www.bio-rad-antibodies.com/SDS/MCA2045PE 20487
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:RPE \(MCA928PE\)](#)

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M440362:250523'

Printed on 23 May 2025