

Datasheet: MCA2046GA

BATCH NUMBER 174622

Description:	MOUSE ANTI HUMAN CD56
Specificity:	CD56
Other names:	NCAM-1
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	MEM-188
Isotype:	IgG2a
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/25 - 1/50
Immunohistology - Frozen (1)	▪			
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting (2)	▪			Non-reducing conditions

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1)The epitope recognised by this antibody is reported to be sensitive to formaldehyde fixation and tissue processing. Bio-Rad recommends the use of acetone fixation for frozen sections.

(2) MEM-188 recognizes CD56 under non-reducing conditions.

Target Species	Human
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant

Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% sodium azide (NaN ₃)
Approx. Protein Concentrations	IgG concentration 1.0mg/ml
External Database Links	UniProt: P13591 Related reagents Entrez Gene: 4684 NCAM1 Related reagents
Synonyms	NCAM
RRID	AB_324588
Specificity	Mouse anti Human CD56 antibody, clone MEM-188 recognizes CD56, a 140kDa cell surface glycoprotein expressed primarily by natural killer cells and a subset of T cells in peripheral blood.
Flow Cytometry	Use 10µl of the suggested working dilution to label 10 ⁶ cells in 100µl
References	- Costa, P.<i>et al</i> (1997) CD56 Workshop Panel Report. In: Leucocyte typing VI. White cell differentiation antigens. Garland Publishing Inc. New York and London. 271-2. - Prager, E. <i>et al</i>. (2001) Induction of hyporesponsiveness and impaired T lymphocyte activation by the CD31 receptor:ligand pathway in T cells. J Immunol. 166 (4): 2364-71. - Staffler G <i>et al</i>. (2003) Selective inhibition of T cell activation via CD147 through novel modulation of lipid rafts. J Immunol. 171 (4): 1707-14. - Crisan, M. <i>et al</i>. (2008) A perivascular origin for mesenchymal stem cells in multiple human organs. Cell Stem Cell. 3: 301-13. - Newman, K.C. <i>et al</i>. (2006) Cross-talk with myeloid accessory cells regulates human natural killer cell interferon-gamma responses to malaria. PLoS Pathog. 2: e118. - Apps, R. <i>et al</i>. (2011) Ex vivo functional responses to HLA-G differ between blood and decidual NK cells. Mol Hum Reprod. 17: 577-86. - Aspinall, A.I. <i>et al</i>. (2010) CX(3)CR1 and vascular adhesion protein-1-dependent recruitment of CD16(+) monocytes across human liver sinusoidal endothelium. Hepatology. 51: 2030-9. - Rodriguez JM <i>et al</i>. (2015) PyNTTTTGT and CpG Immunostimulatory Oligonucleotides: Effect on Granulocyte/Monocyte Colony-Stimulating Factor (GM-CSF) Secretion by Human CD56+ (NK and NKT) Cells. PLoS One. 10 (2): e0117484. - Cabioglu, N. <i>et al</i>. (2025) The Prognostic and Therapeutic Significance of T-Helper-1 Cell Response in Patients with Severe Systemic Inflammatory Response or Sepsis Archives of Clinical and Biomedical Research. 9 (5): 403-411.
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for

short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA2046GA
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG IgA IgM (STAR87...)	HRP
Goat Anti Mouse IgG (STAR70...)	FITC
Goat Anti Mouse IgG (STAR77...)	HRP
Goat Anti Mouse IgG (STAR76...)	RPE
Rabbit Anti Mouse IgG (STAR12...)	RPE
Rabbit Anti Mouse IgG (STAR13...)	HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight®488 , DyLight®550 , DyLight®650 , DyLight®680 , DyLight®800 , FITC , HRP

Recommended Negative Controls

[MOUSE IgG2a NEGATIVE CONTROL \(MCA929\)](#)

Product inquiries: www.bio-rad-antibodies.com/technical-support

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M412602:221114'

Printed on 21 Aug 2026