

Datasheet: MCA2047F BATCH NUMBER 164836

Description:	MOUSE ANTI TUBULIN BETA 3:FITC
Specificity:	TUBULIN BETA 3
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	TU-20
lsotype:	lgG1
Quantity:	0.1 mg

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal					
	communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-</u>					
	rad-antibodies.com/pr					
		Yes No	Not Determined	Suggested Dilution		
	Immunofluorescence	•		1/10 - 1/50		
	Where this product has not been tested for use in a particular technique this does not					
	necessarily exclude its use in such procedures. Suggested working dilutions are given as					
a guide only. It is recommended that the user titrates the product for use in the system using appropriate negative/positive controls.				for use in their own		
Target Species	Human					
Species Cross Reactivity	Reacts with: Mouse, Baboon, Rat, Hamster, Pig, Bovine N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.					
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid					
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)			
	FITC	490	525			
Preparation	Purified IgG prepared by DEAE chromatography from tissue culture supernatant					
Buffer Solution	Phosphate buffered sa	aline				

Preservative Stabilisers	0.09% sodium azide (NaN ₃)		
Approx. Protein Concentrations	IgG concentration 1.0mg/ml		
Immunogen	Synthetic peptide, ESESQGPK, corresponding to amino acids 441-448 of human class beta tubulin coupled to Keyhole Limpet Hemocyanin (KLH).	s III	
	This sequence is widely conserved across species.		
External Database Links	UniProt:		
	Q13509 Related reagents		
	Entrez Gene:		
	10381 TUBB3 Related reagents		
Synonyms	TUBB4		
RRID	AB_2210681		
Specificity	Mouse anti Tubulin beta 3 antibody, clone TU-20 recognizes class III beta-tubulin, restricted to neuronal tissue (Leandro-García 2010; Katsetos 2003).		
	Mouse anti Tubulin beta 3 antibody, clone TU-20 has been used to investigate tumors neuronal origin (<u>Jirásek 2002</u>) including neuroblastoma (<u>Prasannan 2000</u>) and ganglioneuroma (<u>Dráberová 1998</u>). Class III beta tubulin is highly expressed in tumors neuronal origin rather than in non-neuronal tumors (<u>Person 2017</u>).		
References	1. Dráberová, E. <i>et al.</i> (1998) Expression of class III beta-tubulin in normal and neopla human tissues. <u>Histochem Cell Biol. 109 (3): 231-9.</u>	stic	
	 Pěknicová, J. <i>et al.</i> (2001) Differential subcellular distribution of tubulin epitopes in b 	oar	
	spermatozoa: recognition of class III beta-tubulin epitope in sperm tail. <u>Biol Reprod. 65</u> 672-9.	<u>;</u>	
	3. Nicot, A. and DiCicco-Bloom, E. (2001) Regulation of neuroblast mitosis is determin	ed	
	by PACAP receptor isoform expression. Proc Natl Acad Sci U S A. 98: 4758-63.		
	 Carey, R.G. <i>et al.</i> (2002) Pituitary adenylate cyclase activating polypeptide anti-mitogenic signaling in cerebral cortical progenitors is regulated by p57Kip2-dependence 	dont	
	CDK2 activity. <u>J Neurosci. 22 (5): 1583-91.</u>	uem	
	5. Vedin, V. et al. (2010) Organization of the chemosensory neuroepithelium of the		
	vomeronasal organ of the Scandinavian moose <i>Alces alces</i> . <u>Brain Res. 1306: 53-61.</u>		
	 Knerlich-Lukoschus, F. et al. (2010) Chemokine expression in the white matter spina cord precursor niche after force-defined spinal cord contusion injuries in adult rats. <u>Glia</u> 		
	<u>58 (8): 916-31.</u>	<u>.</u>	
	7. Huang, C.L. <i>et al.</i> (2010) Expression of ERCC1 and class III β -tubulin is associated		
	7. Huang, C.L. <i>et al.</i> (2010) Expression of ERCC1 and class III β -tubulin is associated with the survival of resected stage III non-small cell lung cancer patients treated with		
	7. Huang, C.L. <i>et al.</i> (2010) Expression of ERCC1 and class III β -tubulin is associated		

	human glioma cells and mediates antiapoptotic effects. <u>Cancer Res. 70: 3299-308.</u> 9. Zhu, G. <i>et al.</i> (2012) Effects of neurotrophin-3 on the differentiation of neural stem cells			
	into neurons and oligodendrocytes. <u>Neural Regen Res. 7 (19): 1483-7.</u>			
	10. Rosito, M. et al. (2012) CXCL16 Orchestrates Adenosine A3 Receptor and			
	MCP-1/CCL2 Activity to Protect Neurons from Excitotoxic Cell Death in the CNS. \underline{J}			
	<u>Neurosci. 32: 3154-63.</u>			
	11. Volkov, V.A. et al. (2013) Long tethers provide high-force coupling of the Dam1 ring to			
	shortening microtubules. Proc Natl Acad Sci U S A. 110 (19): 7708-13.			
	12. Alexiou, G.A. et al. (2013) Supratentorial ependymomas in children: Analysis of nine			
	cases. <u>J Pediatr Neurosci. 8 (1): 15-8.</u>			
	13. Tarasovetc, E.V. et al. (2021) Permitted and restricted steps of human kinetochore			
	assembly in mitotic cell extracts. Mol Biol Cell. : mbcE20070461.			
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in			
	frost-free freezers is not recommended. This product is photosensitive and should be protected from light.			
Guarantee	Guaranteed for 12 months from the date of despatch or until the date of expiry, whichever comes first. Please see label for expiry date.			
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA2047F 10040			
Regulatory	For research purposes only			

Related Products

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL: FITC (MCA928F)

North & South	Tel: +1 800 265 7376
America	Fax: +1 919 878 3751
	Email: antibody_sales_u

7376 Worldwide 3751 sales_us@bio-rad.com Tel: +44 (0)1865 852 700 **Europe** Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M412605:221114'

Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint