

# Datasheet: MCA2048F BATCH NUMBER 1705

Description:	MOUSE ANTI HUMAN CD222:FITC
Specificity:	CD222
Other names:	IGF-2 RECEPTOR
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	MEM-238
Isotype:	lgG1
Quantity:	0.1 mg

# **Product Details**

Applications This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u>.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry (1)				neat
Immunohistology - Frozen			•	
Immunohistology - Paraffin			•	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1)Membrane permeabilisation is required for this application. Bio-Rad recommends the use of Leucoperm<sup>™</sup> (Product Code <u>BUF09</u>) for this purpose.

Target Species	Human		
Species Cross Reactivity	reactivity is derive	activity and working conditi ed from testing within our land nications from the originato	ons may vary between species. Cross aboratories, peer-reviewed publication ors. Please refer to references indicate
Product Form	Purified IgG - liqu	lid	
Product Form Max Ex/Em	Purified IgG - liqu Fluorophore	uid Excitation Max (nm)	Emission Max (nm)

<ul> <li>kDa transmembrane protein originally identified as the IGF II receptor. CD222 is ubiquitously expressed and is involved with internalization of a variety of ligands.</li> <li>Mouse anti Human CD222 antibody, clone MEM-238 recognizes an epitope located in the region between extracellular domains 2 and 5 (aa 192-697) of CD222 (Roberts <i>et al.</i> 2010).</li> <li>CD222 is primarily expressed intracellularly with a small percentage of molecules being located at the cell surface (5 - 10%).</li> <li>Flow Cytometry Use 10ul of the suggested working dilution to label 1x10<sup>6</sup> cells in 100ul.</li> <li>References 1. Roberts, R.C. <i>et al.</i> (2010) Mistargeting of SH3TC2 away from the recycling endoson causes Charcot-Marie-Tooth disease type 4C. Hum Mol Genet. 19: 1009-18.</li> <li>2. Adachi, A. <i>et al.</i> (2010) Golgi-associated GSK3beta regulates the sorting process of post-Golgi membrane trafficking. J Cell Sci. 123: 3215-25.</li> <li>3. Godar, S. <i>et al.</i> (2002) CD222 (Mannose-6 phosphate / insulin-like growth factor II-receptor) Summary and Workshop Report. In Leucocyte Typing VII: White Cell Differentiation Antigens. Edited by Mason, D. <i>et al.</i> Oxford University Press. pp482-485.</li> <li>4. Rezgui, D. <i>et al.</i> (2009) Structure and function of the human Gly1619Arg polymorphis of M6P/IGF2R domain 11 implicated in IGF2 dependent growth. J Mol Endocrinol. 42: 341-56.</li> </ul>				
Preservative       0.09% Sodium Azide         1%       Bovine Serum Albumin         Approx. Protein       IgG concentration 0.1 mg/ml         Immunogen       Recombinant vaccinia virus containing CD222.         External Database       UniProt:         Links       P11717         Related reagents         Entrez Gene:       3482         3482       IGF2R         RRID       AB_323453         Specificity       Mouse anti Human CD222 antibody, clone MEM-238 recognizes human CD222, a 22         KDa transmembrane protein originally identified as the IGF II receptor. CD222 is ubiquitously expressed and is involved with internalization of a variety of ligands.         Mouse anti Human CD222 antibody, clone MEM-238 recognizes an epitope located in 1 region between extracellular domains 2 and 5 (aa 192-697) of CD222 (Roberts <i>et al.</i> , 2010).         CD222 is primarily expressed intracellularly with a small percentage of molecules being located at the cell surface (5 - 10%).         Flow Cytometry       Use 10ul of the suggested working dilution to label 1x10 <sup>6</sup> cells in 100ul.         References       1. Roberts, R.C. <i>et al.</i> (2010) Mistargeting of SH3TC2 away from the recycling endoson causes Charcot-Marie-Tooth disease type 4C. Hum Mol Genet, 19, 1009-18, 2. Adachi, A. <i>et al.</i> (2010) C0222 (Mannose-6 phosphate / insulin-like growth factor III-receptor) Surmary and Workshop Report. In Leucocyte Typing VII: White Cell Differentiation Antigens. Edited by Mason, D. <i>et al.</i> Oxford University Press. pp482-	Preparation			
Stabilisers       1% Bovine Serum Albumin         Approx. Protein Concentrations       IgG concentration 0.1 mg/ml         Immunogen       Recombinant vaccinia virus containing CD222.         External Database Links       UniProt: P11717         Related reagents       Entrez Gene: 3482         Synonyms       MPRI         RRID       AB_323453         Specificity       Mouse anti Human CD222 antibody, clone MEM-238 recognizes human CD222, a 25 KDa transmembrane protein originally identified as the IGF II receptor. CD222 is ubiquitously expressed and is involved with internalization of a variety of ligands.         Mouse anti Human CD222 antibody, clone MEM-238 recognizes an epitope located in the region between extracellular domains 2 and 5 (aa 192-697) of CD222 (Roberts <i>et al.</i> 2010).         CD2222 is primarily expressed intracellularly with a small percentage of molecules being located at the cell surface (5 - 10%).         Flow Cytometry       Use 10ul of the suggested working dilution to label 1x10 <sup>6</sup> cells in 10ul.         References       1. Roberts, R.C. <i>et al.</i> (2010) Mistargeting of SH3TC2 away from the recycling endoson causes Charcot-Marie-Tooth disease type 4C, Hum Mol Genet. 19. 1009-18.         2. Adachi, A. <i>et al.</i> (2010) Golgi-associated GSK3beta regulates the sorting process of post-Golgi membrane trafficking. J Cell Sci. 123: 3215-25.         3. Godar, S. <i>et al.</i> (2020) CD222 (Mannose-6 phosphate / insulin-like growth factor II-receptor) Summary and Workshop Report. In Leucocyte Typing VII: White Cell Differentiation Antigens. Edited by Mas	Buffer Solution	Phosphate buffered saline		
Stabilisers       1% Bovine Serum Albumin         Approx. Protein Concentrations       IgG concentration 0.1 mg/ml         Immunogen       Recombinant vaccinia virus containing CD222.         External Database Links       P11717 Related reagents         Entrez Gene: 3482 IGF2R Related reagents         Synonyms       MPRI         RRID       AB_323453         Specificity       Mouse anti Human CD222 antibody, clone MEM-238 recognizes human CD222, a 25 KDa transmembrane protein originally identified as the IGF II receptor. CD222 is ubiquitously expressed and is involved with internalization of a variety of ligands.         Mouse anti Human CD222 antibody, clone MEM-238 recognizes an epitope located in the region between extracellular domains 2 and 5 (aa 192-697) of CD222 (Roberts <i>et al.</i> 2010).         CD2222 is primarily expressed intracellularly with a small percentage of molecules being located at the cell surface (5 - 10%).         Flow Cytometry       Use 10ul of the suggested working dilution to label 1x10 <sup>6</sup> cells in 10ul.         References       1. Roberts, R.C. <i>et al.</i> (2010) Mistargeting of SH3TC2 away from the recycling endoson causes Charcot-Marie-Tooth disease type 4C. Hum Mol Genet. 19: 1009-18.         2. Adachi, A. <i>et al.</i> (2010) Golgi-associated GSK3beta regulates the sorting process of post-Golgi membrane trafficking. J Cell Sci. 123: 3215-25.         3. Godar, S. <i>et al.</i> (2002) CD222 (Mannose-6 phosphate / insulin-like growth factor II-receptor) Summary and Workshop Report. In Leucocyte Typing VII: White Cell Differentiation Antigens. Edited by Maso	Preservative	0.09% Sodium Azide		
Approx. Protein Concentrations       IgG concentration 0.1 mg/ml         Immunogen       Recombinant vaccinia virus containing CD222.         External Database Links       UniProt: P11717         Related reagents       Entrez Gene: 3482         Synonyms       MPRI         RRID       AB_323453         Specificity       Mouse anti Human CD222 antibody, clone MEM-238 recognizes human CD222, a 25 kDa transmembrane protein originally identified as the IGF II receptor. CD222 is ubiquitously expressed and is involved with internalization of a variety of ligands.         Mouse anti Human CD222 antibody, clone MEM-238 recognizes an epitope located in ti region between extracellular domains 2 and 5 (aa 192-697) of CD222 (Roberts <i>et al.</i> 2010).         CD2222 is primarily expressed intracellularly with a small percentage of molecules being located at the cell surface (5 - 10%).         Flow Cytometry       Use 10ul of the suggested working dilution to label 1x10 <sup>6</sup> cells in 100ul.         References       1. Roberts, R. C. <i>et al.</i> (2010) Mistargeting of SH3TC2 away from the recycling endoson causes Charcot-Marie-Tooth disease type 4C. Hum Mol Genet. 19: 1009-18, 2. Adachi, A. <i>et al.</i> (2010) Golgi-associated GSK3beta regulates the sorting process of post-Golgi membrane trafficking. J Cell Sci. 123: 3215-25, 3. Godar, S. <i>et al.</i> (2002) CD222 (Mannose-6 phosphate / Insulin-like growth factor II-receptor) Summary and Workshop Report. In Leucocyte Typing VII: White Cell Differentiation Antigens. Edited by Mason, D. <i>et al.</i> Oxford University Press, pp482-485, 4. Rezgui, D. <i>et al.</i> (2009) Structure and function of the human Gly1619Arg polymorphis of MPP/IGF/	Stabilisers			
Concentrations       IgG concentration 0.1 mg/ml         Immunogen       Recombinant vaccinia virus containing CD222.         External Database       UniProt:         Links       P11717         Related reagents         Synonyms       MPRI         RRID       AB_323453         Specificity       Mouse anti Human CD222 antibody, clone MEM-238 recognizes human CD222, a 25 kDa transmembrane protein originally identified as the IGF II receptor. CD222 is ubiquitously expressed and is involved with internalization of a variety of ligands.         Mouse anti Human CD222 antibody, clone MEM-238 recognizes an epitope located in the region between extracellular domains 2 and 5 (aa 192-697) of CD222 (Roberts <i>et al.</i> 2010).         CD2222 is primarily expressed intracellularly with a small percentage of molecules being located at the cell surface (5 - 10%).         Flow Cytometry       Use 10ul of the suggested working dilution to label 1x10 <sup>6</sup> cells in 100ul.         References       1. Roberts, R.C. <i>et al.</i> (2010) Mistargeting of SH3TC2 away from the recycling endoson causes Charcot-Marie-Tooth disease type 4C. Hum Mol Genet. 19: 1009-18.         2. Adachi, A. <i>et al.</i> (2001) Golgi-associated GSK3beta regulates the sorting process of post-Golgi membrane trafficking. J Cell Sci. 123: 3215-25.         3. Godar, S. <i>et al.</i> (2002) CD222 (Mannose-6 phosphate / insulin-like growth factor II-receptor) Summary and Workshop Report. In Leucocyte Typing VII: White Cell Differentiation Antigens. Edited by Mason, D. <i>et al.</i> Oxford University Press. pp482-485. <t< th=""><th></th><th></th></t<>				
External Database Links       UniProt: P11717       P11717       Related reagents         Entrez Gene: 3482       IGF2R       Related reagents         Synonyms       MPRI         RRID       AB_323453         Specificity       Mouse anti Human CD222 antibody, clone MEM-238 recognizes human CD222, a 25 KDa transmembrane protein originally identified as the IGF II receptor. CD222 is ubiquitously expressed and is involved with internalization of a variety of ligands.         Mouse anti Human CD222 antibody, clone MEM-238 recognizes an epitope located in the region between extracellular domains 2 and 5 (aa 192-697) of CD222 (Roberts <i>et al.</i> 2010).         CD222 is primarily expressed intracellularly with a small percentage of molecules being located at the cell surface (5 - 10%).         Flow Cytometry       Use 10ul of the suggested working dilution to label 1x10 <sup>6</sup> cells in 100ul.         References       1. Roberts, R.C. <i>et al.</i> (2010) Mistargeting of SH3TC2 away from the recycling endoson causes Charcot-Marie-Tooth disease type 4C. Hum Mol Genet. 19: 1009-18. 2. Adachi, A. <i>et al.</i> (2010) Goigi-associated GSMsbeta regulates the sorting process of post-Goigi membrane trafficking. J Cell Sci. 123: 3215-25. 3. Godar, S. <i>et al.</i> (2002) CD222 (Manose-6 phosphate / insulin-like growth factor II-receptor) Summary and Workshop Report. In Leucocyte Typing VII: White Cell Differentiation Antigens. Edited by Mason, D. <i>et al.</i> Oxford University Press. pp482-485. 4. Rezgui, D. <i>et al.</i> (2009) Structure and function of the human Gly1619Arg polymorphis of M6P//GF2R domain 11 implicated in IGF2 dependent growth. JMol Endocrinol. 42: 341-56.		IgG concentration 0.1 mg/ml		
Links       UniProt:         P11717       Related reagents         Entrez Gene:       3482         3482       IGF2R         RRID       AB_323453         Specificity       Mouse anti Human CD222 antibody, clone MEM-238 recognizes human CD222, a 25 kDa transmembrane protein originally identified as the IGF II receptor. CD222 is ubiquitously expressed and is involved with internalization of a variety of ligands.         Mouse anti Human CD222 antibody, clone MEM-238 recognizes an epitope located in the region between extracellular domains 2 and 5 (aa 192-697) of CD222 (Roberts <i>et al.</i> 2010).         CD222 is primarily expressed intracellularly with a small percentage of molecules being located at the cell surface (5 - 10%).         Flow Cytometry       Use 10ul of the suggested working dilution to label 1x10 <sup>6</sup> cells in 100ul.         References       1. Roberts, R.C. <i>et al.</i> (2010) Mistargeting of SH3TC2 away from the recycling endoson causes Charcot-Marie-Tooth disease type 4C. Hum Mol Genet. 19: 1009-18, 2. Adachi, A. <i>et al.</i> (2010) Golgi-associated GSK3beta regulates the sorting process of post-Golgi membrane trafficking. J Cell Sci. 123: 3215-25, 3. Godar, S. <i>et al.</i> (2002) CD222 (Mannose-6 phosphate / insulin-like growth factor II-receptor) Summary and Workshop Report. In Leucocyte Typing VII: White Cell Differentiation Antigens. Edited by Mason, D. <i>et al.</i> Oxford University Press, pp482-485, 4. Rezgui, D. <i>et al.</i> (2009) Structure and function of the human Gly1619Arg polymorphils of M6P/IGF2R domain 11 implicated in IGF2 dependent growth. J Mol Endocrinol. 42: 341-56.	Immunogen	Recombinant vaccinia virus containing CD222.		
P11717       Related reagents         Entrez Gene:       3482         3482       IGF2R         RRID       AB_323453         Specificity       Mouse anti Human CD222 antibody, clone MEM-238 recognizes human CD222, a 25 kDa transmembrane protein originally identified as the IGF II receptor. CD222 is ubiquitously expressed and is involved with internalization of a variety of ligands.         Mouse anti Human CD222 antibody, clone MEM-238 recognizes an epitope located in the region between extracellular domains 2 and 5 (aa 192-697) of CD222 (Roberts <i>et al.</i> 2010).         CD222 is primarily expressed intracellularly with a small percentage of molecules being located at the cell surface (5 - 10%).         Flow Cytometry       Use 10ul of the suggested working dilution to label 1x10 <sup>6</sup> cells in 100ul.         References       1. Roberts, R.C. <i>et al.</i> (2010) Mistargeting of SH3TC2 away from the recycling endoson causes Charoct-Marie-Tooth disease type 4C. Hum Mol Genet. 19: 1009-18.         2. Adachi, A. <i>et al.</i> (2010) Golgi-associated GSK3beta regulates the sorting process of post-Golgi membrane trafficking. J Cell Sci. 123: 3215-25.         3. Godar, S. <i>et al.</i> (2002) CD222 (Mannose-6 phosphate / insulin-like growth factor III-receptor) Summary and Workshop Report. In Leucocyte Typing VII: White Cell Differentiation Antigens. Edited by Mason, D. <i>et al.</i> Oxford University Press. p482-485.         4. Rezgui, D. <i>et al.</i> (2009) Structure and function of the human Gly1619Arg polymorphis of M6P/IGF2R domain 11 implicated in IGF2 dependent growth. J Mol Endocrinol. 42: 341-56.	External Database			
Entrez Gene:       3482       IGF2R       Related reagents         Synonyms       MPRI         RRID       AB_323453         Specificity       Mouse anti Human CD222 antibody, clone MEM-238 recognizes human CD222, a 25 kDa transmembrane protein originally identified as the IGF II receptor. CD222 is ubiquitously expressed and is involved with internalization of a variety of ligands.         Mouse anti Human CD222 antibody, clone MEM-238 recognizes an epitope located in the region between extracellular domains 2 and 5 (aa 192-697) of CD222 (Roberts <i>et al.</i> 2010).         CD222 is primarily expressed intracellularly with a small percentage of molecules being located at the cell surface (5 - 10%).         Flow Cytometry       Use 10ul of the suggested working dilution to label 1x10 <sup>6</sup> cells in 100ul.         References       1. Roberts, R.C. <i>et al.</i> (2010) Mistargeting of SH3TC2 away from the recycling endoson causes Charcot-Marie-Tooth disease type 4C. Hum Mol Genet. 19: 1009-18.         2. Adachi, A. <i>et al.</i> (2010) Golgi-associated GSK3beta regulates the sorting process of post-Golgi membrane trafficking. J Cell Sci. 123: 3215-25.         3. Godar, S. <i>et al.</i> (2002) CD222 (Mannose-6 phosphate / insulin-like growth factor II-receptor) Summary and Workshop Report. In Leucocyte Typing VI: White Cell Differentiation Antigens. Edited by Mason, D. <i>et al.</i> 0xford University Press. pp482-485.         4. Rezgui, D. <i>et al.</i> (2009) Structure and function of the human Gly1619Arg polymorphis of M6P/IGF2R domain 11 implicated in IGF2 dependent growth. J Mol Endocrind. 42: 341-56.	Links	UniProt:		
3482       IGF2R       Related reagents         Synonyms       MPRI         RRID       AB_323453         Specificity       Mouse anti Human CD222 antibody, clone MEM-238 recognizes human CD222, a 25 kDa transmembrane protein originally identified as the IGF II receptor. CD222 is ubiquitously expressed and is involved with internalization of a variety of ligands.         Mouse anti Human CD222 antibody, clone MEM-238 recognizes an epitope located in the region between extracellular domains 2 and 5 (aa 192-697) of CD222 (Roberts <i>et al.</i> 2010).         CD222 is primarily expressed intracellularly with a small percentage of molecules being located at the cell surface (5 - 10%).         Flow Cytometry       Use 10ul of the suggested working dilution to label 1x10 <sup>6</sup> cells in 100ul.         References       1. Roberts, R.C. <i>et al.</i> (2010) Mistargeting of SH3TC2 away from the recycling endoson causes Charcot-Marie-Tooth disease type 4C. Hum Mol Genet. 19: 1009-18.         2. Adachi, A. <i>et al.</i> (2010) Golgi-associated GSK3beta regulates the sorting process of post-Golgi membrane trafficking. J Cell Sci. 123: 3215-25.         3. Godar, S. <i>et al.</i> (2000) CD222 (Mannose-6 phosphate / insulin-like growth factor II-receptor) Summary and Workshop Report. In Leucocyte Typing VII: White Cell Differentiation Antigens. Edited by Mason, D. <i>et al.</i> Oxford University Press. pp482-485.         4. Rezgui, D. <i>et al.</i> (2009) Structure and function of the human Gly1619Arg polymorphis of M6P/IGF2R domain 11 implicated in IGF2 dependent growth. J Mol Endocrinol. 42: 341-56.		P11717 Related reagents		
3482       IGF2R       Related reagents         Synonyms       MPRI         RRID       AB_323453         Specificity       Mouse anti Human CD222 antibody, clone MEM-238 recognizes human CD222, a 25 kDa transmembrane protein originally identified as the IGF II receptor. CD222 is ubiquitously expressed and is involved with internalization of a variety of ligands.         Mouse anti Human CD222 antibody, clone MEM-238 recognizes an epitope located in the region between extracellular domains 2 and 5 (aa 192-697) of CD222 (Roberts <i>et al.</i> 2010).         CD222 is primarily expressed intracellularly with a small percentage of molecules being located at the cell surface (5 - 10%).         Flow Cytometry       Use 10ul of the suggested working dilution to label 1x10 <sup>6</sup> cells in 100ul.         References       1. Roberts, R.C. <i>et al.</i> (2010) Mistargeting of SH3TC2 away from the recycling endoson causes Charcot-Marie-Tooth disease type 4C. Hum Mol Genet. 19: 1009-18.         2. Adachi, A. <i>et al.</i> (2010) Golgi-associated GSK3beta regulates the sorting process of post-Golgi membrane trafficking. J Cell Sci. 123: 3215-25.         3. Godar, S. <i>et al.</i> (2000) CD222 (Mannose-6 phosphate / insulin-like growth factor II-receptor) Summary and Workshop Report. In Leucocyte Typing VII: White Cell Differentiation Antigens. Edited by Mason, D. <i>et al.</i> Oxford University Press. pp482-485.         4. Rezgui, D. <i>et al.</i> (2009) Structure and function of the human Gly1619Arg polymorphis of M6P/IGF2R domain 11 implicated in IGF2 dependent growth. J Mol Endocrinol. 42: 341-56.		Entroz Gono:		
Synonyms       MPRI         RRID       AB_323453         Specificity       Mouse anti Human CD222 antibody, clone MEM-238 recognizes human CD222, a 25 kDa transmembrane protein originally identified as the IGF II receptor. CD222 is ubiquitously expressed and is involved with internalization of a variety of ligands.         Mouse anti Human CD222 antibody, clone MEM-238 recognizes an epitope located in the region between extracellular domains 2 and 5 (aa 192-697) of CD222 (Roberts <i>et al.</i> 2010).         CD222 is primarily expressed intracellularly with a small percentage of molecules being located at the cell surface (5 - 10%).         Flow Cytometry       Use 10ul of the suggested working dilution to label 1x10 <sup>6</sup> cells in 100ul.         References       1. Roberts, R.C. <i>et al.</i> (2010) Mistargeting of SH3TC2 away from the recycling endoson causes Charcot-Marie-Tooth disease type 4C. Hum Mol Genet. 19: 1009-18.         2. Adachi, A. <i>et al.</i> (2010) Golgi-associated GSK3beta regulates the sorting process of post-Golgi membrane trafficking. J Cell Sci. 123: 3215-25.         3. Godar, S. <i>et al.</i> (2002) CD222 (Mannose-6 phosphate / insulin-like growth factor II-receptor) Summary and Workshop Report. In Leucocyte Typing VII: White Cell Differentiation Antigens. Edited by Mason, D. <i>et al.</i> Oxford University Press. pp482-485.         4. Rezgui, D. <i>et al.</i> (2009) Structure and function of the human Gly1619Arg polymorphis of M6P/IGF2R domain 11 implicated in IGF2 dependent growth. J Mol Endocrinol. 42: 341-56.				
RRID       AB_323453         Specificity       Mouse anti Human CD222 antibody, clone MEM-238 recognizes human CD222, a 25 kDa transmembrane protein originally identified as the IGF II receptor. CD222 is ubiquitously expressed and is involved with internalization of a variety of ligands.         Mouse anti Human CD222 antibody, clone MEM-238 recognizes an epitope located in the region between extracellular domains 2 and 5 (aa 192-697) of CD222 (Roberts <i>et al.</i> 2010).         CD222 is primarily expressed intracellularly with a small percentage of molecules being located at the cell surface (5 - 10%).         Flow Cytometry       Use 10ul of the suggested working dilution to label 1x10 <sup>6</sup> cells in 100ul.         References       1. Roberts, R.C. <i>et al.</i> (2010) Mistargeting of SH3TC2 away from the recycling endoson causes Charcot-Marie-Tooth disease type 4C. Hum Mol Genet. 19: 1009-18.         2. Adachi, A. <i>et al.</i> (2010) Golgi-associated GSK3beta regulates the sorting process of post-Golgi membrane trafficking. J Cell Sci. 123: 3215-25.         3. Godar, S. <i>et al.</i> (2002) CD222 (Mannose-6 phosphate / insulin-like growth factor II-receptor) Summary and Workshop Report. In Leucocyte Typing VII: White Cell Differentiation Antigens. Edited by Mason, D. <i>et al.</i> 0xford University Press. pp482-485.         4. Rezgui, D. <i>et al.</i> (2009) Structure and function of the human Glyf619Arg polymorphis of M6P/IGF2R domain 11 implicated in IGF2 dependent growth. J Mol Endocrinol. 42: 341-56.		<u>5462</u> IOI 2IX <u>related reagents</u>		
Specificity       Mouse anti Human CD222 antibody, clone MEM-238 recognizes human CD222, a 25 kDa transmembrane protein originally identified as the IGF II receptor. CD222 is ubiquitously expressed and is involved with internalization of a variety of ligands.         Mouse anti Human CD222 antibody, clone MEM-238 recognizes an epitope located in the region between extracellular domains 2 and 5 (aa 192-697) of CD222 (Roberts <i>et al.</i> 2010).         CD222 is primarily expressed intracellularly with a small percentage of molecules being located at the cell surface (5 - 10%).         Flow Cytometry       Use 10ul of the suggested working dilution to label 1x10 <sup>6</sup> cells in 100ul.         References       1. Roberts, R.C. <i>et al.</i> (2010) Mistargeting of SH3TC2 away from the recycling endoson causes Charcot-Marie-Tooth disease type 4C. Hum Mol Genet. 19: 1009-18.         2. Adachi, A. <i>et al.</i> (2010) Golgi-associated GSK3beta regulates the sorting process of post-Golgi membrane trafficking. J Cell Sci. 123: 3215-25.         3. Godar, S. <i>et al.</i> (2002) CD222 (Mannose-6 phosphate / insulin-like growth factor II-receptor) Summary and Workshop Report. In Leucocyte Typing VII: White Cell Differentiation Antigens. Edited by Mason, D. <i>et al.</i> Oxford University Press. pp482-485.         4. Rezgui, D. <i>et al.</i> (2009) Structure and function of the human Gly1619Arg polymorphis of M6P/IGF2R domain 11 implicated in IGF2 dependent growth. J Mol Endocrinol. 42: 341-56.	Synonyms	MPRI		
<ul> <li>KDa transmembrane protein originally identified as the IGF II receptor. CD222 is ubiquitously expressed and is involved with internalization of a variety of ligands.</li> <li>Mouse anti Human CD222 antibody, clone MEM-238 recognizes an epitope located in the region between extracellular domains 2 and 5 (aa 192-697) of CD222 (Roberts <i>et al.</i> 2010).</li> <li>CD222 is primarily expressed intracellularly with a small percentage of molecules being located at the cell surface (5 - 10%).</li> <li>Flow Cytometry Use 10ul of the suggested working dilution to label 1x10<sup>6</sup> cells in 100ul.</li> <li>References 1. Roberts, R.C. <i>et al.</i> (2010) Mistargeting of SH3TC2 away from the recycling endoson causes Charcot-Marie-Tooth disease type 4C. Hum Mol Genet. 19: 1009-18.</li> <li>2. Adachi, A. <i>et al.</i> (2010) Golgi-associated GSK3beta regulates the sorting process of post-Golgi membrane trafficking. J Cell Sci. 123: 3215-25.</li> <li>3. Godar, S. <i>et al.</i> (2002) CD222 (Mannose-6 phosphate / insulin-like growth factor II-receptor) Summary and Workshop Report. In Leucocyte Typing VII: White Cell Differentiation Antigens. Edited by Mason, D. <i>et al.</i> Oxford University Press. pp482-485.</li> <li>4. Rezgui, D. <i>et al.</i> (2009) Structure and function of the human Gly1619Arg polymorphis of M6P/IGF2R domain 11 implicated in IGF2 dependent growth. J Mol Endocrinol. 42: 341-56.</li> </ul>	RRID	AB_323453		
<ul> <li>region between extracellular domains 2 and 5 (aa 192-697) of CD222 (Roberts <i>et al.</i> 2010).</li> <li>CD222 is primarily expressed intracellularly with a small percentage of molecules being located at the cell surface (5 - 10%).</li> <li>Flow Cytometry Use 10ul of the suggested working dilution to label 1x10<sup>6</sup> cells in 100ul.</li> <li>References 1. Roberts, R.C. <i>et al.</i> (2010) Mistargeting of SH3TC2 away from the recycling endoson causes Charcot-Marie-Tooth disease type 4C. Hum Mol Genet. 19: 1009-18.</li> <li>2. Adachi, A. <i>et al.</i> (2010) Golgi-associated GSK3beta regulates the sorting process of post-Golgi membrane trafficking. J Cell Sci. 123: 3215-25.</li> <li>3. Godar, S. <i>et al.</i> (2002) CD222 (Mannose-6 phosphate / insulin-like growth factor II-receptor) Summary and Workshop Report. In Leucocyte Typing VII: White Cell Differentiation Antigens. Edited by Mason, D. <i>et al.</i> Oxford University Press. pp482-485.</li> <li>4. Rezgui, D. <i>et al.</i> (2009) Structure and function of the human Gly1619Arg polymorphis of M6P/IGF2R domain 11 implicated in IGF2 dependent growth. J Mol Endocrinol. 42: 341-56.</li> </ul>	Specificity			
Iocated at the cell surface (5 - 10%).         Flow Cytometry       Use 10ul of the suggested working dilution to label 1x10 <sup>6</sup> cells in 100ul.         References       1. Roberts, R.C. et al. (2010) Mistargeting of SH3TC2 away from the recycling endoson causes Charcot-Marie-Tooth disease type 4C. Hum Mol Genet. 19: 1009-18.         2. Adachi, A. et al. (2010) Golgi-associated GSK3beta regulates the sorting process of post-Golgi membrane trafficking. J Cell Sci. 123: 3215-25.         3. Godar, S. et al. (2002) CD222 (Mannose-6 phosphate / insulin-like growth factor II-receptor) Summary and Workshop Report. In Leucocyte Typing VII: White Cell Differentiation Antigens. Edited by Mason, D. et al. Oxford University Press. pp482-485.         4. Rezgui, D. et al. (2009) Structure and function of the human Gly1619Arg polymorphis of M6P/IGF2R domain 11 implicated in IGF2 dependent growth. J Mol Endocrinol. 42: 341-56.				
References       1. Roberts, R.C. et al. (2010) Mistargeting of SH3TC2 away from the recycling endoson causes Charcot-Marie-Tooth disease type 4C. Hum Mol Genet. 19: 1009-18.         2. Adachi, A. et al. (2010) Golgi-associated GSK3beta regulates the sorting process of post-Golgi membrane trafficking. J Cell Sci. 123: 3215-25.         3. Godar, S. et al. (2002) CD222 (Mannose-6 phosphate / insulin-like growth factor II-receptor) Summary and Workshop Report. In Leucocyte Typing VII: White Cell Differentiation Antigens. Edited by Mason, D. et al. Oxford University Press. pp482-485.         4. Rezgui, D. et al. (2009) Structure and function of the human Gly1619Arg polymorphis of M6P/IGF2R domain 11 implicated in IGF2 dependent growth. J Mol Endocrinol. 42: 341-56.		CD222 is primarily expressed intracellularly with a small percentage of molecules being located at the cell surface (5 - 10%).		
<ul> <li>causes Charcot-Marie-Tooth disease type 4C. <u>Hum Mol Genet. 19: 1009-18.</u></li> <li>2. Adachi, A. <i>et al.</i> (2010) Golgi-associated GSK3beta regulates the sorting process of post-Golgi membrane trafficking. <u>J Cell Sci. 123: 3215-25.</u></li> <li>3. Godar, S. <i>et al.</i> (2002) CD222 (Mannose-6 phosphate / insulin-like growth factor II-receptor) Summary and Workshop Report. In Leucocyte Typing VII: White Cell Differentiation Antigens. Edited by Mason, D. <i>et al.</i> Oxford University Press. pp482-485.</li> <li>4. Rezgui, D. <i>et al.</i> (2009) Structure and function of the human Gly1619Arg polymorphis of M6P/IGF2R domain 11 implicated in IGF2 dependent growth. <u>J Mol Endocrinol. 42: 341-56.</u></li> </ul>	Flow Cytometry	Use 10ul of the suggested working dilution to label 1x10 <sup>6</sup> cells in 100ul.		
5. IVICUOITIICK, P.J. et al. (2008) Paimitoyiation controls recycling in lysosomal softing ar	References	<ol> <li>Adachi, A. <i>et al.</i> (2010) Golgi-associated GSK3beta regulates the sorting process of post-Golgi membrane trafficking. <u>J Cell Sci. 123: 3215-25.</u></li> <li>Godar, S. <i>et al.</i> (2002) CD222 (Mannose-6 phosphate / insulin-like growth factor II-receptor) Summary and Workshop Report. In Leucocyte Typing VII: White Cell Differentiation Antigens. Edited by Mason, D. <i>et al.</i> Oxford University Press. pp482-485.</li> <li>Rezgui, D. <i>et al.</i> (2009) Structure and function of the human Gly1619Arg polymorphism of M6P/IGF2R domain 11 implicated in IGF2 dependent growth. <u>J Mol Endocrinol. 42:</u></li> </ol>		

	<ul> <li>trafficking. <u>Traffic. 9: 1984-97.</u></li> <li>6. Leksa, V. <i>et al.</i> (2002) The N terminus of mannose 6-phosphate/insulinfactor 2 receptor in regulation of fibrinolysis and cell migration. <u>J Biol Cher</u> <u>40575-82.</u></li> <li>7. Osborne, D.G. <i>et al.</i> (2015) Monitoring receptor trafficking following retr WASH deregulation. <u>Methods Cell Biol. 130: 199-213.</u></li> </ul>	<u>m. 277 (43):</u>
Storage	Store at +4°C or at -20°C if preferred.	
	This product should be stored undiluted.	
	Storage in frost free freezers is not recommended. This product is photose should be protected from light.	ensitive and
	Avoid repeated freezing and thawing as this may denature the antibody. S product contain a precipitate we recommend microcentrifugation before us	
Guarantee	12 months from date of despatch	
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/MCA2048F 10041	
Regulatory	For research purposes only	

## **Related Products**

### **Recommended Negative Controls**

MOUSE IgG1 NEGATIVE CONTROL: FITC (MCA928F)

### **Recommended Useful Reagents**

HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-rac	d.com	Email: antibody_sales_uk@bio-rac	d.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M366105:200529'

#### Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint