

Datasheet: MCA2059

| | |
|----------------------|-----------------------|
| Description: | MOUSE ANTI HUMAN CD88 |
| Specificity: | CD88 |
| Other names: | C5aR |
| Format: | Purified |
| Product Type: | Monoclonal Antibody |
| Clone: | P12/1 |
| Isotype: | IgG2a |
| Quantity: | 0.2 mg |

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

| | Yes | No | Not Determined | Suggested Dilution |
|----------------------------|-----|----|----------------|--------------------|
| Flow Cytometry | ▪ | | | 1/50 - 1/100 |
| Immunohistology - Frozen | ▪ | | | |
| Immunohistology - Paraffin | ▪ | | | |
| ELISA | | | ▪ | |
| Immunoprecipitation | | ▪ | | |
| Western Blotting | ▪ | | | |

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species

Human

Species Cross Reactivity

Reacts with: Rhesus Monkey

N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

Product Form

Purified IgG - liquid

Preparation

Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant

| | |
|---------------------------------------|---|
| Buffer Solution | Phosphate buffered saline |
| Preservative Stabilisers | 0.09% Sodium Azide |
| Carrier Free | Yes |
| Approx. Protein Concentrations | IgG concentration 1 mg/ml |
| Immunogen | C5aR - peptide: Met ₁ - Asn ₃₁ |
| External Database Links | <p>UniProt: P21730 Related reagents</p> <p>Entrez Gene: 728 C5AR1 Related reagents</p> |
| Synonyms | C5AR, C5R1 |
| RRID | AB_566906 |
| Fusion Partners | Spleen cells from immunised BALB/c mice were fused with cells of the mouse X63 - Ag8 myeloma cell line. |
| Specificity | <p>Mouse anti Human CD88 antibody, clone P12/1 recognizes the C5a receptor (C5aR) also known as CD88 or C5a anaphylatoxin chemotactic receptor 1. CD88 is predominantly expressed on cells of the myeloid lineage.</p> <p>When C5aR is preincubated with C5a, Mouse anti Human CD88 antibody, clone P12/1 does not bind to the receptor, as the binding site of P12/1 is located in the C5a binding region (Werfel et al. 1996 and Weinman et al. 2003)</p> |
| Flow Cytometry | Use 10ul of the suggested working dilution to label 5 x 10 ⁵ cells in 100ul. |
| References | <ol style="list-style-type: none"> 1. Werfel, T. <i>et al.</i> (1996) CD88 antibodies specifically bind to C5aR on dermal CD117+ and CD14+ cells and react with a desmosomal antigen in human skin. J Immunol. 157: 1729-35. 2. Opperman, M. <i>et al.</i> (1995) Antibodies from the myeloid panel that react with the C5a receptor and antagonize C5a biological activity. In: Schlossman, S.F. (ed.) Leucocyte Typing V. OUP: 955-6. 3. Pollok-Kopp B <i>et al.</i> (2007) Dynamics of protein kinase C-mediated phosphorylation of the complement C5a receptor on serine 334. J Biol Chem. 282 (7): 4345-53. 4. Werfel, T. <i>et al.</i> (1995) Binding of anti-C5a receptor (C5aR) antibodies to cells of clinically normal human skin. In: Schlossman, S.F. (ed.) Leucocyte Typing V. OUP: 957-9. 5. Oppermann, M. (1995) Cluster report: (C5a receptor). In: Schlossman, S.F. (ed.) Leucocyte Typing V. OUP: 953-4. 6. Hüttenrauch, F. <i>et al.</i> (2005) G protein-coupled receptor kinases promote |

phosphorylation and beta-arrestin-mediated internalization of CCR5 homo- and hetero-oligomers. [J Biol Chem. 280: 37503-15.](#)

7. Huber-Lang, M. *et al.* (2005) Changes in the novel orphan, C5a receptor (C5L2), during experimental sepsis and sepsis in humans. [J Immunol. 174 \(2\): 1104-10.](#)

8. Morris, A.C. *et al.* (2011) C5a-mediated neutrophil dysfunction is RhoA-dependent and predicts infection in critically ill patients. [Blood. 117: 5178-88.](#)

9. Visser T *et al.* (2012) Homology in systemic neutrophil response induced by human experimental endotoxemia and by trauma. [Shock. 37 \(2\): 145-51.](#)

10. Nitta, H. *et al.* (2013) Enhancement of human cancer cell motility and invasiveness by anaphylatoxin C5a via aberrantly expressed C5a receptor (CD88). [Clin Cancer Res. 19 \(8\): 2004-13.](#)

11. Patzelt, J. *et al.* (2015) Expression of anaphylatoxin receptors on platelets in patients with coronary heart disease. [Atherosclerosis. 238 \(2\): 289-95.](#)

12. Unnewehr H *et al.* (2013) Changes and regulation of the C5a receptor on neutrophils during septic shock in humans. [J Immunol. 190 \(8\): 4215-25.](#)

13. Visser T *et al.* (2011) Isolated blunt chest injury leads to transient activation of circulating neutrophils. [Eur J Trauma Emerg Surg. 37 \(2\): 177-84.](#)

14. Presicce, P. *et al.* (2015) Neutrophil recruitment and activation in decidua with intra-amniotic IL-1beta in the preterm rhesus macaque. [Biol Reprod. 92 \(2\): 56.](#)

15. Zannetti, C. *et al.* (2016) Characterization of the Inflammasome in Human Kupffer Cells in Response to Synthetic Agonists and Pathogens. [J Immunol. 197 \(1\): 356-67.](#)

16. Werfel, T. *et al.* (1996) The human mast cell line HMC-1 expresses C5a receptors and responds to C5a but not to C5a(desArg). [Scand J Immunol. 44 \(1\): 30-6.](#)

17. Weinmann, O. *et al.* (2003) Up-regulation of C5a receptor expression and function on human monocyte derived dendritic cells by prostaglandin E2. [Immunology. 110 \(4\): 458-65.](#)

18. Hodille, E. *et al.* (2020) Staphylococcal Panton–Valentine Leucocidin and Gamma Haemolysin Target and Lyse Mature Bone Marrow Leucocytes. [Toxins. 12 \(11\): 725.](#)

19. Henes, J.K. *et al.* (2021) C5 Variant rs10985126 is Associated with Mortality in Patients with Symptomatic Coronary Artery Disease. [Pharmgenomics Pers Med. 14: 893-903.](#)

Further Reading

1. Oppermann, M. *et al.* (1993) Probing the human receptor for C5a anaphylatoxin with site-directed antibodies. Identification of a potential ligand binding site on the NH₂-terminal domain. [J Immunol. 151 \(7\): 3785-94.](#)

2. Oppermann, M. & Götze, O. (1994) Plasma clearance of the human C5a anaphylatoxin by binding to leucocyte C5a receptors. [Immunology. 82 \(4\): 516-21.](#)

Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee

12 months from date of despatch

Health And Safety

Material Safety Datasheet documentation #10040 available at:

Information 10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

| | |
|---|---|
| Goat Anti Mouse IgG (STAR77...) | HRP |
| Rabbit Anti Mouse IgG (STAR12...) | RPE |
| Rabbit Anti Mouse IgG (STAR8...) | DyLight@800 |
| Goat Anti Mouse IgG (STAR76...) | RPE |
| Human Anti Mouse IgG2a (HCA037...) | HRP |
| Goat Anti Mouse IgG (Fc) (STAR120...) | FITC , HRP |
| Rabbit Anti Mouse IgG (STAR13...) | HRP |
| Goat Anti Mouse IgG (H/L) (STAR117...) | Alk. Phos. , DyLight@488 , DyLight@550 , DyLight@650 , DyLight@680 , DyLight@800 , FITC , HRP |
| Goat Anti Mouse IgG (STAR70...) | FITC |
| Rabbit Anti Mouse IgG (STAR9...) | FITC |
| Goat Anti Mouse IgG IgA IgM (STAR87...) | Alk. Phos. , HRP |

Recommended Negative Controls

[MOUSE IgG2a NEGATIVE CONTROL \(MCA929\)](#)

| | | | | | |
|----------------------------------|---|------------------|---|---------------|---|
| North & South America | Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com | Worldwide | Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com | Europe | Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com |
|----------------------------------|---|------------------|---|---------------|---|

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M382969:210513'

Printed on 19 Oct 2021