

Datasheet: MCA2061GA

BATCH NUMBER 156419

Description:	MOUSE ANTI HUMAN CD284
Specificity:	CD284
Other names:	TLR4
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	HTA125
Isotype:	IgG2a
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/10 - 1/50
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting	▪			
Immunofluorescence			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human
Species Cross Reactivity	<p>Reacts with: Rhesus Monkey, Guinea Pig, Pig, Dog, Bovine</p> <p>N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.</p>
Product Form	Purified IgG - liquid

Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Ba/F3 cell line expressing TLR4 (CD284).
External Database Links	<p>UniProt: O00206 Related reagents</p> <p>Entrez Gene: 7099 TLR4 Related reagents</p>
RRID	AB_323818
Fusion Partners	Spleen cells from immunised Balb/c mice were fused with cells of the mouse SP2/0 myeloma cell line.
Specificity	<p>Mouse anti Human CD284 antibody, clone HTA125 recognizes the human Toll like receptor 4 (TLR4) cell surface antigen.</p> <p>TLR4, also known as CD284, has been demonstrated to act as a receptor for LPS on human monocytes and macrophages. TLR4 signalling of LPS stimulation requires the presence of the MD-2 molecule.</p> <p>TLR4 is weakly expressed by resting cells, but is upregulated following stimulation with LPS.</p> <p>This antibody has been demonstrated to block activation of monocytes with LPS. The use of a preservative free format of Mouse anti Human CD284 antibody, clone HTA125 (MCA2061EL) is recommended for functional assays.</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells or 100ul whole blood.
References	<ol style="list-style-type: none"> 1. Shimazu, R. <i>et al.</i> (1999) MD-2, a molecule that confers lipopolysaccharide responsiveness on Toll-like receptor 4. J Exp Med. 189 (11): 1777-82. 2. Jiang, Q. <i>et al.</i> (2000) Lipopolysaccharide induces physical proximity between CD14 and toll-like receptor 4 (TLR4) prior to nuclear translocation of NF-kappa B. J Immunol. 165 (7): 3541-4. 3. Yang, S. <i>et al.</i> (2001) Synergistic effect of muramyldipeptide with lipopolysaccharide or

- lipoteichoic acid to induce inflammatory cytokines in human monocytic cells in culture. [Infect Immun. 69 \(4\): 2045-53.](#)
4. Kawahara T *et al.* (2001) Type I *Helicobacter pylori* lipopolysaccharide stimulates toll-like receptor 4 and activates mitogen oxidase 1 in gastric pit cells. [Infect Immun. 69 \(7\): 4382-9.](#)
 5. Devaney, J.M. (2003) Neutrophil elastase up-regulates interleukin-8 via toll-like receptor 4. [FEBS Lett. 544:129-32.](#)
 6. de Kleer, I. (2010) CD30 Discriminates Heat Shock Protein 60-Induced FOXP3+CD4+ T Cells with a Regulatory Phenotype. [J Immunol. 185\(4\):2071-9.](#)
 7. Bieback, K. *et al.* (2002) Hemagglutinin protein of wild-type measles virus activates toll-like receptor 2 signaling. [J Virol. 76: 8729-36.](#)
 8. Brännström, K. *et al.* (2009) The *Schistosoma mansoni* protein Sm16/SmSLP /SmSPO-1 assembles into a nine-subunit oligomer with potential To inhibit Toll-like receptor signaling. [Infect Immun. 77: 1144-54.](#)
 9. Baumgarten, G. *et al.* (2001) *In vivo* expression of proinflammatory mediators in the adult heart after endotoxin administration: the role of toll-like receptor-4. [J Infect Dis. 183: 1617-24.](#)
 10. Cuschieri, J. *et al.* (2006) Endotoxin tolerance attenuates LPS-induced TLR4 mobilization to lipid rafts: a condition reversed by PKC activation. [J Leukoc Biol. 80: 1289-97.](#)
 11. Karlsson, H. *et al.* (2002) Innate immune responses of human neonatal cells to bacteria from the normal gastrointestinal flora. [Infect Immun. 70: 6688-96.](#)
 12. Medvedev, A.E. *et al.* (2001) Induction of tolerance to lipopolysaccharide and mycobacterial components in Chinese hamster ovary/CD14 cells is not affected by overexpression of Toll-like receptors 2 or 4. [J Immunol. 167: 2257-67.](#)
 13. Pioli, P.A. *et al.* (2007) Estradiol attenuates lipopolysaccharide-induced CXC chemokine ligand 8 production by human peripheral blood monocytes. [J Immunol. 179: 6284-90.](#)
 14. Sugawara, S. *et al.* (2000) Proteolysis of human monocyte CD14 by cysteine proteinases (gingipains) from *Porphyromonas gingivalis* leading to lipopolysaccharide hyporesponsiveness. [J Immunol. 165: 411-8.](#)
 15. Lindsay, J.O. *et al.* (2006) Clinical, microbiological, and immunological effects of fructo-oligosaccharide in patients with Crohn's disease. [Gut. 55: 348-55.](#)
 16. Komori, H. *et al.* (2012) $\alpha(1)$ -Acid glycoprotein up-regulates CD163 via TLR4/CD14 protein pathway: possible protection against hemolysis-induced oxidative stress. [J Biol Chem. 287 \(36\): 30688-700.](#)
 17. Maiolini, A. *et al.* (2012) Toll-like receptors 4 and 9 are responsible for the maintenance of the inflammatory reaction in canine steroid-responsive meningitis-arteritis, a large animal model for neutrophilic meningitis. [J Neuroinflammation. 9: 226.](#)
 18. Sels, J.W. *et al.* (2012) Fractional flow reserve is not associated with inflammatory markers in patients with stable coronary artery disease. [PLoS One. 7: e46356.](#)
 19. Prokhorenko, I. *et al.* (2012) Toll-like receptor 4 in phagocytosis of Escherichia coli by endotoxin-activated human neutrophils in whole blood [Critical Care 16: P80](#)
 20. Mazzucchelli, I. *et al.* (2015) Expression and function of toll-like receptors in human circulating endothelial colony forming cells. [Immunol Lett. 168 \(1\): 98-104.](#)
 21. Garbe, K. *et al.* (2012) Plasmacytoid dendritic cells and their Toll-like receptor 9 expression selectively decrease with age. [Hum Immunol. 73 \(5\): 493-7.](#)

22. Zwolak, A. *et al.* (2016) Metformin Changes the Relationship between Blood Monocyte Toll-Like Receptor 4 Levels and Nonalcoholic Fatty Liver Disease-*Ex Vivo* Studies. [PLoS One. 11 \(3\): e0150233.](#)
23. Zwolak, A. *et al.* (2015) Hyperreactivity of Blood Leukocytes in Patients with NAFLD to *ex vivo* Lipopolysaccharide Treatment Is Modulated by Metformin and Phosphatidylcholine but Not by Alpha Ketoglutarate. [PLoS One. 10 \(12\): e0143851.](#)
24. Xu, H. *et al.* (2015) Type 3 innate lymphoid cell depletion is mediated by TLRs in lymphoid tissues of simian immunodeficiency virus-infected macaques. [FASEB J. 29 \(12\): 5072-80.](#)
25. Blagitz, M.G. *et al.* (2015) Expression of CD14 and toll-like receptors 2 and 4 by milk neutrophils in bovine mammary glands infected with *Corynebacterium bovis* [Pesquisa Veterinária Brasileira. 35 \(1\): 1-5.](#)
26. Huang, D. *et al.* (2016) Hyperoxia induces inflammation and regulates cytokine production in alveolar epithelium through TLR2/4-NF-κB-dependent mechanism [Eur Rev Med Pharmacol Sci. 20: 1399-410.](#)
27. Kyrova, K. *et al.* (2014) The response of porcine monocyte derived macrophages and dendritic cells to *Salmonella typhimurium* and lipopolysaccharide. [BMC Vet Res. 10: 244.](#)
28. Ibeagha-Awemu, E.M. *et al.* (2008) Bacterial lipopolysaccharide induces increased expression of toll-like receptor (TLR) 4 and downstream TLR signaling molecules in bovine mammary epithelial cells. [Vet Res. 39 \(2\): 11.](#)
29. Chochi, K. *et al.* (2008) *Helicobacter pylori* augments growth of gastric cancers via the lipopolysaccharide-toll-like receptor 4 pathway whereas its lipopolysaccharide attenuates antitumor activities of human mononuclear cells. [Clin Cancer Res. 14 \(10\): 2909-17.](#)
30. Elner, S.G. *et al.* (2005) TLR4 mediates human retinal pigment epithelial endotoxin binding and cytokine expression. [Invest Ophthalmol Vis Sci. 46 \(12\): 4627-33.](#)

Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the protein. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee

12 months from date of despatch

Health And Safety Information

Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/MCA2061GA>
10040

Regulatory

For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...) [RPE](#)

Goat Anti Mouse IgG IgA IgM (STAR87...) [HRP](#)

Goat Anti Mouse IgG (STAR76...)	RPE
Goat Anti Mouse IgG (STAR70...)	FITC
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight®488 , DyLight®550 , DyLight®650 , DyLight®680 , DyLight®800 , FITC , HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC
Goat Anti Mouse IgG (STAR77...)	HRP
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP
Rabbit Anti Mouse IgG (STAR13...)	HRP

Recommended Negative Controls

[MOUSE IgG2a NEGATIVE CONTROL \(MCA929\)](#)

North & South America	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-rad.com		Email: antibody_sales_uk@bio-rad.com		Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M366131:200529'

Printed on 20 Mar 2024