

Datasheet: MCA2061SBV440

BATCH NUMBER 64574795

Description:	MOUSE ANTI HUMAN CD284:StarBright Violet 440
Specificity:	CD284
Other names:	TLR4
Format:	StarBright Violet 440
Product Type:	Monoclonal Antibody
Clone:	HTA125
Isotype:	IgG2a
Quantity:	100 TESTS/0.5ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species Human

Species Cross Reactivity Reacts with: Rhesus Monkey, Guinea Pig, Pig, Dog, Bovine
N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

Product Form Purified IgG conjugated to StarBright Violet 440 - liquid

Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	StarBright Violet 440	383	436

Preparation Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant

Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃) 1% Bovine Serum Albumin 0.1% Pluronic F68 0.1% PEG 3350 0.05% Tween 20
Immunogen	Ba/F3 cell line expressing TLR4 (CD284).
External Database Links	<p>UniProt: O00206 Related reagents</p> <p>Entrez Gene: 7099 TLR4 Related reagents</p>
Fusion Partners	Spleen cells from immunised Balb/c mice were fused with cells of the mouse SP2/0 myeloma cell line.
Specificity	<p>Mouse anti Human CD284 antibody, clone HTA125 recognizes the human Toll like receptor 4 (TLR4) cell surface antigen.</p> <p>TLR4, also known as CD284, has been demonstrated to act as a receptor for LPS on human monocytes and macrophages. TLR4 signalling of LPS stimulation requires the presence of the MD-2 molecule.</p> <p>TLR4 is weakly expressed by resting cells, but is upregulated following stimulation with LPS.</p> <p>This antibody has been demonstrated to block activation of monocytes with LPS. The use of a preservative free format of Mouse anti Human CD284 antibody, clone HTA125 (MCA2061EL) is recommended for functional assays.</p>
Flow Cytometry	Use 5µl of the suggested working dilution to label 10 ⁶ cells in 100µl. Best practices suggest a 5 minutes centrifugation at 6,000g prior to sample application.
References	<ol style="list-style-type: none"> Shimazu, R. <i>et al.</i> (1999) MD-2, a molecule that confers lipopolysaccharide responsiveness on Toll-like receptor 4. J Exp Med. 189 (11): 1777-82. Sugawara, S. <i>et al.</i> (2000) Proteolysis of human monocyte CD14 by cysteine proteinases (gingipains) from <i>Porphyromonas gingivalis</i> leading to lipopolysaccharide hyporesponsiveness. J Immunol. 165: 411-8. Yang, S. <i>et al.</i> (2001) Synergistic effect of muramyl dipeptide with lipopolysaccharide or lipoteichoic acid to induce inflammatory cytokines in human monocytic cells in culture. Infect Immun. 69 (4): 2045-53. Kawahara T <i>et al.</i> (2001) Type I <i>Helicobacter pylori</i> lipopolysaccharide stimulates toll-like receptor 4 and activates mitogen oxidase 1 in gastric pit cells. Infect Immun. 69 (7): 4382-9.

5. Baumgarten, G. *et al.* (2001) *In vivo* expression of proinflammatory mediators in the adult heart after endotoxin administration: the role of toll-like receptor-4. [J Infect Dis. 183: 1617-24.](#)
6. Medvedev, A.E. *et al.* (2001) Induction of tolerance to lipopolysaccharide and mycobacterial components in Chinese hamster ovary/CD14 cells is not affected by overexpression of Toll-like receptors 2 or 4. [J Immunol. 167: 2257-67.](#)
7. Bieback, K. *et al.* (2002) Hemagglutinin protein of wild-type measles virus activates toll-like receptor 2 signaling. [J Virol. 76: 8729-36.](#)
8. Karlsson, H. *et al.* (2002) Innate immune responses of human neonatal cells to bacteria from the normal gastrointestinal flora. [Infect Immun. 70: 6688-96.](#)
9. Devaney, J.M. (2003) Neutrophil elastase up-regulates interleukin-8 via toll-like receptor 4. [FEBS Lett. 544:129-32.](#)
10. Elner, S.G. *et al.* (2005) TLR4 mediates human retinal pigment epithelial endotoxin binding and cytokine expression. [Invest Ophthalmol Vis Sci. 46 \(12\): 4627-33.](#)
11. Cuschieri, J. *et al.* (2006) Endotoxin tolerance attenuates LPS-induced TLR4 mobilization to lipid rafts: a condition reversed by PKC activation. [J Leukoc Biol. 80: 1289-97.](#)
12. Lindsay, J.O. *et al.* (2006) Clinical, microbiological, and immunological effects of fructo-oligosaccharide in patients with Crohn's disease. [Gut. 55: 348-55.](#)
13. Foster, N. *et al.* (2007) Pivotal advance: vasoactive intestinal peptide inhibits up-regulation of human monocyte TLR2 and TLR4 by LPS and differentiation of monocytes to macrophages. [J Leukoc Biol. 81: 893-903.](#)
14. Pioli, P.A. *et al.* (2007) Estradiol attenuates lipopolysaccharide-induced CXC chemokine ligand 8 production by human peripheral blood monocytes. [J Immunol. 179: 6284-90.](#)
15. Ibeagha-Awemu, E.M. *et al.* (2008) Bacterial lipopolysaccharide induces increased expression of toll-like receptor (TLR) 4 and downstream TLR signaling molecules in bovine mammary epithelial cells. [Vet Res. 39 \(2\): 11.](#)
16. Chochi, K. *et al.* (2008) *Helicobacter pylori* augments growth of gastric cancers via the lipopolysaccharide-toll-like receptor 4 pathway whereas its lipopolysaccharide attenuates antitumor activities of human mononuclear cells. [Clin Cancer Res. 14 \(10\): 2909-17.](#)
17. Brännström, K. *et al.* (2009) The *Schistosoma mansoni* protein Sm16/SmSLP /SmSPO-1 assembles into a nine-subunit oligomer with potential To inhibit Toll-like receptor signaling. [Infect Immun. 77: 1144-54.](#)
18. de Kleer, I. (2010) CD30 Discriminates Heat Shock Protein 60-Induced FOXP3+CD4+ T Cells with a Regulatory Phenotype. [J Immunol. 185\(4\):2071-9.](#)
19. Garbe, K. *et al.* (2012) Plasmacytoid dendritic cells and their Toll-like receptor 9 expression selectively decrease with age. [Hum Immunol. 73 \(5\): 493-7.](#)
20. Komori, H. *et al.* (2012) $\alpha(1)$ -Acid glycoprotein up-regulates CD163 via TLR4/CD14 protein pathway: possible protection against hemolysis-induced oxidative stress. [J Biol Chem. 287 \(36\): 30688-700.](#)
21. Prokhorenko, I. *et al.* (2012) Toll-like receptor 4 in phagocytosis of Escherichia coli by endotoxin-activated human neutrophils in whole blood [Crit Care 16 \(Suppl 3\), P80](#)
22. Sels, J.W. *et al.* (2012) Fractional flow reserve is not associated with inflammatory markers in patients with stable coronary artery disease. [PLoS One. 7: e46356.](#)
23. Maiolini, A. *et al.* (2012) Toll-like receptors 4 and 9 are responsible for the maintenance of the inflammatory reaction in canine steroid-responsive meningitis-arteritis,

- a large animal model for neutrophilic meningitis. [J Neuroinflammation. 9: 226.](#)
24. Kyrova, K. *et al.* (2014) The response of porcine monocyte derived macrophages and dendritic cells to *Salmonella typhimurium* and lipopolysaccharide. [BMC Vet Res. 10: 244.](#)
25. Zwolak, A. *et al.* (2015) Hyperreactivity of Blood Leukocytes in Patients with NAFLD to *ex vivo* Lipopolysaccharide Treatment Is Modulated by Metformin and Phosphatidylcholine but Not by Alpha Ketoglutarate. [PLoS One. 10 \(12\): e0143851.](#)
26. Xu, H. *et al.* (2015) Type 3 innate lymphoid cell depletion is mediated by TLRs in lymphoid tissues of simian immunodeficiency virus-infected macaques. [FASEB J. 29 \(12\): 5072-80.](#)
27. Blagitz, M.G. *et al.* (2015) Expression of CD14 and toll-like receptors 2 and 4 by milk neutrophils in bovine mammary glands infected with *Corynebacterium bovis* [Pesquisa Veterinária Brasileira. 35 \(1\): 1-5.](#)
28. Mazzucchelli, I. *et al.* (2015) Expression and function of toll-like receptors in human circulating endothelial colony forming cells. [Immunol Lett. 168 \(1\): 98-104.](#)
29. Zwolak, A. *et al.* (2016) Metformin Changes the Relationship between Blood Monocyte Toll-Like Receptor 4 Levels and Nonalcoholic Fatty Liver Disease-*Ex Vivo* Studies. [PLoS One. 11 \(3\): e0150233.](#)
30. Huang, D. *et al.* (2016) Hyperoxia induces inflammation and regulates cytokine production in alveolar epithelium through TLR2/4-NF-κB-dependent mechanism [Eur Rev Med Pharmacol Sci. 20: 1399-410.](#)
31. Reineking, W. *et al.* (2018) Canine primary jejunal and colonic epithelial cells predominantly express TLR5 and TLR9 but do not change TLR expression pattern after stimulation with certain Toll-like receptor ligands. [Vet Immunol Immunopathol. 206: 16-24.](#)
32. Awuah, D. *et al.* (2019) The Cross-Talk between miR-511-3p and C-Type Lectin Receptors on Dendritic Cells Affects Dendritic Cell Function. [J Immunol. 203 \(1\): 148-57.](#)

Storage

Store at +4°C. DO NOT FREEZE.
This product should be stored undiluted.

Guarantee

12 months from date of despatch

Acknowledgements

This product is covered by U.S. Patent No. 10,150,841 and related U.S. and foreign counterparts

Health And Safety Information

Material Safety Datasheet documentation #20471 available at: <https://www.bio-rad-antibodies.com/SDS/MCA2061SBV440>

Regulatory

For research purposes only

Related Products

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

Product inquiries: www.bio-rad-antibodies.com/technical-support

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

Printed on 19 May 2026

© 2026 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)