

Datasheet: MCA2071A647

Description:	MOUSE ANTI HUMAN CD80:Alexa Fluor® 647
Specificity:	CD80
Other names:	B7-1
Format:	ALEXA FLUOR® 647
Product Type:	Monoclonal Antibody
Clone:	MEM-233
Isotype:	IgG1
Quantity:	100 TESTS/1ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/5 - 1/10

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human		
Product Form	Purified IgG conjugated to Alexa Fluor® 647 - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	Alexa Fluor®647	650	665
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% sodium azide (NaN ₃)		
Stabilisers	1% bovine serum albumin		
Approx. Protein Concentrations	IgG concentration 0.05 mg/ml		

External Database
Links

UniProt:

[P33681](#) [Related reagents](#)

Entrez Gene:

[941](#) CD80 [Related reagents](#)

Synonyms CD28LG, CD28LG1, LAB7

RRID AB_566899

Specificity

Mouse anti Human CD80 antibody, clone MEM-233 recognizes human CD80, also known as B7-1, a ~60 kDa type 1 trans-membrane protein expressed of macrophages, dendritic cells ([Munro et al. 1994](#)) and activated B-cells ([Ranheim et al. 1993](#)). CD80 is a member of the immunoglobulin superfamily having an extracellular domain bearing both a single [Ig-v-like](#) domain, a single [Ig-c-like](#) domain, a transmembrane sequence and a short cytoplasmic domain. Although the predicted molecular weight for human CD80 is ~33 kDa, the presence of multiple (8) potential N-glycosylation sites ([Chen et al. 1998](#)) results in a migration corresponding to ~60 kDa.

Human CD80 along with CD86 act as co-stimulatory molecules and are both ligands for CD28 and CTLA-4 ([Azuma et al. 1993](#)) involved in T cell activation and proliferation ([Vasu et al. 2003](#)). Although CD80 binds to the same receptors as CD86 it displays quite different characteristics in its avidity and binding kinetics ([van der Merwe et al. 1997](#)).

Mutagenesis indicates residues in both the Ig-v-like and Ig-c-like domains of CD80 are crucial for the interaction with it's receptors CTLA-4 and CD28 ([Peach et al. 1995](#)).

Mouse anti human CD80 antibody, clone MEM-233 binds to residues within the Ig-v-like domain of human CD80 as shown by domain switching assays ([Vasu et al. 2003](#)).

Mouse anti Human CD80, clone MEM-233 in combination with Mouse anti Human CD86, clone Bu63 ([MCA1118](#)) suggest that clone MEM-233 is able to block binding of human CD80 with it's cognate ligands CD28 and CTLA-4 ([Morbach et al. 2011](#)).

Flow Cytometry

Use 10µl of the suggested working dilution to label 10⁶ cells or 100µl whole blood

References

1. Zhan, H. *et al.* (2003) The immunomodulatory role of human conjunctival epithelial cells. [Invest Ophthalmol Vis Sci. 44 \(9\): 3906-10.](#)
2. Tan, P.H. *et al.* (2004) Phenotypic and functional differences between human saphenous vein (HSVEC) and umbilical vein (HUVEC) endothelial cells. [Atherosclerosis. 173: 171-83.](#)
3. Huxley, P. *et al.* (2004) High-affinity small molecule inhibitors of T cell costimulation: compounds for immunotherapy. [Chem Biol. 11: 1651-8.](#)
4. Tan, P.H. *et al.* (2005) Modulation of human dendritic-cell function following transduction with viral vectors: implications for gene therapy. [Blood. 105: 3824-32.](#)
5. Angel, C.E. *et al.* (2006) Cutting edge: CD1a+ antigen-presenting cells in human dermis respond rapidly to CCR7 ligands. [J Immunol. 176 \(10\): 5730-4.](#)

6. Daubenberger, C.A. *et al.* (2007) Flow cytometric analysis on cross-reactivity of human-specific CD monoclonal antibodies with splenocytes of *Aotus nancymaae*, a non-human primate model for biomedical research. [Vet Immunol Immunopathol. 119 \(1-2\): 14-20.](#)
7. Trojan, J. *et al.* (2010) Antisense anti IGF-I cellular therapy of malignant tumours: immune response in cancer patients. [Biomed Pharmacother. 64: 576-8.](#)
8. Piconi, S. *et al.* (2010) Immunological effects of sublingual immunotherapy: clinical efficacy is associated with modulation of programmed cell death ligand 1, IL-10, and IgG4. [J Immunol. 185: 7723-30.](#)
9. John, J. *et al.* (2010) Differential effects of Paclitaxel on dendritic cell function. [BMC Immunol. 11: 14.](#)
10. Hovden, A.O. *et al.* (2011) Maturation of monocyte derived dendritic cells with OK432 boosts IL-12p70 secretion and conveys strong T-cell responses. [BMC Immunol. 12: 2.](#)
11. Silk, K.M. *et al.* (2012) Rapamycin conditioning of dendritic cells differentiated from human ES cells promotes a tolerogenic phenotype. [J Biomed Biotechnol. 2012: 172420.](#)
12. Silk, K.M. *et al.* (2012) Cross-presentation of tumour antigens by human induced pluripotent stem cell-derived CD141+XCR1+ dendritic cells [Gene Ther. 19: 1035-40.](#)
13. Demmers, M.W. *et al.* (2013) Differential effects of activated human renal epithelial cells on T-cell migration. [PLoS One. 8 \(5\): e64916.](#)
14. Scott-Taylor, T.H. *et al.* (2017) Enhanced formation of giant cells in common variable immunodeficiency: Relation to granulomatous disease. [Clin Immunol. 175: 1-9.](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Guarantee 12 months from date of despatch

Acknowledgements This product is provided under an intellectual property licence from Life Technologies Corporation. The transfer of this product is contingent on the buyer using the purchase product solely in research, excluding contract research or any fee for service research, and the buyer must not sell or otherwise transfer this product or its components for (a) diagnostic, therapeutic or prophylactic purposes; (b) testing, analysis or screening services, or information in return for compensation on a per-test basis; (c) manufacturing or quality assurance or quality control, or (d) resale, whether or not resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad CA 92008 USA or outlicensing@thermofisher.com

Health And Safety Information Material Safety Datasheet documentation #10041 available at: <https://www.bio-rad-antibodies.com/SDS/MCA2071A647>
10041

Regulatory For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:Alexa Fluor® 647 \(MCA928A647\)](#)

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M412727:221116'

Printed on 27 Aug 2024

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)