

Datasheet: MCA2086 BATCH NUMBER 150244

| Description: | MOUSE ANTI HUMAN CD18 (ACTIVATION EPITOPE) |
|----------------------|--|
| Specificity: | CD18 (ACTIVATION EPITOPE) |
| Other names: | INTEGRIN BETA 2 CHAIN |
| Format: | Purified |
| Product Type: | Monoclonal Antibody |
| Clone: | MEM-148 |
| Isotype: | lgG1 |
| Quantity: | 0.2 mg |
| | |

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

| | Yes | No | Not Determined | Suggested Dilution |
|----------------------------|-----|----|----------------|-------------------------|
| Flow Cytometry | • | | | |
| Immunohistology - Frozen | | | • | |
| Immunohistology - Paraffin | | | • | |
| ELISA | | | • | |
| Immunoprecipitation | • | | | |
| Western Blotting | • | | | Non-reducing conditions |

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

| Target Species | Human |
|-----------------------------|---|
| Product Form | Purified IgG - liquid |
| Preparation | Purified IgG prepared by affinity chromatography on Protein G |
| Buffer Solution | Phosphate buffered saline |
| Preservative Stabilisers | 0.09% Sodium Azide |

| Approx. Protein Concentrations | IgG concentration 1.0 mg/ml | | | | |
|--------------------------------|---|--|--|--|--|
| Immunogen | Peripheral blood mononuclear cells. | | | | |
| External Database Links | UniProt: P05107 Related reagents | | | | |
| | Entrez Gene: 3689 ITGB2 Related reagents | | | | |
| Synonyms | CD18, MFI7 | | | | |
| RRID | AB_323889 | | | | |
| Specificity | Mayor anti Human CD49 (Activation Enitana) antihady, al | | | | |

Specificity

Mouse anti Human CD18 (Activation Epitope) antibody, clone MEM-148 recognizes an epitope on the human CD18 molecule that is hidden in the CD11/CD18 heterodimer on resting cells. Clone MEM-148 binds very weakly to resting peripheral blood leukocytes and strongly to all leukocytes upon cellular activation. The epitope recognized by Mouse anti Human CD18 (Activation Epitope) antibody, clone MEM-148 is also exposed during dissociation of the CD11/CD18 by low pH.

Flow Cytometry

Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul.

References

- 1. Drbal, K. *et al.* (2001) A proteolytically truncated form of free CD18, the common chain of leukocyte integrins, as a novel marker of activated myeloid cells. <u>Blood. 98 (5): 1561-6.</u>
- 2. Luissint, A.C. *et al.* (2008) JAM-L-mediated leukocyte adhesion to endothelial cells is regulated in cis by alpha4beta1 integrin activation. <u>J Cell Biol. 183 (6): 1159-73.</u>
- 3. Feng, C. *et al.* (2011) Endogenous PMN sialidase activity exposes activation epitope on CD11b/CD18 which enhances its binding interaction with ICAM-1. <u>J Leukoc Biol. 90:</u> 313-21.
- 4. Anogianaki, A. *et al.* (2007) Capsaicin an irritant anti-inflammatory compound. <u>J Biol Regul Homeost Agents</u>. 21: 1-4.
- 5. Arthos, J. *et al.* (2008) HIV-1 envelope protein binds to and signals through integrin alpha4beta7, the gut mucosal homing receptor for peripheral T cells. <u>Nat Immunol. 9:</u> 301-9.
- 6. Cairo, C.W. *et al.* (2006) Cytoskeletal regulation couples LFA-1 conformational changes to receptor lateral mobility and clustering. <u>Immunity. 25: 297-308.</u>
- 7. Tang, X.Y. *et al.* (2008) Intercellular adhesion molecule-3 binding of integrin alphaL beta2 requires both extension and opening of the integrin headpiece. <u>J Immunol. 180:</u> 4793-804.
- 8. Kudlová M. *et al.* (2007) Expression of an activated form of integrin beta2 chain CD18 in cardiac surgical operations. Acta Medica (Hradec Kralove). 50: 187-93.
- 9. Solovjov, D.A. *et al.* (2005) Distinct roles for the alpha and beta subunits in the functions of integrin alphaMbeta2. J Biol Chem. 280: 1336-45.
- 10. Ehirchiou, D. *et al.* (2005) Dual function for a unique site within the beta2l domain of integrin alphaMbeta2. <u>J Biol Chem. 280: 8324-31.</u>

11. Cheng, M. et al. (2007) Mutation of a conserved asparagine in the I-like domain promotes constitutively active integrins alphaLbeta2 and alphaIlbbeta3. J Biol Chem. 282: 18225-32.

12. Shi, M. et al. (2007) A structural hypothesis for the transition between bent and extended conformations of the leukocyte beta2 integrins. J Biol Chem. 282: 30198-206. 13. Dilek, N. et al. (2013) Targeting CD28, CTLA-4 and PD-L1 costimulation differentially controls immune synapses and function of human regulatory and conventional T-cells. PLoS One. 8 (12): e83139.

Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

| Guarantee | 12 months from date of despatch |
|----------------------------------|--|
| Health And Safety Information | Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA2086 10040 |
| Regulatory | For research purposes only |

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...) **RPE**

Goat Anti Mouse IgG IgA IgM (STAR87...) HRP

Goat Anti Mouse IgG (STAR76...) **RPE**

Rabbit Anti Mouse IgG (STAR13...) **HRP**

Goat Anti Mouse IgG (STAR70...) **FITC**

Goat Anti Mouse IgG (H/L) (STAR117...) Alk. Phos., DyLight®488, DyLight®550,

DyLight®650, DyLight®680, DyLight®800,

FITC, HRP

Rabbit Anti Mouse IgG (STAR9...) **FITC**

Goat Anti Mouse IgG (STAR77...) **HRP**

Goat Anti Mouse IgG (Fc) (STAR120...) FITC, HRP

Recommended Negative Controls

Fax: +1 919 878 3751

MOUSE IgG1 NEGATIVE CONTROL (MCA928)

North & South Tel: +1 800 265 7376

America

Worldwide

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_us@bio-rad.com

Email: antibody sales uk@bio-rad.com

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M366146:200529'

Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint