

Datasheet: MCA2086

BATCH NUMBER 150244

Description:	MOUSE ANTI HUMAN CD18 (ACTIVATION EPITOPE)
Specificity:	CD18 (ACTIVATION EPITOPE)
Other names:	INTEGRIN BETA 2 CHAIN
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	MEM-148
Isotype:	IgG1
Quantity:	0.2 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting	▪			Non-reducing conditions

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide

Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Peripheral blood mononuclear cells.
External Database Links	<p>UniProt: P05107 Related reagents</p> <p>Entrez Gene: 3689 ITGB2 Related reagents</p>
Synonyms	CD18, MFI7
RRID	AB_323889
Specificity	<p>Mouse anti Human CD18 (Activation Epitope) antibody, clone MEM-148 recognizes an epitope on the human CD18 molecule that is hidden in the CD11/CD18 heterodimer on resting cells. Clone MEM-148 binds very weakly to resting peripheral blood leukocytes and strongly to all leukocytes upon cellular activation. The epitope recognized by Mouse anti Human CD18 (Activation Epitope) antibody, clone MEM-148 is also exposed during dissociation of the CD11/CD18 by low pH.</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
References	<ol style="list-style-type: none"> Drbal, K. <i>et al.</i> (2001) A proteolytically truncated form of free CD18, the common chain of leukocyte integrins, as a novel marker of activated myeloid cells. Blood. 98 (5): 1561-6. Luissint, A.C. <i>et al.</i> (2008) JAM-L-mediated leukocyte adhesion to endothelial cells is regulated in cis by alpha4beta1 integrin activation. J Cell Biol. 183 (6): 1159-73. Feng, C. <i>et al.</i> (2011) Endogenous PMN sialidase activity exposes activation epitope on CD11b/CD18 which enhances its binding interaction with ICAM-1. J Leukoc Biol. 90: 313-21. Anogianaki, A. <i>et al.</i> (2007) Capsaicin an irritant anti-inflammatory compound. J Biol Regul Homeost Agents. 21: 1-4. Arthos, J. <i>et al.</i> (2008) HIV-1 envelope protein binds to and signals through integrin alpha4beta7, the gut mucosal homing receptor for peripheral T cells. Nat Immunol. 9: 301-9. Cairo, C.W. <i>et al.</i> (2006) Cytoskeletal regulation couples LFA-1 conformational changes to receptor lateral mobility and clustering. Immunity. 25: 297-308. Tang, X.Y. <i>et al.</i> (2008) Intercellular adhesion molecule-3 binding of integrin alphaL beta2 requires both extension and opening of the integrin headpiece. J Immunol. 180: 4793-804. Kudlová M. <i>et al.</i> (2007) Expression of an activated form of integrin beta2 chain CD18 in cardiac surgical operations. Acta Medica (Hradec Kralove). 50: 187-93. Solovjov, D.A. <i>et al.</i> (2005) Distinct roles for the alpha and beta subunits in the functions of integrin alphaMbeta2. J Biol Chem. 280: 1336-45. Ehrichtiou, D. <i>et al.</i> (2005) Dual function for a unique site within the beta2I domain of integrin alphaMbeta2. J Biol Chem. 280: 8324-31.

11. Cheng, M. *et al.* (2007) Mutation of a conserved asparagine in the I-like domain promotes constitutively active integrins alphaLbeta2 and alphaIIb beta3. [J Biol Chem. 282: 18225-32.](#)
12. Shi, M. *et al.* (2007) A structural hypothesis for the transition between bent and extended conformations of the leukocyte beta2 integrins. [J Biol Chem. 282: 30198-206.](#)
13. Dilek, N. *et al.* (2013) Targeting CD28, CTLA-4 and PD-L1 costimulation differentially controls immune synapses and function of human regulatory and conventional T-cells. [PLoS One. 8 \(12\): e83139.](#)

Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee

12 months from date of despatch

Health And Safety Information

Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/MCA2086>
10040

Regulatory

For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...)	RPE
Goat Anti Mouse IgG IgA IgM (STAR87...)	HRP
Goat Anti Mouse IgG (STAR76...)	RPE
Goat Anti Mouse IgG (STAR70...)	FITC
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight®488 , DyLight®550 , DyLight®650 , DyLight®680 , DyLight®800 , FITC , HRP
Goat Anti Mouse IgG (STAR77...)	HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP
Rabbit Anti Mouse IgG (STAR13...)	HRP

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

North & South America

Tel: +1 800 265 7376
Fax: +1 919 878 3751
Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700
Fax: +44 (0)1865 852 739
Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21
Fax: +49 (0) 89 8090 95 50
Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)
'M366146:200529'

