

Datasheet: MCA2087PET

Description:	MOUSE ANTI HUMAN CD11c:RPE
Specificity:	CD11c
Other names:	INTEGRIN ALPHA X CHAIN
Format:	RPE
Product Type:	Monoclonal Antibody
Clone:	BU15
Isotype:	IgG1
Quantity:	25 TESTS/0.25ml

Product Details

RRID AB_2129804

Applications This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat - 1/10

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species Human

Species Cross Reactivity Reacts with: Cynomolgus monkey
N.B. Antibody reactivity and working conditions may vary between species.

Product Form Purified IgG conjugated to R. Phycoerythrin (RPE) - liquid

Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	RPE 488nm laser	496	578

Preparation Purified IgG prepared from tissue culture supernatant

Buffer Solution Phosphate buffered saline

Preservative 0.09% Sodium Azide (NaN₃)
Stabilisers 1% Bovine Serum Albumin
5% Sucrose

External Database Links

UniProt:
[P20702](#) [Related reagents](#)

Entrez Gene:

[3687](#) ITGAX [Related reagents](#)

Synonyms

CD11C

Specificity

Mouse anti Human CD11c antibody, clone Bu15 recognizes the integrin alpha^x subunit (ITGAX), a ~150 kDa glycoprotein also known as CD11 antigen-like family member C or Leu M5, CD11c is expressed by macrophages, monocytes, NK cells and most dendritic cells ([Kohrgruber et al. 1999](#)). CD11c is also expressed at a lower level by granulocytes.

CD11c forms a heterodimeric integral membrane protein with the integrin beta 2 chain to form the leukocyte specific integrin '[inactivated-C3b receptor 4](#)'. CD11c interacts with a number of ligands including the [G-P-R](#) sequence in fibrinogen, ICAM-1 ([Frick et al. 2005](#)), iC3b ([Sadhu et al. 2007](#)) and Junctional adhesion molecule-like ([Bazzoni et al. 2011](#)). CD11c is involved in cell-cell interactions during the inflammatory process and is important for monocyte adhesion and chemotaxis. CD11c also acts as a signalling receptor for polysaccharide ([Inqualls et al. 1995](#))

CD11c is expressed by hairy cell leukaemia cells ([Goodman et al. 2003](#); [Nicolaou et al. 2003](#)).

Flow Cytometry

Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul.

References

1. Hogg, N. *et al.* (1986) The p150,95 molecule is a marker of human mononuclear phagocytes: comparison with expression of class II molecules. [Eur J Immunol. 16 \(3\): 240-8.](#)
2. Loike, J.D. *et al.* (1991) CD11c/CD18 on neutrophils recognizes a domain at the N terminus of the A alpha chain of fibrinogen. [Proc Natl Acad Sci U S A. 88 \(3\): 1044-8.](#)
3. Newman, K.C. *et al.* (2006) Cross-talk with myeloid accessory cells regulates human natural killer cell interferon-gamma responses to malaria. [PLoS Pathog. 2: e118.](#)
4. Sanchez-Madrid, F. and Corbi, A.L. (1992) Leukocyte integrins: structure, function and regulation of their activity. [Seminars Cell Biol. 3: 199-210.](#)
5. Silk, K.M. *et al.* (2012) Rapamycin conditioning of dendritic cells differentiated from human ES cells promotes a tolerogenic phenotype. [J Biomed Biotechnol. 2012:172420.](#)
6. Bird, R.C. *et al.* (2011) An autologous dendritic cell canine mammary tumor hybrid-cell fusion vaccine. [Cancer Immunol Immunother. 60 \(1\): 87-97.](#)
7. Eisenhardt, S.U. *et al.* (2007) Generation of activation-specific human anti-alphaMbeta2 single-chain antibodies as potential diagnostic tools and therapeutic agents. [Blood. 109: 3521-8.](#)
8. Herman, S. *et al.* (2012) Regulatory T cells form stable and long-lasting cell cluster with myeloid dendritic cells (DC). [Int Immunol. 24 \(7\): 417-26.](#)
9. Levy, O. *et al.* (2003) Critical role of the complement system in group B streptococcus-induced tumor necrosis factor alpha release. [Infect Immun. 71: 6344-53.](#)
10. Xie, Z. *et al.* (2016) Human umbilical cord-derived mesenchymal stem cells elicit macrophages into an anti-inflammatory phenotype to alleviate insulin resistance in type 2 diabetic rats. [Stem Cells. 34 \(3\): 627-39.](#)
11. Brown, D.P. *et al.* (2009) The inhibitory receptor LILRB4 (ILT3) modulates antigen presenting cell phenotype and, along with LILRB2 (ILT4), is upregulated in response to *Salmonella* infection. [BMC Immunol. 10: 56.](#)
12. Schroeder JH *et al.* (2017) *Brugia malayi* microfilariae adhere to human vascular endothelial cells in a C3-dependent manner. [PLoS Negl Trop Dis. 11 \(5\): e0005592.](#)

Further Reading

1. Larson, R.S. & Springer, T.A. (1990) Structure and function of leukocyte integrins. [Immunol Rev. 114: 181-217.](#)

Storage

Store at +4°C. DO NOT FREEZE.

This product should be stored undiluted. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee	12 months from date of despatch.
------------------	----------------------------------

Health And Safety Information	Material Safety Datasheet documentation #10306 available at: 10306: https://www.bio-rad-antibodies.com/uploads/MSDS/10306.pdf
--------------------------------------	----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Regulatory	For research purposes only
-------------------	----------------------------

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:RPE \(MCA928PE\)](#)

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	-----------------------------------------------------------------------------------------------------------------------------------------	------------------	-------------------------------------------------------------------------------------------------------------------------------------------------	---------------	-----------------------------------------------------------------------------------------------------------------------------------------------------

'M343074:190110'

Printed on 20 May 2019

© 2019 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)