

## Datasheet: MCA2087SBV790

**BATCH NUMBER 64595345**

<b>Description:</b>	MOUSE ANTI HUMAN CD11c:StarBright Violet 790
<b>Specificity:</b>	CD11c
<b>Other names:</b>	INTEGRIN ALPHA X CHAIN
<b>Format:</b>	StarBright Violet 790
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	BU15
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	100 TESTS/0.5ml

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

**Target Species** Human

**Species Cross Reactivity** Reacts with: Cynomolgus monkey  
**N.B.** Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

**Product Form** Purified IgG conjugated to StarBright Violet 790 - liquid

Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	StarBright Violet 790	402	782

**Preparation** Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant

<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative</b>	0.09% Sodium Azide (NaN <sub>3</sub> )
<b>Stabilisers</b>	1% Bovine Serum Albumin 0.1% Pluronic F68 0.1% PEG 3350 0.05% Tween 20
<b>External Database Links</b>	<p><b>UniProt:</b> <a href="#">P20702</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b> <a href="#">3687</a>    ITGAX    <a href="#">Related reagents</a></p>
<b>Synonyms</b>	CD11C
<b>Specificity</b>	<p><b>Mouse anti Human CD11c antibody, clone Bu15</b> recognizes the integrin alpha<sup>x</sup> subunit (ITGAX), a ~150 kDa glycoprotein also known as CD11 antigen-like family member C or Leu M5, CD11c is expressed by macrophages, monocytes, NK cells and most dendritic cells (<a href="#">Kohrgruber et al. 1999</a>). CD11c is also expressed at a lower level by granulocytes.</p> <p>CD11c forms a heterodimeric integral membrane protein with the integrin beta 2 chain to form the leukocyte specific integrin '<a href="#">inactivated-C3b receptor 4</a>'. CD11c interacts with a number of ligands including the <a href="#">G-P-R</a> sequence in fibrinogen, ICAM-1 (<a href="#">Frick et al. 2005</a>), iC3b (<a href="#">Sadhu et al. 2007</a>) and Junctional adhesion molecule-like (<a href="#">Bazzoni et al. 2011</a>). CD11c is involved in cell-cell interactions during the inflammatory process and is important for monocyte adhesion and chemotaxis. CD11c also acts as a signalling receptor for polysaccharide (<a href="#">Inqualls et al. 1995</a>)</p> <p>CD11c is expressed by hairy cell leukaemia cells (<a href="#">Goodman et al. 2003</a>; <a href="#">Nicolaou et al. 2003</a>).</p>
<b>Flow Cytometry</b>	Use 5µl of the suggested working dilution to label 10 <sup>6</sup> cells in 100µl. Best practices suggest a 5 minutes centrifugation at 6,000g prior to sample application.
<b>References</b>	<ol style="list-style-type: none"> <li>Hogg, N. <i>et al.</i> (1986) The p150,95 molecule is a marker of human mononuclear phagocytes: comparison with expression of class II molecules. <a href="#">Eur J Immunol. 16 (3): 240-8.</a></li> <li>Levy, O. <i>et al.</i> (2003) Critical role of the complement system in group B streptococcus-induced tumor necrosis factor alpha release. <a href="#">Infect Immun. 71: 6344-53.</a></li> <li>Newman, K.C. <i>et al.</i> (2006) Cross-talk with myeloid accessory cells regulates human natural killer cell interferon-gamma responses to malaria. <a href="#">PLoS Pathog. 2: e118.</a></li> <li>Brown, D.P. <i>et al.</i> (2009) The inhibitory receptor LILRB4 (ILT3) modulates antigen presenting cell phenotype and, along with LILRB2 (ILT4), is upregulated in response to <i>Salmonella</i> infection. <a href="#">BMC Immunol. 10: 56.</a></li> <li>Silk, K.M. <i>et al.</i> (2012) Rapamycin conditioning of dendritic cells differentiated from human ES cells promotes a tolerogenic phenotype. <a href="#">J Biomed Biotechnol. 2012:172420.</a></li> </ol>

6. Herman, S. *et al.* (2012) Regulatory T cells form stable and long-lasting cell cluster with myeloid dendritic cells (DC). [Int Immunol. 24 \(7\): 417-26.](#)
7. Xie, Z. *et al.* (2016) Human umbilical cord-derived mesenchymal stem cells elicit macrophages into an anti-inflammatory phenotype to alleviate insulin resistance in type 2 diabetic rats. [Stem Cells. 34 \(3\): 627-39.](#)
8. Schroeder JH *et al.* (2017) *Brugia malayi* microfilariae adhere to human vascular endothelial cells in a C3-dependent manner. [PLoS Negl Trop Dis. 11 \(5\): e0005592.](#)
9. Simões, R.D. *et al.* (2019) Effects of Regulatory T Cell Depletion on NK Cell Responses against *Listeria monocytogenes* in Feline Immunodeficiency Virus Infected Cats. [Viruses. 11 \(11\)Oct 24 \[Epub ahead of print\].](#)
10. de Vries, R.D. *et al.* (2014) Measles vaccination of nonhuman primates provides partial protection against infection with canine distemper virus. [J Virol. 88 \(8\): 4423-33.](#)

---

#### Further Reading

1. Larson, R.S. & Springer, T.A. (1990) Structure and function of leukocyte integrins. [Immunol Rev. 114: 181-217.](#)
2. Loike, J.D. *et al.* (1991) CD11c/CD18 on neutrophils recognizes a domain at the N terminus of the A alpha chain of fibrinogen. [Proc Natl Acad Sci U S A. 88 \(3\): 1044-8.](#)
3. Sanchez-Madrid, F. and Corbi, A.L. (1992) Leukocyte integrins: structure, function and regulation of their activity. [Seminars Cell Biol. 3: 199-210.](#)

---

#### Storage

Store at +4°C. DO NOT FREEZE.  
This product should be stored undiluted.

---

#### Guarantee

12 months from date of despatch

---

#### Acknowledgements

This product is covered by U.S. Patent No. 10,150,841 and related U.S. and foreign counterparts

---

#### Health And Safety Information

Material Safety Datasheet documentation #20471 available at:  
<https://www.bio-rad-antibodies.com/SDS/MCA2087SBV790>  
20471

---

#### Regulatory

For research purposes only

---

## Related Products

### Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

**North & South America** Tel: +1 800 265 7376

Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](http://bio-rad-antibodies.com/datasheets)

'M429781:240429'

Printed on 31 Jan 2025