

Datasheet: MCA2090

BATCH NUMBER 161883

Description:	MOUSE ANTI HUMAN HLA A2
Specificity:	HLA A2
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	BB7.2
Isotype:	IgG2b
Quantity:	0.2 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			10ug/ml
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant.
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide
Carrier Free	Yes

Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Papain solubilized HLA-A2
External Database Links	<p>UniProt: P01892 Related reagents</p> <p>Entrez Gene: 3105 HLA-A Related reagents</p>
Synonyms	HLAA
RRID	AB_324652
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of the mouse NS1 myeloma cell line
Specificity	<p>Mouse anti Human HLA A2 antibody, clone BB7.2 recognizes the human HLA-A2 histocompatibility antigen. The epitope recognized by this antibody has been studied extensively and would appear to include the carboxy-terminus of the alpha-2 helix and a turn on one of the underlying beta strands.</p> <p>Mouse anti Human HLA A2 antibody, clone BB7.2 may be used for the flow cytometric detection of HLA-A2 expression and has also been used for immunoaffinity purification of HLA-A2 molecules. Functionally Mouse anti Human HLA A2 antibody, clone BB7.2 is reported to inhibit MHC restricted cellular cytotoxicity.</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells or 100ul whole blood.
Histology Positive Control Tissue	Tonsil
References	<ol style="list-style-type: none"> Parham, P. & Brodsky, F.M. (1981) Partial purification and some properties of BB7.2. A cytotoxic monoclonal antibody with specificity for HLA-A2 and a variant of HLA-A28. Hum Immunol. 3 (4): 277-99. Hogan, K.T. & Brown, S.L. (1992) Localization and characterization of serologic epitopes on HLA-A2. Hum Immunol. 33 (3): 185-92. Harig, S. <i>et al.</i> (2001) Induction of cytotoxic T-cell responses against immunoglobulin V region-derived peptides modified at human leukocyte antigen-A2 binding residues. Blood. 98 (10): 2999-3005. Pantenburg, B. <i>et al.</i> (2010) Human CD8(+) T cells clear <i>Cryptosporidium parvum</i> from infected intestinal epithelial cells. Am J Trop Med Hyg. 82:600-7. Duncan, L.M. <i>et al.</i> (2010) Stabilization of an E3 ligase-E2-ubiquitin complex increases cell surface MHC class I expression. J Immunol. 184: 6978-85. Wang, B. <i>et al.</i> (2004) Identification of an HLA-A*0201-restricted CD8+ T-cell epitope SSp-1 of SARS-CoV spike protein. Blood. 104: 200-6.

7. Wooldridge, L. *et al.* (2007) Enhanced immunogenicity of CTL antigens through mutation of the CD8 binding MHC class I invariant region. [Eur J Immunol. 37: 1323-33.](#)
8. Wooldridge, L. *et al.* (2010) MHC class I molecules with Superenhanced CD8 binding properties bypass the requirement for cognate TCR recognition and nonspecifically activate CTLs. [J Immunol. 184: 3357-66.](#)
9. Popovic, J. *et al.* (2011) The only proposed T-cell epitope derived from the TEL-AML1 translocation is not naturally processed. [Blood. 118 \(4\): 946-54.](#)
10. Schmidt, J. *et al.* (2011) Reversible Major Histocompatibility Complex I-Peptide Multimers Containing Ni²⁺-Nitrilotriacetic Acid Peptides and Histidine Tags Improve Analysis and Sorting of CD8⁺ T Cells. [J Biol Chem. 286: 41723-35.](#)
11. Brooks, S.E. *et al.* (2015) Application of the pMHC Array to Characterise Tumour Antigen Specific T Cell Populations in Leukaemia Patients at Disease Diagnosis. [PLoS One. 10 \(10\): e0140483.](#)
12. Rothe, K. *et al.* (2016) Latent Cytomegalovirus Infection in Rheumatoid Arthritis and Increased Frequencies of Cytolytic LIR-1+CD8⁺ T Cells. [Arthritis Rheumatol. 68 \(2\): 337-46.](#)
13. De Angelis Rigotti, F. *et al.* (2017) MARCH9-mediated ubiquitination regulates MHC I export from the TGN. [Immunol Cell Biol. 95 \(9\): 753-64.](#)
14. Kim, J.Y. *et al.* (2017) Detection of Donor-Derived Microparticles in the Peripheral Blood of a Hand Transplant Recipient During Rejection. [Transplant Direct. 3 \(3\): e131.](#)
15. Dockree, T. *et al.* (2017) CD8⁺ T-cell specificity is compromised at a defined MHCI/CD8 affinity threshold. [Immunol Cell Biol. 95 \(1\): 68-76.](#)
16. Sun, W. *et al.* (2018) A modified HLA-A*0201-restricted CTL epitope from human oncoprotein (hPEBP4) induces more efficient antitumor responses. [Cell Mol Immunol. 15 \(8\): 768-81.](#)
17. Obenaus, M. *et al.* (2015) Identification of human T-cell receptors with optimal affinity to cancer antigens using antigen-negative humanized mice. [Nat Biotechnol. 33 \(4\): 402-7.](#)
18. Walseng, E. *et al.* (2015) Soluble T-cell receptors produced in human cells for targeted delivery. [PLoS One. 10 \(4\): e0119559.](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/MCA2090>
10040

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR77...)	HRP
Rabbit Anti Mouse IgG (STAR12...)	RPE
Goat Anti Mouse IgG (STAR70...)	FITC
Goat Anti Mouse IgG IgA IgM (STAR87...)	Alk. Phos. , HRP
Goat Anti Mouse IgG (STAR76...)	RPE
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight®488 , DyLight®550 , DyLight®650 , DyLight®680 , DyLight®800 , FITC , HRP
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP
Rabbit Anti Mouse IgG (STAR13...)	HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC

Recommended Negative Controls

[MOUSE IgG2b NEGATIVE CONTROL \(MCA691\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M383726:210513'

Printed on 02 Jun 2023