

Datasheet: MCA2108A647T

Description:	MOUSE ANTI HUMAN BLTR:Alexa Fluor® 647
Specificity:	BLTR
Other names:	LEUKOTRIENE B4 RECEPTOR
Format:	ALEXA FLUOR® 647
Product Type:	Monoclonal Antibody
Clone:	202/7B1
Isotype:	lgG2a
Quantity:	25 TESTS/0.25ml

Product Details

RRID	AB_1100486
------	------------

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	-			1/5 - 1/10

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positiv controls.

	3 1			
Target Species	Human			
Product Form	Purified IgG conjugate	ed to Alexa Fluor® 64	7- liquid	
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)	
	Alexa Fluor®647	650	665	
Preparation	Purified IgG prepared	by affinity chromatog	raphy on Protein G fror	n tissue culture supernatant
Buffer Solution	Phosphate buffered s	aline		
Preservative	0.09%			
Stabilisers	1%			
	Sodium Azide			
	Bovine Serum Albumi	in		
Approx. Protein Concentrations	IgG concentration 0.0	5 mg/ml		
lmmunogen	HeLa cell transfectan	ts expressing human	BLTR.	

External Database Links

UniProt:

Q15722 Related reagents

Entrez Gene:

1241 LTB4R Related reagents

Synonyms

BLT, BLT1, BLTR, CMKRL1, GPR16, P2RY7

Fusion Partners

Spleen cells from immunised Balb/c mice were fused with cells of the mouse SP2/0 myeloma cell line

Specificity

Mouse anti Human BLTR antibody, clone 202/7B1 recognizes Human leukotriene B4 receptor, also known as BLTR, Chemoattractant receptor-like 1, G-protein coupled receptor 16 or P2Y purinoceptor 7. BLTR is a 353 ~40 kDa multipass transmembrane glycoprotein

BLTR is expressed by peripheral blood granulocytes and is important in the pro-inflammatory response.

Flow Cytometry

Use 10ul of the suggested working dilution to label 10⁶ cells or 100ul whole blood.

References

- 1. Pettersson, A. *et al.* (2000) First-generation monoclonal antibodies identifying the human leukotriene B(4) receptor-1. <u>Biochem Biophys Res Commun.</u> 279 (2): 520-5.
- 2. Costa, M.F. *et al.* (2010) Leukotriene B4 mediates gammadelta T lymphocyte migration in response to diverse stimuli. <u>J Leukoc Biol. 87: 323-32.</u>
- 3. Islam, S.A. *et al.* (2006) The leukotriene B4 lipid chemoattractant receptor BLT1 defines antigen-primed T cells in humans. Blood. 107: 444-53.
- 4. Medoff, B.D. *et al.* (2005) BLT1-mediated T cell trafficking is critical for rejection and obliterative bronchiolitis after lung transplantation. <u>J Exp Med. 202: 97-110.</u>
- 5. Serezani, C.H. *et al.* (2009) FcgammaRI ligation leads to a complex with BLT1 in lipid rafts that enhances rat lung macrophage antimicrobial functions. <u>Blood. 114: 3316-24.</u>
- 6. Heller, E.A. *et al.* (2005) Inhibition of atherogenesis in BLT1-deficient mice reveals a role for LTB4 and BLT1 in smooth muscle cell recruitment. <u>Circulation</u>. 112: 578-86.
- 7. Thomas, S.Y. *et al.* (2007) Multiple chemokine receptors, including CCR6 and CXCR3, regulate antigen-induced T cell homing to the human asthmatic airway. <u>J Immunol. 179: 1901-12.</u>
- 8. Sabirsh, A. *et al.* (2005) Fluorescent leukotriene B4: potential applications. <u>J Lipid Res. 46:</u> 1339-46.
- 9. Loell, I. *et al.* (2012) Activated LTB4 pathway in muscle tissue of patients with polymyositis or dermatomyositis. <u>Ann Rheum Dis. 72: 293-9.</u>
- 10. Wang, L. *et al.* (2012) BLT1-Dependent Alveolar Recruitment of CD4+CD25+ Foxp3+ Regulatory T Cells is Important for Resolution of Acute Lung Injury. <u>Am J Respir Crit Care Med.186: 989-98.</u>
- 11. Laarman, A.J. *et al.* (2012) *Staphylococcus aureus* Staphopain A inhibits CXCR2-dependent neutrophil activation and chemotaxis. <u>EMBO J. 31: 3607-19.</u>
- 12. Dixit, N. *et al.* (2014) Leukotriene B4 activates intracellular calcium and augments human osteoclastogenesis. Arthritis Res Ther. 16: 496.

Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend

microcentrifugation before use.

Guarantee	18 months from date of despatch.
Acknowledgements	This product is provided under an intellectual property licence from Life Technologies Corporation. The transfer of this product is contingent on the buyer using the purchase product solely in research, excluding contract research or any fee for service research, and the buyer must not sell or otherwise transfer this product or its components for (a) diagnostic, therapeutic or prophylactic purposes; (b) testing, analysis or screening services, or information in return for compensation on a per-test basis; (c) manufacturing or quality assurance or quality control, or (d) resale, whether or not resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad CA 92008 USA or outlicensing@thermofisher.com
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: 10041: https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

MOUSE IgG2a NEGATIVE CONTROL:Alexa Fluor® 647 (MCA929A647)

Worldwide

Recommended Useful Reagents

HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

North & South Tel: +1 800 265 7376

America

Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M345193:190125'

Printed on 11 Oct 2019

© 2019 Bio-Rad Laboratories Inc | Legal | Imprint