

Datasheet: MCA2108PE

Description:	MOUSE ANTI HUMAN BLTR:RPE
Specificity:	BLTR
Other names:	LEUKOTRIENE B4 RECEPTOR
Format:	RPE
Product Type:	Monoclonal Antibody
Clone:	202/7B1
Isotype:	IgG2a
Quantity:	100 TESTS

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat - 1/2

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive control.

Target Species	Human		
Product Form	Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilized		
Reconstitution	Reconstitute with 1 ml distilled water		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	RPE 488nm laser	496	578
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative Stabilisers	0.09% Sodium Azide		
	1%	Bovine Serum Albumin	
	5%	Sucrose	

Immunogen	HeLa cell transfectants expressing human BLTR.
External Database Links	<p>UniProt: Q15722 Related reagents</p> <p>Entrez Gene: 1241 LTB4R Related reagents</p>
Synonyms	BLT, BLT1, BLTR, CMKRL1, GPR16, P2RY7
RRID	AB_566388
Fusion Partners	Spleen cells from immunised Balb/c mice were fused with cells of the mouse SP2/0 myeloma cell line.
Specificity	<p>Mouse anti Human BLTR antibody, clone 202/7B1 recognizes Human leukotriene B4 receptor, also known as BLTR, Chemoattractant receptor-like 1, G-protein coupled receptor 16 or P2Y purinoceptor 7. BLTR is a 353 ~40 kDa multipass transmembrane glycoprotein</p> <p>BLTR is expressed by peripheral blood granulocytes and is important in the pro-inflammatory response.</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells or 100ul whole blood.
References	<ol style="list-style-type: none"> 1. Pettersson, A. <i>et al.</i> (2000) First-generation monoclonal antibodies identifying the human leukotriene B(4) receptor-1. Biochem Biophys Res Commun. 279 (2): 520-5. 2. Costa, M.F. <i>et al.</i> (2010) Leukotriene B4 mediates gammadelta T lymphocyte migration in response to diverse stimuli. J Leukoc Biol. 87: 323-32. 3. Islam, S.A. <i>et al.</i> (2006) The leukotriene B4 lipid chemoattractant receptor BLT1 defines antigen-primed T cells in humans. Blood. 107: 444-53. 4. Medoff, B.D. <i>et al.</i> (2005) BLT1-mediated T cell trafficking is critical for rejection and obliterative bronchiolitis after lung transplantation. J Exp Med. 202: 97-110. 5. Serezani, C.H. <i>et al.</i> (2009) FcgammaRI ligation leads to a complex with BLT1 in lipid rafts that enhances rat lung macrophage antimicrobial functions. Blood. 114: 3316-24. 6. Heller, E.A. <i>et al.</i> (2005) Inhibition of atherogenesis in BLT1-deficient mice reveals a role for LTB4 and BLT1 in smooth muscle cell recruitment. Circulation. 112: 578-86. 7. Thomas, S.Y. <i>et al.</i> (2007) Multiple chemokine receptors, including CCR6 and CXCR3, regulate antigen-induced T cell homing to the human asthmatic airway. J Immunol. 179: 1901-12. 8. Sabirsh, A. <i>et al.</i> (2005) Fluorescent leukotriene B4: potential applications. J Lipid Res. 46: 1339-46. 9. Loell, I. <i>et al.</i> (2012) Activated LTB4 pathway in muscle tissue of patients with polymyositis or dermatomyositis. Ann Rheum Dis. 72: 293-9. 10. Wang, L. <i>et al.</i> (2012) BLT1-Dependent Alveolar Recruitment of CD4+CD25+ Foxp3+ Regulatory T Cells is Important for Resolution of Acute Lung Injury. Am J Respir Crit Care Med. 186: 989-98.

11. Laarman, A.J. *et al.* (2012) *Staphylococcus aureus* Staphopain A inhibits CXCR2-dependent neutrophil activation and chemotaxis. [EMBO J. 31: 3607-19.](#)
12. Dixit, N. *et al.* (2014) Leukotriene B4 activates intracellular calcium and augments human osteoclastogenesis. [Arthritis Res Ther. 16: 496.](#)
13. Sezin, T. *et al.* (2019) Dual inhibition of complement factor 5 and leukotriene B4 synergistically suppresses murine pemphigoid disease. [JCI Insight. 4 \(15\)Aug 08 \[Epub ahead of print\].](#)

Storage Prior to reconstitution store at +4°C. Following reconstitution store at +4°C.

DO NOT FREEZE.

This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10075 available at: 10075: <https://www.bio-rad-antibodies.com/uploads/MSDS/10075.pdf>

Regulatory For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG2a NEGATIVE CONTROL:RPE \(MCA929PE\)](#)

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

From March 15, 2021, we will no longer supply printed datasheets with our products. Look out for updates on how to access your digital version at bio-rad-antibodies.com

'M374555:201112'

Printed on 10 Feb 2021