

Datasheet: MCA2112

Description:	MOUSE ANTI BOVINE INTERFERON GAMMA
Specificity:	IFN GAMMA
Other names:	Immune interferon
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	CC330
Isotype:	IgG1
Quantity:	0.5 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry (1)	▪			1/50 - 1/100
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA	▪			5µg/ml as capture antibody
Immunoprecipitation			▪	
Western Blotting			▪	
ELISpot	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1) Membrane permeabilisation is required for this application. Bio-Rad recommends the use of Leucoperm (Product Code [BUF09](#)) for this purpose.

Target Species	Bovine
Species Cross Reactivity	<p>Reacts with: Sheep, Goat</p> <p>N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.</p>

Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% sodium azide (NaN ₃)
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Recombinant bovine IFN-gamma.
External Database Links	<p>UniProt: P07353 Related reagents</p> <p>Entrez Gene: 281237 IFNG Related reagents</p>
RRID	AB_2123456
Specificity	Mouse anti Bovine interferon γ antibody, clone CC330 recognizes bovine interferon gamma, also known as immune interferon. Interferon γ is a 143 amino acid ~20 kDa immunomodulatory cytokine secreted predominantly by T lymphocytes in response to infection by invading microorganisms. Interferon γ enhances cytotoxic activities of many cell types including macrophages, NK cells and T lymphocytes (Weiss et al. 2002).
Flow Cytometry	Use 10µl of the suggested working dilution to label 10 ⁶ cells or 100µl whole blood
ELISA	Mouse anti Bovine interferon-γ antibody, clone CC330 may be used as a capture reagent in a sandwich ELISA in combination with Mouse anti Bovine interferon-γ antibody, clone CC302 (MCA1783B) as a detection reagent and recombinant bovine interferon-γ (PBP007A) as a protein standard.
References	<ol style="list-style-type: none"> 1. Mwangi, W. <i>et al.</i> (2002) DNA-encoded fetal liver tyrosine kinase 3 ligand and granulocyte macrophage-colony-stimulating factor increase dendritic cell recruitment to the inoculation site and enhance antigen-specific CD4+ T cell responses induced by DNA vaccination of outbred animals. J Immunol. 169 (7): 3837-46. 2. Baszler, T.V. <i>et al.</i> (2008) Bovine immune response to inoculation with <i>Neospora caninum</i> surface antigen SRS2 lipopeptides mimics immune response to infection with live parasites. Clin Vaccine Immunol. 15: 659-67. 3. Pascale, F. <i>et al.</i> (2008) Plasmacytoid dendritic cells migrate in afferent skin lymph. J Immunol. 180: 5963-72. 4. Tourais-Esteves, I. <i>et al.</i> (2008) Neonatal goats display a stronger TH1-type cytokine

- response to TLR ligands than adults [Dev Comp Immunol. 32: 1231-41.](#)
5. Begg, D.J. *et al.* (2009) Enzyme-linked immunospot: an alternative method for the detection of interferon gamma in Johne's disease. [J Vet Diagn Invest. 21: 187-96.](#)
6. Panadero, R. *et al.* (2009) Immunomodulatory effect of *Hypoderma lineatum* antigens: in vitro effect on bovine lymphocyte proliferation and cytokine production. [Parasite Immunol. 31: 72-7.](#)
7. Elhmouzi-Younes, J. *et al.* (2009) Bovine neonate natural killer cells are fully functional and highly responsive to interleukin-15 and to NKp46 receptor stimulation. [Vet Res. 2009 40: 54.](#)
8. Ferret-Bernard, S. *et al.* (2010) Cellular and molecular mechanisms underlying the strong neonatal IL-12 response of lamb mesenteric lymph node cells to R-848. [PLoS One. 5: e13705.](#)
9. Contreras, V. *et al.* (2010) Existence of CD8 α -like dendritic cells with a conserved functional specialization and a common molecular signature in distant mammalian species. [J Immunol. 185: 3313-25.](#)
10. Elhmouzi-Younes, J. *et al.* (2010) Ovine CD16+/CD14- blood lymphocytes present all the major characteristics of natural killer cells. [Vet Res. 41 \(1\): 4.](#)
11. McLaughlin, K. *et al.* (2010) Hsp70 enhances presentation of FMDV antigen to bovine CD4+ T cells *in vitro*. [Vet Res. 41: 36.](#)
12. Ferret-Bernard, S. *et al.* (2011) Mesenteric lymph node cells from neonates present a prominent IL-12 response to CpG oligodeoxynucleotide via an IL-15 feedback loop of amplification. [Vet Res. 42:19.](#)
13. Sow, F.B. *et al.* (2011) Respiratory syncytial virus is associated with an inflammatory response in lungs and architectural remodeling of lung-draining lymph nodes of newborn lambs. [Am J Physiol Lung Cell Mol Physiol. 300 \(1\): L12-24.](#)
14. Skyberg, J.A. *et al.* (2011) Murine and bovine $\gamma\delta$ T cells enhance innate immunity against *Brucella abortus* infections. [PLoS One. 6:e21978.](#)
15. Appana, G. *et al.* (2013) Antemortem and postmortem examinations of the cattle calf naturally infected with *Mycobacterium avium* subsp. *paratuberculosis*. [Eur J Microbiol and Immunol. 3: 241-51.](#)
16. Verhelst, D. *et al.* (2014) Parasite distribution and associated immune response during the acute phase of *Toxoplasma gondii* infection in sheep. [BMC Vet Res. 10: 293.](#)
17. Köhler H *et al.* (2015) Characterization of a caprine model for the subclinical initial phase of *Mycobacterium avium* subsp. *paratuberculosis* infection. [BMC Vet Res. 11 \(1\): 74.](#)
18. Bergmann, A. *et al.* (2015) *In Vivo* Volatile Organic Compound Signatures of *Mycobacterium avium* subsp. *paratuberculosis*. [PLoS One. 10 \(4\): e0123980.](#)
19. Cassady-Cain, R.L. *et al.* (2017) Inhibition of Antigen-Specific and Nonspecific Stimulation of Bovine T and B Cells by Lymphostatin from Attaching and Effacing *Escherichia coli*. [Infect Immun. 85\(2\):e00845-16.](#)
20. Rodrigues, V. *et al.* (2017) Development of a bead-based multiplexed assay for simultaneous quantification of five bovine cytokines by flow cytometry. [Cytometry A. 91 \(9\): 901-7.](#)
21. Villa-Mancera, A. *et al.* (2021) Phage display-based vaccine with cathepsin L and excretory-secretory products mimotopes of *Fasciola hepatica*. induces protective cellular and humoral immune responses in sheep. [Vet Parasitol. 289: 109340.](#)
22. Risalde, M.Á. *et al.* (2017) Development and evaluation of an interferon gamma assay

for the diagnosis of tuberculosis in red deer experimentally infected with *Mycobacterium bovis*. [BMC Vet Res. 13 \(1\): 341.](#)

23. Ciliberti, M.G. *et al.* (2022) Green extraction of bioactive compounds from wine lees and their bio-responses on immune modulation using in vitro sheep model. [J Dairy Sci. 105 \(5\): 4335-53.](#)

24. Santillo, A. *et al.* (2022) Feeding tannins to dairy cows in different seasons improves the oxidative status of blood plasma and the antioxidant capacity of cheese. [J Dairy Sci. 105 \(11\): 8609-20.](#)

25. Bouroutzika, E. *et al.* (2023) Melatonin Administration to Pregnant Ewes for Coccidiosis Control in Their Offspring. [Animals \(Basel\). 13 \(14\): 2381.](#)

26. Tulu, B. *et al.* (2020) Cellular and Cytokine Responses in the Granulomas of Asymptomatic Cattle Naturally Infected with *Mycobacterium bovis* in Ethiopia. [Infect Immun. 88 \(12\): e00507-20.](#)

27. Edmans, M.D. *et al.* (2024) MAIT cell-MR1 reactivity is highly conserved across multiple divergent species. [J Biol Chem. 107338. 3 May \[Epub ahead of print\].](#)

28. Burucúa, M.M. *et al.* (2024) Immunoregulatory and antiviral effect mediated by TLR7 and BMAP28 interaction in bovine alphaherpesvirus-infected respiratory primary cultures [Veterinary Microbiology. : 110342.](#)

Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.
----------------	---

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee	12 months from date of despatch
------------------	---------------------------------

Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA2112
--------------------------------------	--

Regulatory	For research purposes only
-------------------	----------------------------

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...)	RPE
Goat Anti Mouse IgG IgA IgM (STAR87...)	HRP
Goat Anti Mouse IgG (STAR70...)	FITC
Rabbit Anti Mouse IgG (STAR13...)	HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC
Goat Anti Mouse IgG (STAR77...)	HRP
Goat Anti Mouse IgG (STAR76...)	RPE
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight®488 , DyLight®550 , DyLight®650 , DyLight®680 , DyLight®800 , FITC , HRP

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

Product inquiries: www.bio-rad-antibodies.com/technical-support

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M437822:250319'

Printed on 03 Jul 2025

© 2025 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)