

## Datasheet: MCA2120

**BATCH NUMBER 158956**

<b>Description:</b>	MOUSE ANTI GRANZYME B
<b>Specificity:</b>	GRANZYME B
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	GB11
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	0.1 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry (1)	▪			1/10 - 1/100
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA	▪			2ug/ml - 5ug/ml
Immunoprecipitation	▪			
Western Blotting			▪	
Immunofluorescence	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

**(1) Membrane permeabilisation is required for this application. Bio-Rad recommends the use of Leucoperm™ (Product Code [BUF09](#)) for this purpose.**

<b>Target Species</b>	Human
<b>Species Cross Reactivity</b>	Reacts with: Chimpanzee, Monkey, Rhesus Monkey <b>N.B.</b> Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.
<b>Product Form</b>	Purified IgG - liquid

<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
<b>Buffer Solution</b>	Tris buffered saline
<b>Preservative Stabilisers</b>	0.09% Sodium azide
<b>Approx. Protein Concentrations</b>	IgG concentration 1.0 mg/ml
<b>Immunogen</b>	Purified human Granzyme B.
<b>External Database Links</b>	<p><b>UniProt:</b>  <a href="#">P10144</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">3002</a>    GZMB    <a href="#">Related reagents</a></p>
<b>Synonyms</b>	CGL1, CSPB, CTLA1, GRB
<b>RRID</b>	AB_2114582
<b>Fusion Partners</b>	Spleen cells from immunised Balb/c mice were fused with cells of the mouse SP2/0 myeloma cell line.
<b>Specificity</b>	<p><b>Mouse anti Granzyme B antibody, clone GB11</b> recognizes the serine protease Granzyme B, important in the induction of apoptosis in target cells by cytolytic lymphocytes (CTLs).</p> <p>Granzyme B plays a key role in the induction of apoptosis by CTLs. After delivery to the target cell, Granzyme B activates the cascade of caspases that finally results in cell death.</p> <p>Mouse anti Granzyme B antibody, clone GB11 is suitable for the detection of Granzyme B expressing cells by flow cytometry. In normal peripheral blood approximately 20% of CD8+ve T cells have been found to express Granzyme B.</p>
<b>Flow Cytometry</b>	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells or 100ul whole blood.
<b>ELISA</b>	This antibody has been reported to function as a capture reagent in sandwich ELISA assays for soluble Granzyme B in conjunction with biotin conjugated <a href="#">GB10</a> as detection reagent.
<b>References</b>	<ol style="list-style-type: none"> <li>Spaeny-Dekking, E.H. <i>et al.</i> (1998) Extracellular granzymes A and B in humans: detection of native species during CTL responses <i>in vitro</i> and <i>in vivo</i>. <a href="#">J Immunol. 160 (7): 3610-6.</a></li> <li>Tschopp, C.M. (2006) Granzyme B, a novel mediator of allergic inflammation: its induction and release in blood basophils and human asthma. <a href="#">Blood.108:2290-9.</a></li> </ol>

3. Wever, D. C. *et al.* (1998) The CD8+ Granzyme B+ T cell subset in peripheral blood from healthy individuals contains activated and apoptosis-prone cells. [Immunol. 93: 383](#)
4. Hsi, B.L. & Yeh, C.J. (1986) Monoclonal antibodies to human amnion. [J Reprod Immunol. 9 \(1\): 11-21.](#)
5. Mahrus, S. & Craik, C.S. (2005) Selective chemical functional probes of granzymes A and B reveal granzyme B is a major effector of natural killer cell-mediated lysis of target cells. [Chem Biol. 12: 567-77.](#)
6. Davis, C.C. *et al.* (2010) Interleukin-7 permits Th1/Tc1 maturation and promotes ex vivo expansion of cord blood T cells: a critical step toward adoptive immunotherapy after cord blood transplantation. [Cancer Res. 70: 5249-58.](#)
7. Hallermalm, K. *et al.* (2008) Modulation of the tumor cell phenotype by IFN-gamma results in resistance of uveal melanoma cells to granule-mediated lysis by cytotoxic lymphocytes. [J Immunol. 180: 3766-74.](#)
8. Hufner, K. *et al.* (2009) Fewer latent herpes simplex virus type 1 and cytotoxic T cells occur in the ophthalmic division than in the maxillary and mandibular divisions of the human trigeminal ganglion and nerve. [J Virol. 83: 3696-703.](#)
9. Kumar, D. *et al.* (2009) JNK MAPK pathway regulates constitutive transcription of CCL5 by human NK cells through SP1. [J Immunol. 182: 1011-20.](#)
10. Clayton, A. *et al.* (2008) Human tumor-derived exosomes down-modulate NKG2D expression. [J Immunol. 180: 7249-58.](#)
11. Schleypen, J.S. *et al.* (2006) Cytotoxic markers and frequency predict functional capacity of natural killer cells infiltrating renal cell carcinoma. [Clin Cancer Res. 12: 718-25.](#)
12. Hodge, S. *et al.* (2015) Increased CD8 T-cell granzyme B in COPD is suppressed by treatment with low-dose azithromycin. [Respirology. 20 \(1\): 95-100.](#)
13. Kumagai-Takei, N. *et al.* (2016) The Suppressed Induction of Human Mature Cytotoxic T Lymphocytes Caused by Asbestos Is Not due to Interleukin-2 Insufficiency. [J Immunol Res. 2016: 7484872.](#)
14. Kumagai-Takei, N. *et al.* (2021) Effect of IL-15 addition on asbestos-induced suppression of human cytotoxic T lymphocyte induction. [Environ Health Prev Med. 26 \(1\): 50.](#)
15. Jimenez, O. *et al.* (2019) M1-like macrophage polarization prevails in young children with classic Hodgkin Lymphoma from Argentina. [Sci Rep. 9 \(1\): 12687.](#)

<b>Storage</b>	<p>This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.</p> <p>Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.</p>
<b>Guarantee</b>	12 months from date of despatch
<b>Health And Safety Information</b>	<p>Material Safety Datasheet documentation #10057 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA2120">https://www.bio-rad-antibodies.com/SDS/MCA2120</a></p> <p>10057</p>
<b>Regulatory</b>	For research purposes only

## Related Products

### Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...)	<a href="#">RPE</a>
Goat Anti Mouse IgG IgA IgM (STAR87...)	<a href="#">HRP</a>
Goat Anti Mouse IgG (STAR76...)	<a href="#">RPE</a>
Goat Anti Mouse IgG (STAR70...)	<a href="#">FITC</a>
Rabbit Anti Mouse IgG (STAR13...)	<a href="#">HRP</a>
Goat Anti Mouse IgG (Fc) (STAR120...)	<a href="#">FITC</a> , <a href="#">HRP</a>
Rabbit Anti Mouse IgG (STAR9...)	<a href="#">FITC</a>
Goat Anti Mouse IgG (STAR77...)	<a href="#">HRP</a>
Goat Anti Mouse IgG (H/L) (STAR117...)	<a href="#">Alk. Phos.</a> , <a href="#">DyLight®488</a> , <a href="#">DyLight®550</a> , <a href="#">DyLight®650</a> , <a href="#">DyLight®680</a> , <a href="#">DyLight®800</a> , <a href="#">FITC</a> , <a href="#">HRP</a>

### Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://bio-rad-antibodies.com/datasheets)  
'M384263:210513'

Printed on 19 Mar 2025