

Datasheet: MCA2126APC

| | |
|----------------------|---------------------------|
| Description: | MOUSE ANTI HUMAN CD58:APC |
| Specificity: | CD58 |
| Other names: | LFA-3 |
| Format: | APC |
| Product Type: | Monoclonal Antibody |
| Clone: | MEM-63 |
| Isotype: | IgG1 |
| Quantity: | 100 TESTS |

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

| | Yes | No | Not Determined | Suggested Dilution |
|----------------|-----|----|----------------|--------------------|
| Flow Cytometry | ■ | | | Neat - 1/10 |

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

| | | | |
|---------------------------------|----------------------------------------------------------------------------------------|----------------------------|--------------------------|
| Target Species | Human | | |
| Product Form | Purified IgG conjugated to allophycocyanin (APC) - liquid | | |
| Max Ex/Em | Fluorophore | Excitation Max (nm) | Emission Max (nm) |
| | APC | 650 | 661 |
| Preparation | Purified IgG prepared by size-exclusion chromatography from tissue culture supernatant | | |
| Buffer Solution | TRIS buffered saline | | |
| Preservative Stabilisers | 0.09% sodium azide (NaN ₃) | | |
| External Database Links | UniProt: P19256 Related reagents | | |

Entrez Gene:[965](#) CD58 [Related reagents](#)

| | |
|-------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Synonyms | LFA3 |
| RRID | AB_2291397 |
| Specificity | Mouse anti Human CD58 antibody, clone MEM-63 recognizes human CD58, also known as LFA-3. CD58 is a membrane glycoprotein of ~55-70 kDa. It occurs in two forms, one transmembrane with a cytoplasmic domain, the other form anchored in the membrane via a glycosylphosphatidylinositol tail. The complete amino acid sequence of both forms has been deduced from cDNA. CD58 is a heavily N-glycosylated cell adhesion molecule which plays a critical role in facilitation of antigen specific recognition through interaction with CD2 on T lymphocytes (Makgoba <i>et al.</i> 1989). CD58 has a wide tissue distribution, being present on erythrocytes, platelets, monocytes, a subset of lymphocytes, bone marrow cells, epithelium and endothelial cells. There are approximately 5,000 CD58 molecules on each erythrocyte. There is reduced expression of CD58 on haemopoietic cells in individuals with paroxysmal nocturnal haemoglobinuria. |
| Flow Cytometry | Use 10µl of the suggested working dilution to label 10 ⁶ cells in 100µl |
| References | <ol style="list-style-type: none">1. Makgoba, M.W. <i>et al.</i> (1989) The CD2-LFA-3 and LFA-1-ICAM pathways: relevance to T-cell recognition. Immunol Today. 10 (12): 417-22.2. Shaw, S., Johnson, J.P., (1989) In Leucocyte Typing IV: White Cell Differentiation Antigens. Edited by Knapp, W., Dorken, B., Gilks, W.R., Rieber, E.P., Schmidt, R.E., Stein, H. and von dem Borne, A.E.G.Kr. Oxford University Press. pp 714-716.3. Grundy, J.E. <i>et al.</i> (1993) Increased adherence of CD2 peripheral blood lymphocytes to cytomegalovirus-infected fibroblasts is blocked by anti-LFA-3 antibody. Immunology. 78 (3): 413-20.4. Megyola, C. <i>et al.</i> (2011) Identification of a sub-population of B cells that proliferates after infection with Epstein-Barr virus. Virol J. 8: 84. |
| Storage | Store at +4°C. DO NOT FREEZE. This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use. |
| Guarantee | Guaranteed for 12 months from the date of despatch or until the date of expiry, whichever comes first. Please see label for expiry date. |
| Health And Safety Information | Material Safety Datasheet documentation #10057 available at: https://www.bio-rad-antibodies.com/SDS/MCA2126APC 10057 |
| Regulatory | For research purposes only |

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:APC \(MCA928APC\)](#)

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

**North & South
America**

Tel: +1 800 265 7376

Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M412970:221117'

Printed on 06 Oct 2023