

Datasheet: MCA2126F

BATCH NUMBER 163814

Description:	MOUSE ANTI HUMAN CD58:FITC
Specificity:	CD58
Other names:	LFA-3
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	MEM-63
Isotype:	IgG1
Quantity:	100 TESTS/2ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human		
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant		
Buffer Solution	TRIS buffered saline		
Preservative Stabilisers	0.09% Sodium Azide		
External Database	UniProt:		

Links

[P19256](#) [Related reagents](#)

Entrez Gene:

[965](#) CD58 [Related reagents](#)

Synonyms

LFA3

RRID

AB_323491

Specificity

Mouse anti Human CD58 antibody, clone MEM-63 recognizes human CD58, also known as LFA-3. CD58 is a membrane glycoprotein of ~55-70 kDa. It occurs in two forms, one transmembrane with a cytoplasmic domain, the other form anchored in the membrane via a glycosylphosphatidylinositol tail. The complete amino acid sequence of both forms has been deduced from cDNA. CD58 is a heavily N-glycosylated cell adhesion molecule which plays a critical role in facilitation of antigen specific recognition through interaction with CD2 on T lymphocytes ([Makgoba *et al.* 1989](#)). CD58 has a wide tissue distribution, being present on erythrocytes, platelets, monocytes, a subset of lymphocytes, bone marrow cells, epithelium and endothelial cells. There are approximately 5,000 CD58 molecules on each erythrocyte. There is reduced expression of CD58 on haemopoietic cells in individuals with paroxysmal nocturnal haemoglobinuria.

Flow Cytometry

Use 20ul of the suggested working dilution to label 10⁶ cells in 100ul.

References

1. Makgoba, M.W. *et al.* (1989) The CD2-LFA-3 and LFA-1-ICAM pathways: relevance to T-cell recognition. [Immunol Today. 10 \(12\): 417-22.](#)
2. Shaw, S., Johnson, J.P., (1989) In Leucocyte Typing IV: White Cell Differentiation Antigens. Edited by Knapp, W., Dorken, B., Gilks, W.R., Rieber, E.P., Schmidt, R.E., Stein, H. and von dem Borne, A.E.G.Kr. Oxford University Press. pp 714-716.
3. Grundy, J.E. *et al.* (1993) Increased adherence of CD2 peripheral blood lymphocytes to cytomegalovirus-infected fibroblasts is blocked by anti-LFA-3 antibody. [Immunology. 78 \(3\): 413-20.](#)
4. Megyola, C. *et al.* (2011) Identification of a sub-population of B cells that proliferates after infection with Epstein-Barr virus. [Virol J. 8: 84.](#)

Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Guarantee

Guaranteed for 12 months from the date of despatch or until the date of expiry, whichever comes first. Please see label for expiry date.

Health And Safety Information

Material Safety Datasheet documentation #10057 available at: <https://www.bio-rad-antibodies.com/SDS/MCA2126F10057>

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:FITC \(MCA928F\)](#)

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M410512:221028'

Printed on 18 Jan 2024