

## Datasheet: MCA2127

**BATCH NUMBER 1608**

<b>Description:</b>	MOUSE ANTI HUMAN CD25
<b>Specificity:</b>	CD25
<b>Other names:</b>	IL-2R ALPHA CHAIN
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	MEM-181
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	0.2 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/50 - 1/200
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Human
<b>Product Form</b>	Purified IgG - liquid
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein A
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.09% Sodium Azide

<b>Approx. Protein Concentrations</b>	IgG concentration 1 mg/ml
<b>Immunogen</b>	Human PHA blasts; day 3 of culture.
<b>External Database Links</b>	<p><b>UniProt:</b>  <a href="#">P01589</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">3559</a>    IL2RA    <a href="#">Related reagents</a></p>
<b>RRID</b>	AB_324271
<b>Fusion Partners</b>	Spleen cells from immunised BALB/c mice were fused with cells of the mouse P3.X63 Ag8.653 myeloma cell line.
<b>Specificity</b>	<p><b>Mouse anti Human CD25, clone MEM-181</b> recognizes the ~55 kDa alpha subunit of the human IL-2 receptor, also known as p55 or TAC antigen, CD25 is a type 1 transmembrane protein with <a href="#">two Sushi domains</a>, also known as short consensus repeats (SCRs) or complement control protein (CCP) modules (<a href="#">Norman et al. 1991</a>) located within its extracellular domain.</p> <p>The IL-2 receptor exists in three forms. A high affinity form consisting of a non-covalently linked heterodimer composed of the alpha subunit (CD25) and the IL-2 receptor beta subunit also known as CD122 or p75, a medium affinity beta subunit (CD122) monomer or a low affinity alpha (CD25) subunit monomer.</p> <p>CD25 is expressed by activated T lymphocytes and activated B lymphocytes responding to antigen or mitogen stimulation. CD25 is also expressed in some thymocytes and oligodendrocytes. In disease, elevated expression of CD25 is noted in a number of chronic inflammatory conditions, tuberculoid leprosy patients demonstrate markedly elevated levels of circulating CD25<sup>high</sup> FoxP3<sup>+</sup> regulatory T cells (T-regs) (<a href="#">Attia et al. 2010</a>).</p> <p>Elevated levels of CD25 antigen expression are often seen in cases of <a href="#">non-Hodgkin 's lymphoma</a> and diffuse large B cell lymphoma (<a href="#">Fujiwara et al.2013</a>).</p>
<b>Flow Cytometry</b>	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul.
<b>References</b>	<ol style="list-style-type: none"> <li>1. Prager, E. <i>et al.</i> (2001) Induction of hyporesponsiveness and impaired T lymphocyte activation by the CD31 receptor:ligand pathway in T cells. <a href="#">J Immunol. 166 (4): 2364-71.</a></li> <li>2. Thorborn, G. <i>et al.</i> (2010) Increased sensitivity of CD4<sup>+</sup> T-effector cells to CD4<sup>+</sup>CD25<sup>+</sup> Treg suppression compensates for reduced Treg number in asymptomatic HIV-1 infection. <a href="#">PLoS One. 5: e9254.</a></li> <li>3. Cutler, A.J. <i>et al.</i> (2010) Umbilical cord-derived mesenchymal stromal cells modulate monocyte function to suppress T cell proliferation. <a href="#">J Immunol. 185: 6617-23.</a></li> <li>4. Lawson, J.M. <i>et al.</i> (2008) Increased resistance to CD4<sup>+</sup>CD25<sup>hi</sup> regulatory T</li> </ol>

- cell-mediated suppression in patients with type 1 diabetes. [Clin Exp Immunol. 154: 353-9.](#)
5. Holderness, J. *et al.* (2007) Select plant tannins induce IL-2Ralpha up-regulation and augment cell division in gammadelta T cells. [J Immunol. 179: 6468-78.](#)
  6. Zhang, Y. *et al.* (2013) Accelerated *in vivo* proliferation of memory phenotype CD4+ T-cells in human HIV-1 infection irrespective of viral chemokine co-receptor tropism. [PLoS Pathog. 9 \(4\): e1003310.](#)
  7. Nocentini, G. *et al.* (2014) Expansion of regulatory GITR + CD25 Low/- CD4 + T cells in systemic lupus erythematosus patients. [Arthritis Res Ther. 16: 444.](#)
  8. Soukup, K. *et al.* (2015) The MAPK-Activated Kinase MK2 Attenuates Dendritic Cell-Mediated Th1 Differentiation and Autoimmune Encephalomyelitis. [J Immunol. 195 \(2\): 541-52.](#)
  9. Kusunoki, Y. *et al.* (2010) T-cell immunosenescence and inflammatory response in atomic bomb survivors. [Radiat Res. 174 \(6\): 870-6.](#)
  10. Bughani, U. *et al.* (2017) T cell activation and differentiation is modulated by a CD6 domain 1 antibody Itolizumab. [PLoS One. 12 \(7\): e0180088.](#)
  11. Knutson, K.L. *et al.* (2015) Regulatory T cells, inherited variation, and clinical outcome in epithelial ovarian cancer. [Cancer Immunol Immunother. 64 \(12\): 1495-504.](#)
  12. Boland, J.W. *et al.* (2014) A preliminary evaluation of the effects of opioids on innate and adaptive human *in vitro* immune function. [BMJ Support Palliat Care. 4 \(4\): 357-67.](#)
  13. Luger, R. *et al.* (2013) Toll-like receptor 4 engagement drives differentiation of human and murine dendritic cells from a pro- into an anti-inflammatory mode. [PLoS One. 8 \(2\): e54879.](#)

---

**Storage**

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

---

**Guarantee**

12 months from date of despatch

---

**Health And Safety Information**

Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/MCA2127>  
10040

---

**Regulatory**

For research purposes only

---

## Related Products

### Recommended Secondary Antibodies

- |   |  |
|---|--|
| Goat Anti Mouse IgG (STAR77...)         | <a href="#">HRP</a>                              |
| Rabbit Anti Mouse IgG (STAR12...)       | <a href="#">RPE</a>                              |
| Goat Anti Mouse IgG IgA IgM (STAR87...) | <a href="#">Alk. Phos.</a> , <a href="#">HRP</a> |
| Goat Anti Mouse IgG (STAR76...)         | <a href="#">RPE</a>                              |
| Rabbit Anti Mouse IgG (STAR13...)       | <a href="#">HRP</a>                              |

Goat Anti Mouse IgG (STAR70...)

[FITC](#)

Goat Anti Mouse IgG (H/L) (STAR117...)

[Alk. Phos.](#), [DyLight®488](#), [DyLight®550](#),  
[DyLight®650](#), [DyLight®680](#), [DyLight®800](#),  
[FITC](#), [HRP](#)

Rabbit Anti Mouse IgG (STAR9...)

[FITC](#)

Goat Anti Mouse IgG (Fc) (STAR120...)

[FITC](#), [HRP](#)

## Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://bio-rad-antibodies.com/datasheets)

'M366230:200529'

Printed on 12 Sep 2023

---

© 2023 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)