

Datasheet: MCA2127SBV515

BATCH NUMBER 100004706

Description:	MOUSE ANTI HUMAN CD25:StarBright Violet 515
Specificity:	CD25
Other names:	IL-2R ALPHA CHAIN
Format:	StarBright Violet 515
Product Type:	Monoclonal Antibody
Clone:	MEM-181
Isotype:	IgG1
Quantity:	100 TESTS/0.5ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human		
Product Form	Purified IgG conjugated to StarBright Violet 515 - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	StarBright Violet 515	401	516
Preparation	Purified IgG prepared by affinity chromatography on Protein A		
Buffer Solution	Phosphate buffered saline		
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃) 1% Bovine Serum Albumin 0.1% Pluronic F68 0.1% PEG 3350		

Immunogen Human PHA blasts; day 3 of culture.

External Database Links

UniProt:

[P01589](#) [Related reagents](#)

Entrez Gene:

[3559](#) IL2RA [Related reagents](#)

Fusion Partners

Spleen cells from immunised BALB/c mice were fused with cells of the mouse P3.X63 Ag8.653 myeloma cell line.

Specificity

Mouse anti Human CD25, clone MEM-181 recognizes the ~55 kDa alpha subunit of the human IL-2 receptor, also known as p55 or TAC antigen, CD25 is a type 1 transmembrane protein with [two Sushi domains](#), also known as short consensus repeats (SCRs) or complement control protein (CCP) modules ([Norman et al. 1991](#)) located within its extracellular domain.

The IL-2 receptor exists in three forms. A high affinity form consisting of a non-covalently linked heterodimer composed of the alpha subunit (CD25) and the IL-2 receptor beta subunit also known as CD122 or p75, a medium affinity beta subunit (CD122) monomer or a low affinity alpha (CD25) subunit monomer.

CD25 is expressed by activated T lymphocytes and activated B lymphocytes responding to antigen or mitogen stimulation. CD25 is also expressed in some thymocytes and oligodendrocytes. In disease, elevated expression of CD25 is noted in a number of chronic inflammatory conditions, tuberculoid leprosy patients demonstrate markedly elevated levels of circulating CD25^{high} FoxP3⁺ regulatory T cells (T-regs) ([Attia et al. 2010](#)).

Elevated levels of CD25 antigen expression are often seen in cases of [non-Hodgkin's lymphoma](#) and diffuse large B cell lymphoma ([Fujiwara et al. 2013](#)).

Flow Cytometry

Use 5ul of the suggested working dilution to label 10⁶ cells in 100ul. Best practices suggest a 5 minutes centrifugation at 6,000g prior to sample application.

References

1. Prager, E. *et al.* (2001) Induction of hyporesponsiveness and impaired T lymphocyte activation by the CD31 receptor:ligand pathway in T cells. [J Immunol. 166 \(4\): 2364-71.](#)
2. Thorborn, G. *et al.* (2010) Increased sensitivity of CD4⁺ T-effector cells to CD4⁺CD25⁺ Treg suppression compensates for reduced Treg number in asymptomatic HIV-1 infection. [PLoS One. 5: e9254.](#)
3. Cutler, A.J. *et al.* (2010) Umbilical cord-derived mesenchymal stromal cells modulate monocyte function to suppress T cell proliferation. [J Immunol. 185: 6617-23.](#)
4. Lawson, J.M. *et al.* (2008) Increased resistance to CD4⁺CD25^{hi} regulatory T cell-mediated suppression in patients with type 1 diabetes. [Clin Exp Immunol. 154: 353-9.](#)
5. Holderness, J. *et al.* (2007) Select plant tannins induce IL-2R α up-regulation and augment cell division in gammadelta T cells. [J Immunol. 179: 6468-78.](#)
6. Zhang, Y. *et al.* (2013) Accelerated *in vivo* proliferation of memory phenotype CD4⁺

T-cells in human HIV-1 infection irrespective of viral chemokine co-receptor tropism. [PLoS Pathog. 9 \(4\): e1003310.](#)

7. Nocentini, G. *et al.* (2014) Expansion of regulatory GITR + CD25 Low/- CD4 + T cells in systemic lupus erythematosus patients. [Arthritis Res Ther. 16: 444.](#)

8. Soukup, K. *et al.* (2015) The MAPK-Activated Kinase MK2 Attenuates Dendritic Cell-Mediated Th1 Differentiation and Autoimmune Encephalomyelitis. [J Immunol. 195 \(2\): 541-52.](#)

9. Kusunoki, Y. *et al.* (2010) T-cell immunosenescence and inflammatory response in atomic bomb survivors. [Radiat Res. 174 \(6\): 870-6.](#)

10. Bughani, U. *et al.* (2017) T cell activation and differentiation is modulated by a CD6 domain 1 antibody Itolizumab. [PLoS One. 12 \(7\): e0180088.](#)

11. Knutson, K.L. *et al.* (2015) Regulatory T cells, inherited variation, and clinical outcome in epithelial ovarian cancer. [Cancer Immunol Immunother. 64 \(12\): 1495-504.](#)

12. Boland, J.W. *et al.* (2014) A preliminary evaluation of the effects of opioids on innate and adaptive human *in vitro* immune function. [BMJ Support Palliat Care. 4 \(4\): 357-67.](#)

13. Luger, R. *et al.* (2013) Toll-like receptor 4 engagement drives differentiation of human and murine dendritic cells from a pro- into an anti-inflammatory mode. [PLoS One. 8 \(2\): e54879.](#)

Storage	Store at +4°C. DO NOT FREEZE. This product should be stored undiluted.
Guarantee	12 months from date of despatch
Acknowledgements	This product is covered by U.S. Patent No. 10,150,841 and related U.S. and foreign counterparts
Health And Safety Information	Material Safety Datasheet documentation #20438 available at: https://www.bio-rad-antibodies.com/SDS/MCA2127SBV515 20438
Regulatory	For research purposes only

Related Products

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)
'M373243:200901'

Printed on 08 Mar 2024