

## Datasheet: MCA2141FT

<b>Description:</b>	MOUSE ANTI HUMAN CD146:FITC
<b>Specificity:</b>	CD146
<b>Other names:</b>	MUC18
<b>Format:</b>	FITC
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	OJ79c
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	25 µg

### Product Details

#### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat - 1/10

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

#### Target Species

Human

#### Species Cross Reactivity

Reacts with: Pig, Dog

**N.B.** Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

#### Product Form

Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid

#### Max Ex/Em

Fluorophore	Excitation Max (nm)	Emission Max (nm)
FITC	490	525

#### Preparation

Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant

#### Buffer Solution

Phosphate buffered saline

<b>Preservative Stabilisers</b>	0.09% sodium azide (NaN <sub>3</sub> ) 1% bovine serum albumin
<b>Approx. Protein Concentrations</b>	IgG concentration 0.1 mg/ml
<b>Immunogen</b>	Recombinant human MUC18 (D1-D5) Fc protein.
<b>External Database Links</b>	<p><b>UniProt:</b>  <a href="#">P43121</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">4162</a>    MCAM    <a href="#">Related reagents</a></p>
<b>Synonyms</b>	MUC18
<b>RRID</b>	AB_2143381
<b>Fusion Partners</b>	Spleen cells from immunized mice were fused with cells of the mouse Sp2/0 Ag.14 myeloma cell line.
<b>Specificity</b>	<p><b>Mouse anti Human CD146 antibody, clone OJ79c</b> recognizes human Cell surface glycoprotein MUC18, also known as CD146, Cell surface glycoprotein P1H12, Melanoma cell adhesion molecule (MCAM) or S-endo 1 endothelial-associated antigen. CD146 is a 646 amino acid single pass type 1 transmembrane glycoprotein with a calculated molecular mass of ~72 kDa. However due to extensive N-linked glycosylation CD146 migrates in polyacrylamide gels with an apparent molecular mass of ~118 kDa. CD146 is a member of the immunoglobulin superfamily bearing 2 <a href="#">V-type Ig-like</a> and 3 <a href="#">C-type Ig-like domains</a>. CD146 is expressed by all endothelial cells and by melanoma cells and appears to act as an adhesion molecule (<a href="#">UniProt: P43121</a>). Expression in melanoma may be linked to tumor progression (<a href="#">Lehmann et al. 1989</a>).</p> <p>Mouse anti Human CD146 antibody, clone OJ79c is highly expressed on pericytes and has been utilized for the identification of perivascular mesenchymal precursor cells from cardiac muscle using flow cytometry (<a href="#">Chen et al. 2014</a>).</p>
<b>Flow Cytometry</b>	Use 10µl of the suggested working dilution to label 10 <sup>6</sup> cells in 100µl
<b>References</b>	<ol style="list-style-type: none"> <li>Paul, G. <i>et al.</i> (2012) The adult human brain harbors multipotent perivascular mesenchymal stem cells. <a href="#">PLoS One. 7: e35577.</a></li> <li>Crisan, M. <i>et al.</i> (2008) A perivascular origin for mesenchymal stem cells in multiple human organs. <a href="#">Cell Stem Cell. 3: 301-13.</a></li> <li>Iohara, K. <i>et al.</i> (2008) A novel stem cell source for vasculogenesis in ischemia: subfraction of side population cells from dental pulp. <a href="#">Stem Cells. 26 (9): 2408-18.</a></li> <li>Park, T.S. <i>et al.</i> (2010) Placental Perivascular Cells for Human Muscle Regeneration. <a href="#">Stem Cells Dev. 20: 451-63.</a></li> <li>Smith, K. <i>et al.</i> (2011) Mono- and tri-cationic porphyrin-monoclonal antibody conjugates:</li> </ol>

- photodynamic activity and mechanism of action. [Immunology. 132: 256-65.](#)
6. James, A.W. *et al.* (2012) Perivascular stem cells: a prospectively purified mesenchymal stem cell population for bone tissue engineering. [Stem Cells Transl Med. 1 \(6\): 510-9.](#)
7. Wassmer, S.C. *et al.* (2011) Vascular endothelial cells cultured from patients with cerebral or uncomplicated malaria exhibit differential reactivity to TNF. [Cell Microbiol. 13: 198-209.](#)
8. Lee, J.H. *et al.* (2012) Generation of osteogenic construct using periosteal-derived osteoblasts and polydioxanone/pluronic F127 scaffold with periosteal-derived CD146 positive endothelial-like cells. [J Biomed Mater Res A.101: 942-53.](#)
9. Boneberg, E.M. *et al.* (2009) Soluble CD146 is generated by ectodomain shedding of membrane CD146 in a calcium-induced, matrix metalloprotease-dependent process. [Microvasc Res. 78: 325-31.](#)
10. Nielsen, C.T. *et al.* (2011) Distinct features of circulating microparticles and their relationship to clinical manifestations in systemic lupus erythematosus. [Arthritis Rheum. 63: 3067-77.](#)
11. Iversen, L.V. *et al.* (2013) A heparin-based method for flow cytometric analysis of microparticles directly from platelet-poor plasma in calcium containing buffer. [J Immunol Methods. 388 \(1-2\): 49-59.](#)
12. Meireles, A.L. *et al.* (2011) Increased levels of circulating endothelial progenitor cells in human T-cell lymphotropic virus type I carriers. [Arch Med Res. 42: 34-7.](#)
13. Iversen, L.V. *et al.* (2013) Circulating microparticles and plasma levels of soluble E- and P-selectins in patients with systemic sclerosis. [Scand J Rheumatol. 42 \(6\): 473-82.](#)
14. Murakami, M. *et al.* (2013) The use of granulocyte-colony stimulating factor induced mobilization for isolation of dental pulp stem cells with high regenerative potential. [Biomaterials. pii: S0142-9612\(13\)00942-3.](#)
15. Ruetze, M. *et al.* (2013) A novel niche for skin derived precursors in non-follicular skin. [J Dermatol Sci. 69: 132-9.](#)
16. Dokić, J. *et al.* (2013) Mesenchymal stem cells from periapical lesions modulate differentiation and functional properties of monocyte-derived dendritic cells. [Eur J Immunol. 43: 1862-72.](#)
17. Chen, W.C. *et al.* (2015) Human myocardial pericytes: multipotent mesodermal precursors exhibiting cardiac specificity. [Stem Cells. 33 \(2\): 557-73.](#)
18. Iversen, L.V. *et al.* (2015) Cross-sectional study of soluble selectins, fractions of circulating microparticles and their relationship to lung and skin involvement in systemic sclerosis. [BMC Musculoskelet Disord. 16: 191.](#)
19. Tawonsawatruk T *et al.* (2016) Adipose derived pericytes rescue fractures from a failure of healing - non-union. [Sci Rep. 6: 22779.](#)
20. Boissier, R. *et al.* (2016) Histologic and urodynamic effects of autologous stromal vascular fraction extracted from fat tissue with minimal *ex vivo* manipulation on a porcine model of intrinsic sphincter deficiency [J Urology. Jun 2 \[Epub ahead of print\]](#)
21. Esteves, C.L. *et al.* (2017) Equine Mesenchymal Stromal Cells Retain a Pericyte-Like Phenotype. [Stem Cells Dev. 26 \(13\): 964-72.](#)
22. Stefanska, A. *et al.* (2016) Human kidney pericytes produce renin. [Kidney Int. 90 \(6\): 1251-61.](#)
23. Lapin, M. *et al.* (2016) MINDEC-An Enhanced Negative Depletion Strategy for Circulating Tumour Cell Enrichment. [Sci Rep. 6: 28929.](#)

24. Muerza-Cascante, M.L. *et al.* (2016) Endosteal-like extracellular matrix expression on melt electrospun written scaffolds. [Acta Biomater. pii: S1742-7061\(16\)30706-1.](#)
25. Eliasberg, C.D. *et al.* (2017) Perivascular Stem Cells Diminish Muscle Atrophy Following Massive Rotator Cuff Tears in a Small Animal Model. [J Bone Joint Surg Am. 99 \(4\): 331-41.](#)
26. James, A.W. *et al.* (2017) Isolation and characterization of canine perivascular stem/stromal cells for bone tissue engineering. [PLoS One. 12 \(5\): e0177308.](#)
27. Shen, J. *et al.* (2018) Effects of WNT3A and WNT16 on the Osteogenic and Adipogenic Differentiation of Perivascular Stem/Stromal Cells. [Tissue Eng Part A. 24 \(1-2\): 68-80.](#)
28. Lapin, M. *et al.* (2017) Single-cell mRNA profiling reveals transcriptional heterogeneity among pancreatic circulating tumour cells. [BMC Cancer. 17 \(1\): 390.](#)
29. Esteves, C.L. *et al.* (2017) Isolation and characterization of equine native MSC populations. [Stem Cell Res Ther. 8 \(1\): 80.](#)
30. Gaceb, A. *et al.* (2017) Pericytes secrete pro-regenerative molecules in response to platelet-derived growth factor-BB. [J Cereb Blood Flow Metab. : 271678X17719645.](#)
31. Noda, S. *et al.* (2019) Effect of cell culture density on dental pulp-derived mesenchymal stem cells with reference to osteogenic differentiation. [Sci Rep. 9 \(1\): 5430.](#)
32. Gaceb, A. *et al.* (2020) An *In Vitro*. Partial Lesion Model of Differentiated Human Mesencephalic Neurons: Effect of Pericyte Secretome on Phenotypic Markers. [J Mol Neurosci. 70 \(11\): 1914-25.](#)
33. Stefanska, A. *et al.* (2021) Role of Pericytes in the Development of the Renin/Angiotensin System: Induction of Functional Renin in Cultures of Pericytes. [Methods Mol Biol. 2235: 169-180.](#)
34. Orikasa, S. *et al.* (2022) Hypoxia-inducible factor 1 $\alpha$  induces osteo/odontoblast differentiation of human dental pulp stem cells via Wnt/ $\beta$ -catenin transcriptional cofactor BCL9. [Sci Rep. 12 \(1\): 682.](#)
35. Menon, R. *et al.* (2023) Human Induced Pluripotent Stem Cell-Derived Pericytes as Scalable and Editable Source to Study Direct Lineage Reprogramming Into Induced Neurons. [Cell Reprogram. 25 \(5\): 212-23.](#)
36. Molinos, M. *et al.* (2023) Alterations of bovine nucleus pulposus cells with aging. [Aging Cell. 22 \(8\): e13873.](#)
37. An, H.J. *et al.* (2021) Pro-Angiogenic and Osteogenic Effects of Adipose Tissue-Derived Pericytes Synergistically Enhanced by Nel-like Protein-1. [Cells. 10 \(9\): 2244.](#)
38. Mendoza-Zambrano, E. *et al.* (2025) Role of Extracellular Vesicles in Chronic Post-Embolic Pulmonary Hypertension: Data from an Experimental Animal Model and Patients. [Biomedicines. 13 \(6\): 1499.](#)

---

**Further Reading**

1. Kuzu, I. *et al.* (1993) Expression of adhesion molecules on the endothelium of normal tissue vessels and vascular tumors. [Lab Invest. 69 \(3\): 322-8.](#)
2. Piriou-Guzylack, L. (2008) Membrane markers of the immune cells in swine: an update. [Vet Res. 39: 54.](#)

---

**Storage**

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

---

<b>Guarantee</b>	12 months from date of despatch
<b>Health And Safety Information</b>	Material Safety Datasheet documentation #10041 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA2141FT">https://www.bio-rad-antibodies.com/SDS/MCA2141FT</a>
<b>Regulatory</b>	For research purposes only

---

## Related Products

### Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:FITC \(MCA928F\)](#)

### Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

**Product inquiries:** [www.bio-rad-antibodies.com/technical-support](http://www.bio-rad-antibodies.com/technical-support)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](http://bio-rad-antibodies.com/datasheets)  
'M437815:250319'

**Printed on 28 Apr 2026**