

Datasheet: MCA2142APC

BATCH NUMBER 170607

Description:	MOUSE ANTI HUMAN CD63:APC
Specificity:	CD63
Format:	APC
Product Type:	Monoclonal Antibody
Clone:	MEM-259
Isotype:	IgG1
Quantity:	100 TESTS

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human		
Product Form	Purified IgG conjugated to allophycocyanin (APC) - lyophilized		
Reconstitution	Reconstitute with 1ml distilled water		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	APC	650	661
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% Sodium Azide (NaN ₃)		
Stabilisers	1% Bovine Serum Albumin 5% Sucrose		

Approx. Protein Concentrations	IgG concentration 0.1 mg/ml
Immunogen	HPB-ALL cell line.
External Database Links	<p>UniProt: P08962 Related reagents</p> <p>Entrez Gene: 967 CD63 Related reagents</p>
Synonyms	MLA1, TSPAN30
RRID	AB_2229161
Specificity	<p>Mouse anti Human CD63 antibody, clone MEM-259 recognizes the CD63 cell surface antigen, also known as granulophysin, lysosomal associated membrane protein 3 (LAMP-3), Melanoma-associated antigen ME491, Ocular melanoma-associated antigen or tetraspanin-30. CD63 is a 238 amino acid, tetraspanin glycoprotein of ~40-60 kDa.</p> <p>CD63 is expressed by monocytes, macrophages, endothelial cells where it is a component of Weibel-Palade bodies (Vischer et al. 1993) and by a variety of cell lines. It is located intracellularly in lysosomal granules of platelets, being translocated to the surface upon activation. It may therefore be useful as a marker of platelet activation (Hamamoto et al. 1994). CD63 expression is also associated with melanoma development (Radford et al. 1997) and expression is rapidly down-regulated by progesterone in endometrial tissues (Okada et al. 1999).</p>
Flow Cytometry	Use 10µl of the suggested working dilution to label 10 ⁶ cells in 100µl or 100µl whole blood
References	<ol style="list-style-type: none"> Casey, T.M. <i>et al.</i> (2007) Organelle proteomics: identification of the exocytic machinery associated with the natural killer cell secretory lysosome. Mol Cell Proteomics. 6 (5): 767-80. Durand-Panteix, S. <i>et al.</i> (2012) B7-H1, which represses EBV-immortalized B cell killing by autologous T and NK cells, is oppositely regulated by c-Myc and EBV latency III program at both mRNA and secretory lysosome levels. J Immunol. 189 (1): 181-90. Abdel-Latif, D. <i>et al.</i> (2004) Rac2 is critical for neutrophil primary granule exocytosis. Blood. 104: 832-9. Norling, L.V. <i>et al.</i> (2012) Resolvin D1 limits polymorphonuclear leukocyte recruitment to inflammatory loci: receptor-dependent actions. Arterioscler Thromb Vasc Biol. 32 (8): 1970-8. McKechnie, N.M. <i>et al.</i> (2006) Fas-ligand is stored in secretory lysosomes of ocular barrier epithelia and released with microvesicles. Exp Eye Res. 83: 304-14. Spring, F.A. <i>et al.</i> (2013) Tetraspanins CD81 and CD82 facilitate α4β1-mediated adhesion of human erythroblasts to vascular cell adhesion molecule-1. PLoS One. 8(5):e62654. Pliyev, B.K. (2008) Activated human neutrophils rapidly release the chemotactically

active D2D3 form of the urokinase-type plasminogen activator receptor (uPAR/CD87). [Mol Cell Biochem. 321: 111-22.](#)

8. Suraeva, N.M. *et al.* (2015) Changes in the Morphological and Immunological Characteristics of Mel Ibr Melanoma Cells in Response to Chicken Embryo Extract. [Bull Exp Biol Med. 159 \(4\): 520-3.](#)

9. Schäfer, T. *et al.* (2010) A granular variant of CD63 is a regulator of repeated human mast cell degranulation. [Allergy. 65 \(10\): 1242-55.](#)

10. Jakhria, T. *et al.* (2014) β 2-microglobulin amyloid fibrils are nanoparticles that disrupt lysosomal membrane protein trafficking and inhibit protein degradation by lysosomes. [J Biol Chem. 289 \(52\): 35781-94.](#)

11. Ethier, C. *et al.* (2016) Calcitriol Reduces Eosinophil Necrosis Which Leads to the Diminished Release of Cytotoxic Granules. [Int Arch Allergy Immunol. 171 \(2\): 119-29.](#)

12. Nielsen, M.H. *et al.* (2023) Increased extracellular vesicles (EVs) related to T cell-mediated inflammation and vascular function in familial hypercholesterolemia. [Atheroscler Plus. 53: 16-25.](#)

13. Ilaraza, R. *et al.* (2023) Rac2 regulates immune complex-mediated granule polarization and exocytosis in neutrophils. [J Leukoc Biol. 114 \(2\): 116-125.](#)

14. Vrancx, C. *et al.* (2021) Mechanism of Cellular Formation and *In Vivo* Seeding Effects of Hexameric β -Amyloid Assemblies. [Mol Neurobiol. 58 \(12\): 6647-69.](#)

15. Gratpain, V. *et al.* (2024) Influence of a pro-inflammatory stimulus on the miRNA and lipid content of human dental stem cell-derived extracellular vesicles and their impact on microglial activation [Heliyon. : e27025.](#)

16. Cinato, M. *et al.* (2021) Apilimod alters TGF β signaling pathway and prevents cardiac fibrotic remodeling. [Theranostics. 11 \(13\): 6491-506.](#)

Storage

Prior to reconstitution store at +4°C.

After reconstitution store at +4°C.

DO NOT FREEZE.

This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee

12 months from date of despatch

Health And Safety Information

Material Safety Datasheet documentation #20487 available at: <https://www.bio-rad-antibodies.com/SDS/MCA2142APC>

Regulatory

For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:APC \(MCA928APC\)](#)

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

Product inquiries: www.bio-rad-antibodies.com/technical-support

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M436944:250305'

Printed on 29 Apr 2026

© 2026 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)