

Datasheet: MCA2152

Description:	MOUSE ANTI HUMAN CD282
Specificity:	CD282
Other names:	TLR2
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	TLR2.3
Isotype:	IgG2a
Quantity:	0.2 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting			▪	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide
Approx. Protein Concentrations	IgG concentration 0.5 mg/ml
Immunogen	CHO cell line transfected with human TLR2 (CD282).
External Database Links	UniProt:

[O60603](#) [Related reagents](#)

Entrez Gene:

[7097](#) TLR2 [Related reagents](#)

Synonyms	TIL4
-----------------	------

Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the NS0 myeloma cell line.
------------------------	---

Specificity	<p>Mouse anti Human CD282 antibody, clone TL2.3 recognizes human TLR2, otherwise known as CD282. TLR2 is a member of the Toll-like receptor (TLR) family and is expressed primarily by peripheral blood monocytes.</p> <p>TLRs are expressed on the cell surface and the endocytic compartment and recognize pathogen-associated molecular patterns (PAMPs) that are expressed on infectious agents. They also initiate cell signalling to induce production of cytokines necessary for the innate immunity and subsequent adaptive immunity. TLR2 is reported to respond to a diverse range of bacterial cell wall components, mediating the innate immune response in co-operation with MD-2.</p> <p>Mouse anti Human CD282 antibody, clone TL2.1 is reported to block TLR2 function.</p>
--------------------	--

Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
-----------------------	---

Histology Positive Control Tissue	Tonsil.
--	---------

References	<ol style="list-style-type: none">Lien, E. <i>et al.</i> (1999) Toll-like receptor 2 functions as a pattern recognition receptor for diverse bacterial products. J Biol Chem. 274 (47): 33419-25.Flo, T.H. <i>et al.</i> (2001) Differential expression of Toll-like receptor 2 in human cells. J Leukoc Biol. 69 (3): 474-81.Sels, J.W. <i>et al.</i> (2012) Fractional flow reserve is not associated with inflammatory markers in patients with stable coronary artery disease. PLoS One. 7 (10): e46356.
-------------------	--

Storage	<p>Store at +4°C or at -20°C if preferred.</p> <p>This product should be stored undiluted.</p> <p>Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.</p>
----------------	--

Shelf Life	18 months from date of despatch.
-------------------	----------------------------------

Health And Safety Information	Material Safety Datasheet documentation #10040 available at: 10040: https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf
--------------------------------------	--

Regulatory	For research purposes only
-------------------	----------------------------

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR76...)

[RPE](#)

Goat Anti Mouse IgG IgA IgM (STAR87...) [Alk. Phos.](#), [HRP](#)
Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight@488](#), [DyLight@549](#),
[DyLight@649](#), [DyLight@680](#), [DyLight@800](#),
[FITC](#), [HRP](#)
Rabbit Anti Mouse IgG (STAR9...) [FITC](#)
Goat Anti Mouse IgG (STAR77...) [HRP](#)
Rabbit Anti Mouse IgG (STAR12...) [RPE](#)
Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)
Rabbit Anti Mouse IgG (STAR8...) [DyLight@800](#)
Goat Anti Mouse IgG (STAR70...) [FITC](#)
Human Anti Mouse IgG2a (HCA037...) [FITC](#), [HRP](#)
Rabbit Anti Mouse IgG (STAR13...) [HRP](#)

Recommended Negative Controls

[MOUSE IgG2a NEGATIVE CONTROL \(MCA929\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

'M313802:180403'

Printed on 26 May 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)