

Datasheet: MCA2152GA

BATCH NUMBER 160373

Description:	MOUSE ANTI HUMAN CD282
Specificity:	CD282
Other names:	TLR2
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	TLR2.3
Isotype:	IgG2a
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/10 - 1/50
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant.
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	<0.1% Sodium Azide (NaN ₃)

Approx. Protein Concentrations	IgG concentration 0.5mg/ml
Immunogen	CHO cell line transfected with human TLR2 (CD282).
External Database Links	<p>UniProt: O60603 Related reagents</p> <p>Entrez Gene: 7097 TLR2 Related reagents</p>
Synonyms	TIL4
RRID	AB_931710
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the NS0 myeloma cell line.
Specificity	<p>Mouse anti Human CD282 antibody, clone TL2.3 recognizes human TLR2, otherwise known as CD282. TLR2 is a member of the Toll-like receptor (TLR) family and is expressed primarily by peripheral blood monocytes.</p> <p>TLRs are expressed on the cell surface and the endocytic compartment and recognize pathogen-associated molecular patterns (PAMPs) that are expressed on infectious agents. They also initiate cell signalling to induce production of cytokines necessary for the innate immunity and subsequent adaptive immunity. TLR2 is reported to respond to a diverse range of bacterial cell wall components, mediating the innate immune response in co-operation with MD-2.</p> <p>Mouse anti Human CD282 antibody, clone TL2.1 is reported to block TLR2 function.</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
Histology Positive Control Tissue	Tonsil
References	<ol style="list-style-type: none"> 1. Flo, T.H. <i>et al.</i> (2001) Differential expression of Toll-like receptor 2 in human cells. J Leukoc Biol. 69 (3): 474-81. 2. Yilmaz, A. <i>et al.</i> (2006) Differential effects of statins on relevant functions of human monocyte-derived dendritic cells. J Leukoc Biol. 79 (3): 529-38. 3. Lee, R.M. <i>et al.</i> (2006) Influenza A viruses upregulate neutrophil toll-like receptor 2 expression and function. Scand J Immunol. 63 (2): 81-9. 4. Karlsson, K.R. <i>et al.</i> (2008) Homogeneous monocytes and macrophages from human embryonic stem cells following coculture-free differentiation in M-CSF and IL-3. Exp Hematol. 36 (9): 1167-75. 5. Sels, J.W. <i>et al.</i> (2012) Fractional flow reserve is not associated with inflammatory markers in patients with stable coronary artery disease. PLoS One. 7 (10): e46356. 6. Jankovicova, K. <i>et al.</i> (2012) TLR2 in pleural fluid is modulated by talc particles during

- pleurodesis. [Clin Dev Immunol. 2012: 158287.](#)
7. Krejsek, J. *et al.* (2013) TLR2 and TLR4 expression on blood monocytes and granulocytes of cardiac surgical patients is not affected by the use of cardiopulmonary bypass. [Acta Medica \(Hradec Kralove\). 56 \(2\): 57-66.](#)
 8. Krejsek, J. *et al.* (2013) TLR2 and TLR4 expression on blood monocytes and granulocytes of cardiac surgical patients is not affected by the use of cardiopulmonary bypass. [Acta Medica \(Hradec Kralove\). 56 \(2\): 57-66.](#)
 9. Jaedicke, K.M. *et al.* (2013) Leptin up-regulates TLR2 in human monocytes. [J Leukoc Biol. 93 \(4\): 561-71.](#)
 10. Al-Hassi, H.O. *et al.* (2014) Altered human gut dendritic cell properties in ulcerative colitis are reversed by *Lactobacillus plantarum* extracellular encrypted peptide STp. [Mol Nutr Food Res. 58 \(5\): 1132-43.](#)
 11. Mann, E.R. *et al.* (2014) Human gut dendritic cells drive aberrant gut-specific t-cell responses in ulcerative colitis, characterized by increased IL-4 production and loss of IL-22 and IFN γ . [Inflamm Bowel Dis. 20 \(12\): 2299-307.](#)
 12. Larsson, O. *et al.* (2018) Substance P represents a novel first-line defense mechanism in the nose. [J Allergy Clin Immunol. 141 \(1\): 128-136.e3.](#)

Further Reading 1. Lien, E. *et al.* (1999) Toll-like receptor 2 functions as a pattern recognition receptor for diverse bacterial products. [J Biol Chem. 274 \(47\): 33419-25.](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/MCA2152GA>
10040

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

- | | |
|---|--|
| Goat Anti Mouse IgG (STAR77...) | HRP |
| Rabbit Anti Mouse IgG (STAR12...) | RPE |
| Goat Anti Mouse IgG IgA IgM (STAR87...) | Alk. Phos. , HRP |
| Goat Anti Mouse IgG (STAR76...) | RPE |
| Goat Anti Mouse IgG (Fc) (STAR120...) | FITC , HRP |
| Rabbit Anti Mouse IgG (STAR13...) | HRP |
| Goat Anti Mouse IgG (STAR70...) | FITC |

Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight®488](#), [DyLight®550](#),
[DyLight®650](#), [DyLight®680](#), [DyLight®800](#),
[FITC](#), [HRP](#)

Rabbit Anti Mouse IgG (STAR9...) [FITC](#)

Recommended Negative Controls

[MOUSE IgG2a NEGATIVE CONTROL \(MCA929\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M389384:210806'

Printed on 12 Aug 2023

© 2023 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)