

## Datasheet: MCA2155T

<b>Description:</b>	MOUSE ANTI HUMAN CD206
<b>Specificity:</b>	CD206
<b>Other names:</b>	MANNOSE RECEPTOR C TYPE 1
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	15-2
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	25 µg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting (1)	▪			
Immunofluorescence	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

**(1)Clone 15-2 recognises a protein of approximately 175kDa under non-reducing conditions.**

<b>Target Species</b>	Human
<b>Product Form</b>	Purified IgG - liquid
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.09% Sodium Azide
<b>Approx. Protein Concentrations</b>	IgG concentration 0.5 mg/ml

<b>Immunogen</b>	Purified human mannose receptor.
<b>External Database Links</b>	<p><b>UniProt:</b>  <a href="#">P22897</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">4360</a>    MRC1    <a href="#">Related reagents</a></p>
<b>Synonyms</b>	CLEC13D
<b>RRID</b>	AB_2144910
<b>Fusion Partners</b>	Spleen cells from immunised Balb/c mice where fused with cells of the SP2/0 Ag.14 mouse myeloma cell line.
<b>Specificity</b>	<p><b>Mouse anti Human CD206 monoclonal antibody, clone 15-2</b> recognizes human macrophage mannose receptor 1, also known as CD206 or C-type lectin domain family 13 member D-like. CD206 is a ~175 kDa single pass type I transmembrane glycoprotein belonging to the group of pattern recognition receptors (<a href="#">Paveley et al. 2011</a>). CD206 has multiple carbohydrate recognition motifs and acts as a receptor for bacteria, fungi and other pathogens (<a href="#">Ezekowitz et al. 1990</a>). CD206 is predominantly expressed in tissue macrophages and dendritic cells (<a href="#">Engering et al. 1997</a>) and can also be found in a subpopulation of endothelial cells (<a href="#">Pack et al. 2007</a>) and sperm cells (<a href="#">Cardona-Maya et al. 2006</a>). CD206 can also be detected in a soluble form in human plasma and is elevated in patients with acute sepsis (<a href="#">Rødgaard-Hansen et al. 2013</a>).</p> <p>Mouse anti CD206, clone 15-2 has been used extensively to monitor mannose receptor modulation in macrophages treated with a wide range of cytokines and growth factors (<a href="#">Chang et al. 2004</a>) and to indicate CD206 as a marker for alternative activation of macrophages (<a href="#">Joerink et al. 2011</a>).</p>
<b>Flow Cytometry</b>	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul.
<b>References</b>	<ol style="list-style-type: none"> <li>1. Barrett-Bergshoeff, M. <i>et al.</i> (1997) Monoclonal antibodies against the human mannose receptor that inhibit the binding of tissue-type plasminogen activator. <a href="#">Thromb Haemost. 77: 718-24.</a></li> <li>2. Koning, N. <i>et al.</i> (2009) Distribution of the immune inhibitory molecules CD200 and CD200R in the normal central nervous system and multiple sclerosis lesions suggests neuron-glia and glia-glia interactions. <a href="#">J Neuropathol Exp Neurol. 68: 159-67.</a></li> <li>3. Emara, M. <i>et al.</i> (2011) Recognition of the major cat allergen Fel d 1 through the cysteine-rich domain of the mannose receptor determines its allergenicity. <a href="#">J Biol Chem. 286:13033-40.</a></li> <li>4. Chang, S.K. <i>et al.</i> (2008) B lymphocyte stimulator regulates adaptive immune responses by directly promoting dendritic cell maturation. <a href="#">J Immunol. 180: 7394-403.</a></li> <li>5. MacKinnon, A.C. <i>et al.</i> (2008) Regulation of alternative macrophage activation by galectin-3. <a href="#">J Immunol. 180: 2650-8.</a></li> <li>6. Lai, W.K. <i>et al.</i> (2006) Expression of DC-SIGN and DC-SIGNR on human sinusoidal</li> </ol>

- endothelium: a role for capturing hepatitis C virus particles. [Am J Pathol. 169: 200-8.](#)
7. Ueno, N. *et al.* (2009) Differences in human macrophage receptor usage, lysosomal fusion kinetics and survival between logarithmic and metacyclic *Leishmania infantum chagasi* promastigotes. [Cell Microbiol. 11: 1827-41.](#)
  8. Kato, M. *et al.* (2000) Expression of multilectin receptors and comparative FITC-dextran uptake by human dendritic cells. [Int Immunol. 12:1511-9.](#)
  9. Chang, Y.C. *et al.* (2004) Modulation of macrophage differentiation and activation by decoy receptor 3. [J Leukoc Biol. 75: 486-94.](#)
  10. Yamamoto, H. *et al.* (2011) Sphingosylphosphorylcholine and lysosulfatide have inverse regulatory functions in monocytic cell differentiation into macrophages. [Arch Biochem Biophys. 506: 83-91.](#)
  11. He, L.Z. *et al.* (2007) Antigenic targeting of the human mannose receptor induces tumor immunity. [J Immunol. 178: 6259-67.](#)
  12. Sturge, J. *et al.* (2007) Mannose receptor regulation of macrophage cell migration. [J Leukoc Biol. 82: 585-93.](#)
  13. Torrelles, J.B. *et al.* (2006) Fine discrimination in the recognition of individual species of phosphatidyl-myo-inositol mannosides from *Mycobacterium tuberculosis* by C-type lectin pattern recognition receptors. [J Immunol. 177 \(3\): 1805-16.](#)
  14. Larsson, K. *et al.* (2015) COX/mPGES-1/PGE2 pathway depicts an inflammatory-dependent high-risk neuroblastoma subset. [Proc Natl Acad Sci U S A. 112 \(26\): 8070-5.](#)
  15. Stankevich, K.S. *et al.* (2015) Surface modification of biomaterials based on high-molecular polylactic acid and their effect on inflammatory reactions of primary human monocyte-derived macrophages: perspective for personalized therapy. [Mater Sci Eng C Mater Biol Appl. 51: 117-26.](#)
  16. Yamane, K. & Leung, K.P. (2016) Rabbit M1 and M2 macrophages can be induced by human recombinant GM-CSF and M-CSF. [FEBS Open Bio. 6 \(9\): 945-53.](#)
  17. Karna, S.L. *et al.* (2016) RNA-Seq Transcriptomic Responses of Full-Thickness Dermal Excision Wounds to *Pseudomonas aeruginosa* Acute and Biofilm Infection. [PLoS One. 11 \(10\): e0165312.](#)
  18. Özçelik H *et al.* (2015) Harnessing the multifunctionality in nature: a bioactive agent release system with self-antimicrobial and immunomodulatory properties. [Adv Healthc Mater. 4 \(13\): 2026-36.](#)
  19. Argueta-Donohué J *et al.* (2016) Differential phagocytosis of *Leishmania mexicana* promastigotes and amastigotes by monocyte-derived dendritic cells. [Microbiol Immunol. 60 \(6\): 369-81.](#)

---

**Storage**

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

---

**Guarantee**

12 months from date of despatch

---

**Health And Safety Information**

Material Safety Datasheet documentation #10040 available at: 10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

## Related Products

### Recommended Secondary Antibodies

Goat Anti Mouse IgG IgA IgM (STAR87...)	<a href="#">Alk. Phos.</a> , <a href="#">HRP</a>
Goat Anti Mouse IgG (STAR77...)	<a href="#">HRP</a>
Rabbit Anti Mouse IgG (STAR12...)	<a href="#">RPE</a>
Rabbit Anti Mouse IgG (STAR8...)	<a href="#">DyLight@800</a>
Rabbit Anti Mouse IgG (STAR13...)	<a href="#">HRP</a>
Goat Anti Mouse IgG (STAR76...)	<a href="#">RPE</a>
Goat Anti Mouse IgG (STAR70...)	<a href="#">FITC</a>
Goat Anti Mouse IgG (Fc) (STAR120...)	<a href="#">FITC</a> , <a href="#">HRP</a>
Rabbit Anti Mouse IgG (STAR9...)	<a href="#">FITC</a>
Goat Anti Mouse IgG (H/L) (STAR117...)	<a href="#">Alk. Phos.</a> , <a href="#">DyLight@488</a> , <a href="#">DyLight@680</a> , <a href="#">DyLight@800</a> , <a href="#">FITC</a> , <a href="#">HRP</a>

### Recommended Negative Controls

#### [MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

From March 15, 2021, we will no longer supply printed datasheets with our products.  
Look out for updates on how to access your digital version at [bio-rad-antibodies.com](http://bio-rad-antibodies.com)

'M366268:200529'

**Printed on 12 Feb 2021**