

Datasheet: MCA216

BATCH NUMBER 154776

Description:	MOUSE ANTI HUMAN CD66b
Specificity:	CD66b
Other names:	BILIARY GLYCOPROTEIN, CARCINOEMBRYONIC ANTIGEN RELATED CELL ADHESION MOLECULE 8
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	80H3
Isotype:	IgG1
Quantity:	0.2 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/20 - 1/100
Immunohistology - Frozen (1)	▪			1/25 - 1/100
Immunohistology - Paraffin		▪		
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting (2)	▪			
Functional Assays (3)	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1) **The epitope recognised by this antibody is reported to be sensitive to formaldehyde fixation and tissue processing. Bio-Rad recommends the use of acetone fixation for frozen sections.**

(2) See [Feuk-Lagerstedt et al.](#)

(3) **This product contains sodium azide, removal by dialysis is recommended prior to use in functional assays. Dialysis cassettes [EQU003](#) are suitable for this purpose.**

Target Species	Human
-----------------------	-------

Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
Buffer Solution	Phosphate buffered saline.
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃) 0.5% Bovine Serum Albumin
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Human leucocytes from individuals with CML.
External Database Links	<p>UniProt: P31997 Related reagents</p> <p>Entrez Gene: 1088 CEACAM8 Related reagents</p>
Synonyms	CGM6
RRID	AB_2229261
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the mouse MOPC 315-43 myeloma cell line.
Specificity	Mouse anti Human CD66b antibody, clone 80H3 reacts with the CEACAM8 or CGM6 gene product, from the carcinoembryonic gene family. The CEACAM8 gene product is called CD66b, Non-specific cross-reacting antigen NCA-95 or Carcinoembryonic antigen CGM6. CD66b is a 286 amino acid mature ~95-100 kDa GPI anchored molecule with an N-terminal 34 amino acid signal peptide and a 29 amino acid C-terminal propeptide, shed in the mature form. CD66b is expressed strongly by mature granulocytes and metamyelocytes, and weakly by bone marrow myelocytes. CD66b is not expressed by peripheral blood monocytes or lymphocytes.
Flow Cytometry	Use 10ul of the suggested working dilution to label 5 x 10 ⁵ cells or 100ul whole blood.
Histology Positive Control Tissue	Bone Marrow
References	<ol style="list-style-type: none"> Mannoni, P. <i>et al.</i> (1982) Monoclonal antibodies against human granulocytes and myeloid differentiation antigens. Hum Immunol. 5 (4): 309-23. Ionita, M.G. <i>et al.</i> (2010) High neutrophil numbers in human carotid atherosclerotic plaques are associated with characteristics of rupture-prone lesions. Arterioscler Thromb Vasc Biol. 30 (9): 1842-8. Simard, J.C. <i>et al.</i> (2010) Induction of neutrophil degranulation by S100A9 via a

- MAPK-dependent mechanism. [J Leukoc Biol. 87 \(5\): 905-14.](#)
4. Feuk-Lagerstedt, E. *et al.* (1999) Identification of CD66a and CD66b as the major galectin-3 receptor candidates in human neutrophils. [J Immunol. 163: 5592-8.](#)
 5. Jankowski, A. *et al.* (2002) Determinants of the phagosomal pH in neutrophils. [J Biol Chem. 277: 6059-66.](#)
 6. Jinnouchi, A. *et al.* (2005) Local anesthetics inhibit priming of neutrophils by lipopolysaccharide for enhanced release of superoxide: suppression of cytochrome b558 expression by disparate mechanisms. [J Leukoc Biol. 78: 1356-65.](#)
 7. Thickett, D.R. *et al.* (2002) A role for vascular endothelial growth factor in acute and resolving lung injury. [Am J Respir Crit Care Med. 166: 1332-7.](#)
 8. Sekine, K. *et al.* (2006) Panning of multiple subsets of leukocytes on antibody-decorated poly(ethylene) glycol-coated glass slides. [J Immunol Methods. 313: 96-109.](#)
 9. Ottonello, L. *et al.* (1999) Monoclonal Lym-1 antibody-dependent cytolysis by neutrophils exposed to granulocyte-macrophage colony-stimulating factor: intervention of FcγRII (CD32), CD11b-CD18 integrins, and CD66b glycoproteins. [Blood. 93: 3505-11.](#)
 10. Ottonello, L. *et al.* (2000) Monoclonal Lym-1 antibody-targeted lysis of B lymphoma cells by neutrophils. Evidence for two mechanisms of FcγRII-dependent cytolysis. [J Leukoc Biol. 68: 662-8.](#)
 11. Moraes, T.J. *et al.* (2006) Abnormalities in the pulmonary innate immune system in cystic fibrosis. [Am J Respir Cell Mol Biol. 34: 364-74.](#)
 12. Nijhuis, J. *et al.* (2009) Neutrophil Activation in Morbid Obesity, Chronic Activation of Acute Inflammation [Obesity \(Silver Spring\). 17: 2014-8.](#)
 13. Pliyev, B.K. and Menshikov, M.Y. (2010) Release of the soluble urokinase-type plasminogen activator receptor (suPAR) by activated neutrophils in rheumatoid arthritis. [Inflammation. 33: 1-9.](#)
 14. Binet, F. and Girard, D. (2008) Novel human neutrophil agonistic properties of arsenic trioxide: involvement of p38 mitogen-activated protein kinase and/or c-jun NH2-terminal MAPK but not extracellular signal-regulated kinases-1/2. [J Leukoc Biol. 84: 1613-22.](#)
 15. Rosenkranz, A.R. *et al.* (1999) Novel C5-dependent mechanism of neutrophil stimulation by bioincompatible dialyzer membranes. [J Am Soc Nephrol. 10: 128-35.](#)
 16. Simard, J.C. *et al.* (2010) Induction of neutrophil degranulation by S100A9 via a MAPK-dependent mechanism. [J Leukoc Biol. 87: 905-14.](#)
 17. Sekine, K. *et al.* (2006) Panning of multiple subsets of leukocytes on antibody-decorated poly(ethylene) glycol-coated glass slides [J Immunol Methods. 313: 96-109.](#)
 18. Orozco-Uribe, M. *et al.* (2015) Early Exposure of Human Neutrophils to Mycobacteria Triggers Cell Damage and Pro-Inhibitory Molecules, but not Activation [Tuberculosis Ch4](#)
 19. Whitmore, L.C. *et al.* (2016) A Common Genetic Variant in TLR1 Enhances Human Neutrophil Priming and Impacts Length of Intensive Care Stay in Pediatric Sepsis. [J Immunol. 196 \(3\): 1376-86.](#)
 20. Bastian, O.W. *et al.* (2016) Neutrophils contribute to fracture healing by synthesizing fibronectin+ extracellular matrix rapidly after injury. [Clin Immunol. 164: 78-84.](#)
 21. Dyugovskaya, L. *et al.* (2016) Intermittent Hypoxia Affects the Spontaneous Differentiation *In Vitro* of Human Neutrophils into Long-Lived Giant Phagocytes. [Oxid Med Cell Longev. 2016: 9636937.](#)
 22. Medford, A.J. *et al.* (2019) Blood-based monitoring identifies acquired and targetable driver *HER2* mutations in endocrine-resistant metastatic breast cancer. [NPJ Precis Oncol.](#)

[3: 18.](#)

23. Ebright, R.Y. *et al.* (2020) HIF1A signaling selectively supports proliferation of breast cancer in the brain. [Nat Commun. 11 \(1\): 6311.](#)

Storage Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10041 available at: <https://www.bio-rad-antibodies.com/SDS/MCA216>
10041

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR77...) [HRP](#)
Rabbit Anti Mouse IgG (STAR12...) [RPE](#)
Goat Anti Mouse IgG IgA IgM (STAR87...) [Alk. Phos.](#), [HRP](#)
Goat Anti Mouse IgG (STAR76...) [RPE](#)
Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)
Rabbit Anti Mouse IgG (STAR13...) [HRP](#)
Goat Anti Mouse IgG (STAR70...) [FITC](#)
Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight®488](#), [DyLight®550](#),
[DyLight®650](#), [DyLight®680](#), [DyLight®800](#),
[FITC](#), [HRP](#)
Rabbit Anti Mouse IgG (STAR9...) [FITC](#)

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M373003:200811'

Printed on 12 Aug 2023

© 2023 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)