

Datasheet: MCA2164GA

BATCH NUMBER 161440

Description:	MOUSE ANTI CHICKEN CD4
Specificity:	CD4
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	2-35
Isotype:	IgG2b
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/50 - 1/100
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting			▪	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Chicken
Species Cross Reactivity	<p>Reacts with: Turkey</p> <p>N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.</p>
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant

Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Chicken embryonic thymocytes
Fusion Partners	Lymph node cells from immunised Balb/c mice were fused with cells of the SP2/0 myeloma cell line
Specificity	Mouse anti Chicken CD4, clone 2-35 recognizes the chicken homologue of human CD4, a ~64 kDa cell surface protein expressed by thymocytes and a subset of T cells (Luhtala et al. 1993). Mouse anti Chicken CD4, clone 2-35 has been demonstrated to recognize turkey CD4 (Li et al. 1998).
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul
References	<ol style="list-style-type: none"> Vainio, O. <i>et al.</i> (1989) Characterization of chicken CD4-expressing cells. Prog Clin Biol Res. 307: 45-56. Li, Z. <i>et al.</i> (1999) Cross-reactive anti-chicken CD4 and CD8 monoclonal antibodies suggest polymorphism of the turkey CD8alpha molecule. Poult Sci. 78 (11): 1526-31. Koskinen, R. <i>et al.</i> (1999) Cloning and modeling of the first nonmammalian CD4. J Immunol. 162 (7): 4115-21. Pavlova, S.P. <i>et al.</i> (2010) <i>In vitro</i> and <i>in vivo</i> characterization of glycoprotein C-deleted infectious laryngotracheitis virus. J Gen Virol. 91 (Pt 4): 847-57. Luhtala, M. <i>et al.</i> (1993) Analysis of chicken CD4 by monoclonal antibodies indicates evolutionary conservation between avian and mammalian species. Hybridoma. 12: 633-46. Rosa, A.C. <i>et al.</i> (2014) Isolation and molecular characterization of Brazilian turkey reovirus from immunosuppressed young poults. Arch Virol. 159 (6): 1453-7. Blohm, U. <i>et al.</i> (2016) Immunological Competence of Different Domestic Chicken Breeds Against Avian Influenza Infection. Avian Dis. 60 (1 Suppl): 262-8. Röhe, I. <i>et al.</i> (2017) Effect of feeding soybean meal and differently processed peas on the gut mucosal immune system of broilers. Poult Sci. 96 (7): 2064-73. Sachan, S. <i>et al.</i> (2015) Adjuvant potential of resiquimod with inactivated Newcastle disease vaccine and its mechanism of action in chicken. Vaccine. 33 (36): 4526-32. Kannan, T.A. <i>et al.</i> (2017) Age Related Changes in T Cell Subsets in Thymus and Spleen of Layer Chicken (<i>Gallus domesticus</i>) Int J Curr Microbiol App Sci. 6 (1): 15-19. Tang, Y. <i>et al.</i> (2020) Immune Modulation and the Development of Fowl Typhoid: A Model of Human Disease? Pathogens. 9 (10): 843. Konieczka, P. <i>et al.</i> (2022) Increased arginine, lysine, and methionine levels can improve the performance, gut integrity and immune status of turkeys but the effect is interactive and depends on challenge conditions. Vet Res. 53 (1): 59.

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/MCA2164GA>
10040

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...) [RPE](#)

Goat Anti Mouse IgG (H/L) (STAR117...) [FITC](#)

Rabbit Anti Mouse IgG (STAR13...) [HRP](#)

Rabbit Anti Mouse IgG (STAR9...) [FITC](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)
'M383139:210513'

Printed on 25 Mar 2023

© 2023 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)