

Datasheet: MCA2166F

Description:	MOUSE ANTI CHICKEN CD8 ALPHA:FITC
Specificity:	CD8 ALPHA
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	11-39
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

RRID AB_2075649

Applications This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat - 1/5

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species Chicken

Species Cross Reactivity Reacts with: Turkey
N.B. Antibody reactivity and working conditions may vary between species.

Product Form Purified IgG - liquid

Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525

Preparation Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant

Buffer Solution Phosphate buffered saline

Preservative 0.09% Sodium Azide
Stabilisers 1% Bovine Serum Albumin

Approx. Protein Concentrations IgG concentration 0.1 mg/ml

Immunogen Chicken T-cells.

Fusion Partners	Lymph node cells from immunised Balb/c mice were fused with cells of the SP2/0 myeloma cell line.
Specificity	<p>Mouse anti chicken CD8 alpha, clone 11-39 recognizes the alpha chain of the chicken CD8 homologue, a 33-35 kDa cell surface protein. CD8 is expressed as either alpha/alpha homodimers or alpha/beta heterodimers on a subpopulation of T cells and NK cells. Mouse anti chicken CD8 alpha, clone 11-39 recognises all polymorphic forms of chicken CD8 alpha.</p> <p>Mouse anti chicken CD8 alpha, clone 11-39 has been demonstrated to cross react with Turkey (Li et al. 1999).</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
References	<ol style="list-style-type: none"> 1. Luhtala, M. <i>et al.</i> (1995) Characterization of chicken CD8-specific monoclonal antibodies recognizing novel epitopes. Scand J Immunol. 42 (1): 171-4. 2. Luhtala, M. <i>et al.</i> (1997) Polymorphism of chicken CD8-alpha, but not CD8-beta. Immunogenetics. 46 (5): 396-401. 3. Li, Z. <i>et al.</i> (1999) Cross-reactive anti-chicken CD4 and CD8 monoclonal antibodies suggest polymorphism of the turkey CD8alpha molecule. Poult Sci. 78 (11): 1526-31. 4. McKenna, G.F. (2003) Immunopathologic investigations with an attenuated chicken anemia virus in day-old chickens. Avian Dis. 47: 1339-45. 5. Morimura, T. <i>et al.</i> (1996) Apoptosis and CD8-down-regulation in the thymus of chickens infected with Marek's disease virus. Arch Virol. 141 (11): 2243-9. 6. Luhtala M (1998) Chicken CD4, CD8alphabeta, and CD8alphaalpha T cell co-receptor molecules. Poult Sci. 77 (12): 1858-73. 7. Imhof, B.A. <i>et al.</i> (2000) Intestinal CD8 alpha alpha and CD8 alpha beta intraepithelial lymphocytes are thymus derived and exhibit subtle differences in TCR beta repertoires. J Immunol. 165 (12): 6716-22. 8. Arstila, T.P. & Lassila, O. (1993) Androgen-induced expression of the peripheral blood gamma delta T cell population in the chicken. J Immunol. 151 (12): 6627-33. 9. Bohls, R.L. <i>et al.</i> (2006) The use of flow cytometry to discriminate avian lymphocytes from contaminating thrombocytes. Dev Comp Immunol. 30 (9): 843-50. 10. Powell, F.L. <i>et al.</i> (2009) The turkey, compared to the chicken, fails to mount an effective early immune response to <i>Histomonas meleagridis</i> in the gut. Parasite Immunol. 31 (6): 312-27. 11. Katevuo, K. & Vainio, O. (1996) Thymocyte emigration in the chicken: an over-representation of CD4+ cells over CD8+ in the periphery. Immunology. 89 (3): 419-23. 12. Morimura, T. <i>et al.</i> (1995) Immunomodulation of peripheral T cells in chickens infected with Marek's disease virus: involvement in immunosuppression. J Gen Virol. 76 (Pt 12): 2979-85. 13. Powell, F. <i>et al.</i> (2009) Development of reagents to study the turkey's immune response: Identification and molecular cloning of turkey CD4, CD8α and CD28. Dev Comp Immunol. 33 (4): 540-6. 14. Juul-Madsen, H.R. <i>et al.</i> (2002) Major histocompatibility complex-linked immune response of young chickens vaccinated with an attenuated live infectious bursal disease virus vaccine followed by an infection. Poult Sci. 81 (5): 649-56. 15. Wang, Y. <i>et al.</i> (2003) A novel method to analyze viral antigen-specific cytolytic activity in the chicken utilizing flow cytometry. Vet Immunol Immunopathol. 95 (1-2): 1-9. 16. Arstila, T.P. <i>et al.</i> (1995) Primed avian $\gamma\delta$ T cells respond to mycobacterial antigens, but show no preference for the 65-kDa heat shock protein. Cell Immunol. 162 (1): 74-9. 17. Arstila, T.P. <i>et al.</i> (1994) $\gamma\delta$ and $\alpha\beta$ T cells are equally susceptible to apoptosis. Scand J Immunol. 40 (2): 209-15. 18. Rosa, A.C. <i>et al.</i> (2014) Isolation and molecular characterization of Brazilian turkey reovirus from immunosuppressed young poults. Arch Virol. 159 (6): 1453-7. 19. Röhe I. <i>et al.</i> (2017) Effect of feeding soybean meal and differently processed peas on the gut

mucosal immune system of broilers [Poultry Science. Feb 23 \[Epub ahead of print\]](#)
20. Abd El-Hack, M. & Alagawany, M. (2015) Performance, egg quality, blood profile, immune function, and antioxidant enzyme activities in laying hens fed diets with thyme powder [Journal of Animal and Feed Sciences. 24 \(2\): 127-33.](#)
21. Kannan, T.A. *et al.* (2017) Age Related Changes in T Cell Subsets in Thymus and Spleen of Layer Chicken (*Gallus domesticus*) [Int J Curr Microbiol Appl Sci. 6 \(1\): 15-9.](#)

Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee

18 months from date of despatch.

Health And Safety Information

Material Safety Datasheet documentation #10041 available at:
10041: <https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf>

Regulatory

For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:FITC \(MCA928F\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M334529:181203'

Printed on 20 May 2019

© 2019 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)