Datasheet: MCA2175 BATCH NUMBER 164458

Description:	RAT ANTI HUMAN CD195
Specificity:	CD195
Other names:	CCR5
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	HEK/1/85a
Isotype:	lgG2a
Quantity:	0.2 mg

Product Details

Applications This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u>.

		0010.			
		Yes	No	Not Determined	Suggested Dilution
	Flow Cytometry				1/50 - 1/100
	Immunohistology - Frozen			•	
	Immunohistology - Paraffin			•	
	ELISA			•	
	Immunoprecipitation			•	
	Western Blotting			•	
	Where this product has n	ot been t	ested for	use in a particular tech	nique this does not
	necessarily exclude its us			•	•
	•				
	a guide only. It is recomm	nended th	hat the us	er titrates the product f	or use in their own
	system using appropriate	negative	/positive	controls.	
Target Species	Human				
Product Form	Purified IgG - liquid				
Preparation	Purified IgG prepared by	affinity cl	nromatog	aphy on Protein G fro	m tissue culture
	supernatant				
	•				
Buffer Solution	Phosphate buffered salin	е			
	· · · · · · · · · · · · · · · · · · ·				
Preservative					
Stabilisers	0.09% sodium azide (Nal	N ₃)			

Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	CHO cells transfected with human CCR5
External Database Links	UniProt: <u>P51681</u> <u>Related reagents</u> Entrez Gene: <u>1234</u> CCR5 <u>Related reagents</u>
Synonyms	CMKBR5
RRID	AB_324334
Fusion Partners	Spleen cells from immunized rats were fused with cell of the rat Y3 myeloma cell line
Specificity	Rat anti Human CD195 antibody, clone HEK/1/85a recognizes the human CD195 cell surface antigen, a 45 kDa glycoprotein also known as CCR5.
	CD195 acts as a receptor for a number of chemokines including RANTES and eotaxin and serves as a co-receptor for the entry of HIV into cells. CD195 is expressed by a subset of T lymphocytes and by monocytes.
Flow Cytometry	Use 10µl of the suggested working dilution to label 100µl fresh whole blood
References	 Mueller, A. <i>et al.</i> (2002) Pathways for internalization and recycling of the chemokine receptor CCR5. <u>Blood 99 (3): 785-91.</u> Ramirez, R. <i>et al.</i> (2006) Microinflammation in hemodialysis is related to a preactivated subset of monocytes. <u>Hemodial Int. 10 Suppl 1:S24-7.</u> Liptrott, N.J. <i>et al.</i> (2009) The impact of cytokines on the expression of drug transporters, cytochrome P450 enzymes and chemokine receptors in human PBMC. <u>Br J Pharmacol. 156: 497-508.</u> Pilch-Cooper, H.A. <i>et al.</i> (2011) Circulating human CD4 and CD8 T cells do not have large intracellular pools of CCR5. <u>Blood 118: 1015-9.</u> Bönsch, C. <i>et al.</i> (2015) Potent Anti-HIV Chemokine Analogs Direct Post-Endocytic Sorting of CCR5. <u>PLoS One 10 (4): e0125396.</u> Holmannova, D. <i>et al.</i> (2017) Effects of conventional CPB and mini-CPB on neutrophils CD162, CD166 and CD195 expression. <u>Perfusion. 32 (2): 141-50.</u>
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

e 12 mor				
-	•			
y For res	earch purpose			
l Products				
ended Secondary	Antibodies			
ti Rat IgG (STAR16.)	DyLight®800		
iti Rat IgG (STAR17.)	<u>FITC</u>		
Rat IgG (STAR69)		<u>FITC</u>		
Rat IgG (STAR73)		<u>RPE</u>		
ti Rat IgG (STAR21.)	<u>HRP</u>		
Rat IgG (MOUSE A	DSORBED) (S	TAR71) <u>DyLight®550</u> ,	DyLight®650, [DyLight®800
Rat IgG (STAR131	.)	<u>Alk. Phos., Bic</u>	otin	
Rat IgG (STAR72)		HRP		
Tel: +1 800 265 7376 Fax: +1 919 878 3751	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739	•	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50
	https://v 10040 ry For res cl Products bended Secondary ati Rat IgG (STAR16. ati Rat IgG (STAR17. Rat IgG (STAR69) Rat IgG (STAR73) ati Rat IgG (STAR73) ati Rat IgG (STAR73) Rat IgG (STAR73) Ati Rat IgG (STAR73) Ati Rat IgG (STAR73) Ati Rat IgG (STAR73) Ati Rat IgG (STAR72) Tel: +1 800 265 7376	Image: marked secondary Antibodies Image: marked secondary Antis Image: marked secon	Image: Instant sector of the sector of th	https://www.bio-rad-antibodies.com/SDS/MCA2175 10040 ry For research purposes only d Products beended Secondary Antibodies nti Rat IgG (STAR16) DyLight®800 nti Rat IgG (STAR17) FITC Rat IgG (STAR69) FITC Rat IgG (STAR73) RPE nti Rat IgG (STAR17) HRP Rat IgG (STAR17) HRP Rat IgG (STAR17) HRP Rat IgG (STAR131) Alk. Phos., Biotin Rat IgG (STAR131) Alk. Phos., Biotin Rat IgG (STAR131) HRP Rat IgG (STAR72) HRP Tet: +1 800 265 7376 Worldwide Tet: +44 (0)1865 852 700 Europe

'M413164:221119'

Printed on 23 Jan 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint