

Datasheet: MCA2175F

Description:	RAT ANTI HUMAN CD195:FITC
Specificity:	CD195
Other names:	CCR5
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	HEK/1/85a
lsotype:	lgG2a
Quantity:	0.1 mg

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .						
		Yes	No	Not Determined	Suggested Dilution		
	Flow Cytometry	-			Neat - 1/10		
	Where this product has not been tested for use in a particular technique this does not						
	necessarily exclude its a guide only. It is recor system using appropria	nmended that	the use	r titrates the product	ng dilutions are given as for use in their own		
Target Species	Human						
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid						
Max Ex/Em	Fluorophore FITC	Excitation Ma 490	x (nm)	Emission Max (nm) 525			
Preparation	Purified IgG prepared b supernatant	by affinity chro	omatogra	aphy on Protein G fro	om tissue culture		
Buffer Solution	Phosphate buffered saline						
Preservative Stabilisers	0.09% sodium azide (NaN ₃) 1% bovine serum albumin						
Approx. Protein Concentrations	IgG concentration 0.1 r	mg/ml					

Immunogen	CHO cells transfected with human CCR5	
External Database Links	UniProt: <u>P51681</u> <u>Related reagents</u> Entrez Gene: <u>1234</u> CCR5 <u>Related reagents</u>	
Synonyms	CMKBR5	
RRID	AB_324066	
Fusion Partners	Spleen cells from immunized rats were fused with cell of the rat Y3 myeloma cell line	
Specificity	Rat anti Human CD195 antibody, clone HEK/1/85a recognizes the human CD195 cell surface antigen, a 45 kDa glycoprotein also known as CCR5.	
	CD195 acts as a receptor for a number of chemokines including RANTES and eotaxin and serves as a co-receptor for the entry of HIV into cells. CD195 is expressed by a subset of T lymphocytes and by monocytes.	
Flow Cytometry	Use 10μ I of the suggested working dilution to label 100μ I fresh whole blood	
References	 Mueller, A. <i>et al.</i> (2002) Pathways for internalization and recycling of the chemokine receptor CCR5. <u>Blood 99 (3): 785-91.</u> Ramirez, R. <i>et al.</i> (2006) Microinflammation in hemodialysis is related to a preactivated subset of monocytes. <u>Hemodial Int. 10 Suppl 1:S24-7.</u> Liptrott, N.J. <i>et al.</i> (2009) The impact of cytokines on the expression of drug transporters, cytochrome P450 enzymes and chemokine receptors in human PBMC. <u>Br J Pharmacol. 156: 497-508.</u> Pilch-Cooper, H.A. <i>et al.</i> (2011) Circulating human CD4 and CD8 T cells do not have large intracellular pools of CCR5. <u>Blood 118: 1015-9.</u> Bönsch, C. <i>et al.</i> (2015) Potent Anti-HIV Chemokine Analogs Direct Post-Endocytic Sorting of CCR5. <u>PLoS One 10 (4): e0125396.</u> Holmannova, D. <i>et al.</i> (2017) Effects of conventional CPB and mini-CPB on neutrophils CD162, CD166 and CD195 expression. <u>Perfusion. 32 (2): 141-50.</u> 	
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.	
Guarantee	12 months from date of despatch	

Health And Safety	Material Safety Datasheet documentation #10041 available at:	
Information	https://www.bio-rad-antibodies.com/SDS/MCA2175F	
	10041	

Regulatory

For research purposes only

Related Products

Recommended Negative Controls

RAT IgG2a NEGATIVE CONTROL:FITC (MCA6005F)

Recommended Useful Reagents

HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-ra	id.com	Email: antibody_sales_uk@bio-	rad.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M413161:221119'

Printed on 23 Jan 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint