

Datasheet: MCA2179

BATCH NUMBER 150117

Description:	RAT ANTI CCT ETA
Specificity:	CCT ETA
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	PK/16/8/a
Isotype:	IgG2a
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen			▪	
Immunohistology - Paraffin (1)	▪			1/100
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting	▪			1/500

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1) This product requires antigen retrieval using heat treatment prior to staining of paraffin sections.

Species Cross Reactivity

Reacts with: Mouse, Human, Bovine, Rabbit

N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

Product Form

Purified IgG - liquid

Preparation

Purified IgG prepared by affinity chromatography on Protein G from tissue culture

	supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
External Database Links	<p>UniProt: Q2NKZ1 Related reagents</p> <p>Entrez Gene: 514355 CCT7 Related reagents</p>
RRID	AB_324836
Specificity	<p>Rat anti CCTeta antibody, clone PK/16/8/a recognizes the eta polypeptide of the CCT chaperonin molecule complex. CCTη is a 543 amino acid ~60 kDa molecule encoded by the CCT7 gene, a member of the TCP-1 chaperonin family (UniProt: Q2NKZ1)</p> <p>The intact CCT complex is composed of eight polypeptides in a double-ring structure. CCT is important within cells in aiding the folding of proteins including actin, tubulin and the VHL tumour suppressor protein.</p> <p>Rat anti CCTeta antibody, clone PK/16/8/a detects a band of ~60 kDa in HeLa cell lysates using western blotting under reducing conditions.</p>
Histology Positive Control Tissue	Human prostate.
Western Blotting	MCA2179 detects a band of approximately 60kD in Hela cell lysates.
References	<ol style="list-style-type: none"> 1. Satish, L. <i>et al.</i> (2010) Cloning and expression of rabbit CCT subunits eta and beta in healing cutaneous wounds. Cell Stress Chaperones. 15 (6): 819-26. 2. Hynes, G. <i>et al.</i> (1996) Analysis of chaperonin-containing TCP-1 subunits in the human keratinocyte two-dimensional protein database: further characterisation of antibodies to individual subunits. Electrophoresis. 17 (11): 1720-7. 3. Satish L <i>et al.</i> (2011) Chaperonin containing T-complex polypeptide (CCT) subunit expression in oral mucosal wounds and fibroblasts. Cell Stress Chaperones. 16 (6): 675-80. 4. Hynes, G.M. & Willison, K.R. (2000) Individual subunits of the eukaryotic cytosolic chaperonin mediate interactions with binding sites located on subdomains of beta-actin. J Biol Chem. 275 (25): 18985-94. 5. Satish, L. <i>et al.</i> (2010) Chaperonin containing T-complex polypeptide subunit eta

(CCT-eta) is a specific regulator of fibroblast motility and contractility. [PLoS One. 5\(4\):e10063.](#)

6. Satish L *et al.* (2013) Increased CCT-eta expression is a marker of latent and active disease and a modulator of fibroblast contractility in Dupuytren's contracture. [Cell Stress Chaperones. 18 \(4\): 397-404.](#)

Further Reading	1. Llorca, O. <i>et al.</i> (2000) Eukaryotic chaperonin CCT stabilizes actin and tubulin folding intermediates in open quasi-native conformations. EMBO J. 19 (22): 5971-9.
------------------------	--

Storage	Store at +4°C or at -20°C if preferred.
----------------	---

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee	12 months from date of despatch
------------------	---------------------------------

Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA2179 10040
--------------------------------------	---

Regulatory	For research purposes only
-------------------	----------------------------

Related Products

Recommended Secondary Antibodies

Rabbit Anti Rat IgG (STAR16...)	DyLight®800
Rabbit Anti Rat IgG (STAR17...)	FITC
Goat Anti Rat IgG (STAR72...)	HRP
Goat Anti Rat IgG (STAR69...)	FITC
Goat Anti Rat IgG (STAR73...)	RPE
Rabbit Anti Rat IgG (STAR21...)	HRP
Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...)	DyLight®550 , DyLight®650 , DyLight®800
Goat Anti Rat IgG (STAR131...)	Alk. Phos. , Biotin

North & South America Tel: +1 800 265 7376
Fax: +1 919 878 3751
Email: antibody_sales_us@bio-rad.com

Worldwide Tel: +44 (0)1865 852 700
Fax: +44 (0)1865 852 739
Email: antibody_sales_uk@bio-rad.com

Europe Tel: +49 (0) 89 8090 95 21
Fax: +49 (0) 89 8090 95 50
Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M366298:200529'

Printed on 19 Jan 2024