

Datasheet: MCA2183PE

BATCH NUMBER INN1607

Description:	RAT ANTI MOUSE CD13:RPE
Specificity:	CD13
Other names:	AMINOPEPTIDASE N
Format:	RPE
Product Type:	Monoclonal Antibody
Clone:	R3-63
Isotype:	IgG2a
Quantity:	100 TESTS

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their system using appropriate negative/positive controls.

Target Species	Mouse		
Product Form	Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilized		
Reconstitution	Reconstitute with 1.0 ml distilled water		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	RPE 488nm laser	496	578
Preparation	Purified IgG prepared by affinity chromatography on Protein G supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative Stabilisers	0.09% Sodium Azide		
	1% Bovine Serum Albumin		

5% Sucrose

Immunogen	Mouse intestinal APN
External Database Links	UniProt: P97449 Related reagents Entrez Gene: 16790 Anpep Related reagents
Synonyms	Lap1, Lap-1
RRID	AB_324138
Fusion Partners	Spleen cells from immunized mice were fused with cells of the IR983F rat myeloma cell line.
Specificity	<p>Rat anti Mouse CD13 antibody, clone R3-63 recognizes mouse aminopeptidase N (APN), a cell surface protein homologous with human CD13. In the mouse, CD13 is a non-covalently linked homodimer of approximately 150 kDa subunits expressed by a variety of cells including monocytes, macrophages, dendritic cell and veiled cells.</p> <p>Rat anti Mouse CD13 antibody, clone R3-63 has been reported to block mouse APN enzyme activity (Hansen <i>et al.</i> 1993).</p>
Flow Cytometry	<p>Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul.</p> <p>The Fc region of monoclonal antibodies may bind non-specifically to cells expressing low affinity fc receptors. This may be reduced by using SeroBlock FcR (BUF041A/B).</p>
References	<ol style="list-style-type: none">1. Kamoun, W.S. <i>et al.</i> (2009) Edema control by cediranib, a vascular endothelial growth factor receptor-targeted kinase inhibitor, prolongs survival despite persistent brain tumor growth in mice. J Clin Oncol. 27: 2542-52.2. Hansen, A.S. <i>et al.</i> (1993) A mouse aminopeptidase N is a marker for antigen-presenting cells and appears to be co-expressed with major histocompatibility complex class II molecules. Eur J Immunol. 23 (9): 2358-64.3. Larsen, S.L. <i>et al.</i> (1996) T cell responses affected by aminopeptidase N (CD13)-mediated trimming of major histocompatibility complex class II-bound peptides. J Exp Med. 184 (1): 183-9.4. Rangel, R. <i>et al.</i> (2007) Impaired angiogenesis in aminopeptidase N-null mice. Proc Natl Acad Sci U S A. 104: 4588-93.5. Lahdenranta, J. <i>et al.</i> (2007) Treatment of hypoxia-induced retinopathy with targeted proapoptotic peptidomimetic in a mouse model of disease. FASEB J. 21: 3272-8.6. Li, P. <i>et al.</i> (2010) Use of adenoviral vectors to target chemotherapy to tumor vascular endothelial cells suppresses growth of breast cancer and melanoma. Mol Ther. 18: 921-8.7. van Deventer, H.W. <i>et al.</i> (2008) C-C chemokine receptor 5 on pulmonary fibrocytes facilitates migration and promotes metastasis via matrix metalloproteinase 9. Am J Pathol.

[173: 253-64.](#)

8. Gabrilovac, J. *et al.* (2011) Expression, regulation and functional activities of aminopeptidase N (EC 3.4.11.2; APN; CD13) on murine macrophage J774 cell line. [Immunobiology. 216: 132-44.](#)
9. Ozawa, M.G. *et al.* (2008) Beyond receptor expression levels: the relevance of target accessibility in ligand-directed pharmacodelivery systems. [Trends Cardiovasc Med. 18: 126-32.](#)
10. Bertilaccio, M.T. *et al.* (2008) Vasculature-targeted tumor necrosis factor-alpha increases the therapeutic index of doxorubicin against prostate cancer. [Prostate. 68: 1105-15.](#)
11. Boström, M. *et al.* (2014) The hippocampal neurovascular niche during normal development and after irradiation to the juvenile mouse brain. [Int J Radiat Biol. 90: 778-89.](#)
12. Mayer-Barber, K.D. *et al.* (2011) Innate and adaptive interferons suppress IL-1 α and IL-1 β production by distinct pulmonary myeloid subsets during *Mycobacterium tuberculosis* infection. [Immunity. 35: 1023-34.](#)
13. Winnicka, B. *et al.* (2010) CD13 is dispensable for normal hematopoiesis and myeloid cell functions in the mouse. [J Leukoc Biol. 88: 347-59.](#)
14. Ridder, D.A. *et al.* (2015) Brain endothelial TAK1 and NEMO safeguard the neurovascular unit. [J Exp Med. 212 \(10\): 1529-49.](#)
15. Vanlandewijck, M. *et al.* (2015) Functional Characterization of Germline Mutations in PDGFB and PDGFRB in Primary Familial Brain Calcification. [PLoS One. 10 \(11\): e0143407.](#)
16. Körbelin J *et al.* (2016) A brain microvasculature endothelial cell-specific viral vector with the potential to treat neurovascular and neurological diseases. [EMBO Mol Med. 8 \(6\): 609-25.](#)
17. Zötz, J.S. *et al.* (2016) CD13/aminopeptidase N is a negative regulator of mast cell activation. [FASEB J. 30 \(6\): 2225-35.](#)
18. Sung, S.J. *et al.* (2017) Proximal Tubule CD73 Is Critical in Renal Ischemia-Reperfusion Injury Protection. [J Am Soc Nephrol. 28 \(3\): 888-902.](#)
19. Yanagida, K. *et al.* (2017) Size-selective opening of the blood-brain barrier by targeting endothelial sphingosine 1-phosphate receptor 1. [Proc Natl Acad Sci U S A. 114 \(17\): 4531-6.](#)
20. Elabi, O. *et al.* (2021) Human α -synuclein overexpression in a mouse model of Parkinson's disease leads to vascular pathology, blood brain barrier leakage and pericyte activation. [Sci Rep. 11 \(1\): 1120.](#)
21. Kato, T. *et al.* (2020) Excessive Production of Transforming Growth Factor β 1 Causes Mural Cell Depletion From Cerebral Small Vessels. [Front Aging Neurosci. 12: 151.](#)

Storage

Prior to reconstitution store at +4°C. Following reconstitution store at +4°C.

DO NOT FREEZE.

This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #20487 available at: https://www.bio-rad-antibodies.com/SDS/MCA2183PE 20487
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

[RAT IgG2a NEGATIVE CONTROL:RPE \(MCA1212PE\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)
'M375426:210104'

Printed on 19 Jan 2024