

Datasheet: MCA2189

Description:	MOUSE ANTI MOUSE MHC CLASS I
Specificity:	MHC CLASS I
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	2G5
Isotype:	IgG2b
Quantity:	0.25 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/10 - 1/25
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting			▪	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species

Mouse

Species Cross Reactivity

Reacts with: Rat, Guinea Pig, Sheep, Bovine, Pig, Human, Hamster

N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

Product Form

Purified IgG - liquid

Preparation

Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant

Buffer Solution

Phosphate buffered saline

Preservative Stabilisers	0.09% sodium azide (NaN ₃)
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Purified H-2K ^b and H-2D ^b MHC-I molecules.
RRID	AB_324082
Fusion Partners	Spleen cells from immunised C1D mice were fused with cells of the X63 myeloma cell line.
Specificity	<p>Mouse anti Mouse MHC Class I antibody, clone 2G5 recognizes a monomorphic epitope present on murine MHC class I molecules, expressed at varying levels on the majority of nucleated cells. The major histocompatibility complex (MHC) is a cluster of genes that are important in the immune response to infections. In mice, this complex is referred to as the histocompatibility 2 (H-2) region.</p> <p>The epitope recognized by clone 2G5 is conformation dependent and is reported to be phylogenetically conserved (Claesson et al. 1994). Reactivity has been observed with some canine samples suggesting that this antibody may recognize a polymorphic epitope of canine MHC class I.</p>
Flow Cytometry	Use 10µl of the suggested working dilution to label 10 ⁶ cells in 100µl
References	<ol style="list-style-type: none"> Perone, M.J. <i>et al.</i> (2006) Dendritic cells expressing transgenic galectin-1 delay onset of autoimmune diabetes in mice. J Immunol. 177 (8): 5278-89. Vitadello, M. <i>et al.</i> (2010) Myofiber stress-response in myositis: parallel investigations on patients and experimental animal models of muscle regeneration and systemic inflammation. Arthritis Res Ther. 12 (2): R52. Huang, Y.C. <i>et al.</i> (2008) CD5-low expression lymphocytes in canine peripheral blood show characteristics of natural killer cells. J Leukoc Biol. 84 (6): 1501-10. Liu, C.C. <i>et al.</i> (2008) Transient downregulation of monocyte-derived dendritic-cell differentiation, function, and survival during tumoral progression and regression in an <i>in vivo</i> canine model of transmissible venereal tumor. Cancer Immunol Immunother. 57 (4): 479-91. Letellier, M. <i>et al.</i> (2008) Normal adult climbing fiber mono-innervation of cerebellar Purkinje cells in mice lacking MHC class I molecules. Dev Neurobiol. 68 (8): 997-1006. Gupta, A. <i>et al.</i> (2012) Efficacy of <i>Mycobacterium indicus pranii</i> immunotherapy as an adjunct to chemotherapy for tuberculosis and underlying immune responses in the lung. PLoS One. 7 (7): e39215. Giunchetti, R.C. <i>et al.</i> (2007) Immunogenicity of a killed <i>Leishmania</i> vaccine with saponin adjuvant in dogs. Vaccine. 25 (44): 7674-86. Cenci, E. <i>et al.</i> (2006) Modulation of phenotype and function of dendritic cells by a therapeutic synthetic killer peptide. J Leukoc Biol. 79 (1): 40-5.

9. Giunchetti RC *et al.* (2008) A killed *Leishmania* vaccine with sand fly saliva extract and saponin adjuvant displays immunogenicity in dogs. [Vaccine. 26 \(5\): 623-38.](#)
10. Patel, G.K. *et al.* (2012) A humanized stromal bed is required for engraftment of isolated human primary squamous cell carcinoma cells in immunocompromised mice. [J Invest Dermatol. 132 \(2\): 284-90.](#)
11. Zuza, A.L. *et al.* (2016) Astrocyte response to St. Louis encephalitis virus. [Virus Res. 217: 92-100.](#)
12. Lohan, P. *et al.* (2016) Culture expanded primary chondrocytes have potent immunomodulatory properties and do not induce an allogeneic immune response. [Osteoarthritis Cartilage. 24 \(3\): 521-33.](#)
13. Gupta, A. *et al.* (2012) Protective efficacy of *Mycobacterium indicus pranii* against tuberculosis and underlying local lung immune responses in guinea pig model. [Vaccine. 30 \(43\): 6198-209.](#)
14. Reid E *et al.* (2016) Type I and III IFNs Produced by Plasmacytoid Dendritic Cells in Response to a Member of the Flaviviridae Suppress Cellular Immune Responses. [J Immunol. 196 \(10\): 4214-26.](#)
15. Iwasaki, Y. *et al.* (2016) Differentiation/Purification Protocol for Retinal Pigment Epithelium from Mouse Induced Pluripotent Stem Cells as a Research Tool. [PLoS One. 11 \(7\): e0158282.](#)
16. Wang, Y. *et al.* (2020) Characterization of a rhodanese homologue from *Haemonchus contortus* and its immune-modulatory effects on goat immune cells *in vitro*. [Parasit Vectors. 13 \(1\): 454.](#)
17. Ehsan, M. *et al.* (2021) *Fasciola gigantica* tegumental calcium-binding EF-hand protein 4 exerts immunomodulatory effects on goat monocytes. [Parasit Vectors. 14 \(1\): 276.](#)

Further Reading

1. Piriou-Guzylack, L. (2008) Membrane markers of the immune cells in swine: an update. [Vet Res. 39: 54.](#)

Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee

12 months from date of despatch

Health And Safety Information

Material Safety Datasheet documentation #10040 available at: 10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

Regulatory

For research purposes only

Related Products

Recommended Secondary Antibodies

- | | |
|-----------------------------------|---------------------|
| Goat Anti Mouse IgG (STAR77...) | HRP |
| Rabbit Anti Mouse IgG (STAR12...) | RPE |

Goat Anti Mouse IgG (STAR70...) [FITC](#)
Goat Anti Mouse IgG IgA IgM (STAR87...) [Alk. Phos.](#), [HRP](#)
Rabbit Anti Mouse IgG (STAR9...) [FITC](#)
Goat Anti Mouse IgG (STAR76...) [RPE](#)
Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight®488](#), [DyLight®550](#),
[DyLight®650](#), [DyLight®680](#), [DyLight®800](#),
[FITC](#), [HRP](#)
Rabbit Anti Mouse IgG (STAR13...) [HRP](#)
Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M413236:221121'

Printed on 21 Nov 2022

© 2022 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)